Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	29 (1983)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	SOME KNOT THEORY OF COMPLEX PLANE CURVES
Autor:	Rudolph, Lee
Kapitel:	§6. Global knot theory in brief—the affine case
DOI:	https://doi.org/10.5169/seals-52979

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. <u>Siehe Rechtliche Hinweise.</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. <u>Voir Informations légales.</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. <u>See Legal notice.</u>

Download PDF: 25.05.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

= 1). The rest of the relations needed for the van Kampen presentation of $\pi_1(\mathbb{CP}^2 - \Gamma; *)$ come, then, by declaring this representation trivial. One obtains a finite presentation, of course, by choosing generators of the acting free group; Moishezon's problem of "normal forms" is essentially the problem of making a good choice. Several modernizations [Abe], [Che], [Cha] of van Kampen's proof have been published in recent years.

In a standard van Kampen presentation (where the generators of the acting free group are free generators), each relation corresponds either to a singularity of Γ or to a simple vertical tangent to Γ ; and (up to the action of the corresponding free generator) each relation is of a certain canonical form, which depends only on the closed braid type (§7) of the link of the branch(es) at the point of Γ , through which the line in the pencil passes that gives the relation in question, where this line itself is used to find the axis of the closed braid. In particular, the knot group of a node curve always has a standard van Kampen presentation in which each relation either sets conjugates of two x_i equal (from a simple vertical tangent) or says that two such conjugates commute (from a node); if "conjugates" could be deleted, the Zariski Conjecture would be trivially true.

There is also a great body of work on "knot groups" of curves in (compact, smooth) complex surfaces other than \mathbb{CP}^2 , and on the related issue of fundamental groups of surfaces; we cannot touch on these topics here.

§6. GLOBAL KNOT THEORY IN BRIEF—THE AFFINE CASE

Little appears to be known about algebraic curves in affine space, from the knot-theoretical viewpoint. The gross algebraic topology (even just homology theory) of \mathbb{CP}^2 is implicated with the quite rigid geometry; but affine space is contractible, and on the other hand its geometry is "infinite" (for instance in the sense that there are Lie groups of arbitrarily high dimension contained in the group of birgular automorphisms of \mathbb{C}^2), so that the conspirators have fallen out and neither can give away much about the other.

One might think, for example, to study the embedding of a curve Γ in \mathbb{C}^2 by first embedding \mathbb{C}^2 itself into \mathbb{CP}^2 . Then the affine complement $\mathbb{C}^2 - \Gamma$ becomes the projective complement $\mathbb{CP}^2 - (\Gamma \cup \mathbb{CP}^1_{\infty})$, where $\Gamma \cup \mathbb{CP}^1_{\infty}$ is a (reducible) projective algebraic curve. The obstacle to this program is the unfortunate fact that \mathbb{C}^2 , just as an algebraic surface, without distinguished coordinates, is not uniquely embedded as $\mathbb{CP}^2 - \mathbb{CP}^1_{\infty}$. Any biregular automorphism of \mathbb{C}^2 (in particular, one of the vast majority which cannot be extended biregularly to \mathbb{CP}^2) will move Γ around, and so the configuration of $\Gamma \cup \mathbb{CP}^1_{\infty}$ is not determined by the embedding of Γ in \mathbb{C}^2 . (For instance, though the geometric number of points at infinity on Γ is determined by Γ , the algebraic intersection number of the closure of Γ with the line at infinity can be made arbitrarily large. Likewise the local singularities at infinity are not determined by the affine curve.)

The main theorems known here have been proved by Abhyankar and his collaborators [A-M, A-S]. They are unknotting theorems, in the sense that they take this form: "Let Γ be a certain curve in \mathbb{C}^2 , and let $i: \Gamma \to \mathbb{C}^2$ be any algebraic embedding; then there is a biregular automorphism of \mathbb{C}^2 returning *i* to the inclusion map". Briefly, such a curve Γ cannot be knotted in \mathbb{C}^2 .

However, for most of the curves they deal with, these theorems are not genuinely topological, for the reimbedding *i* is required to be an embedding of Γ with its given structure as a variety, and generally there might be moduli. Only in the original theorem [A-M] (which had been stated, but not correctly proved, by Segre) are there no conceivable moduli, when Γ is a straight line. Then the theorem is this.

THEOREM. Let $\Gamma \subset \mathbb{C}^2$ be an algebraic curve without singularities, homeomorphic to \mathbb{C} . Then there is a biregular change of coordinates $A : \mathbb{C}^2 \to \mathbb{C}^2$ so that $A\Gamma$ is a straight (complex) line.

A topological proof has been given in [Ru 4]. It goes like this. One shows (just as for a singular point) that the intersection of Γ (which we can assume to be parametrized by z = p(t), w = q(t), $p, q \in \mathbb{C}[t]$) with a very large bidisk boundary is an iterated torus knot $K = O\{m_1, n_1; ...; m_s, n_s\}$, with m_1 $= m/GCD(m, n), n_1 = n/GCD(m, n), m = \deg p, n = \deg q$. By hypothesis, K is a slice knot. This forces K = O, in particular, one of m_1, n_1 is 1. Thereafter the argument is as in [A-M]—if (say) $m_1 = 1$ and p and q are monic, then the biregular change of coordinates $(z, w) \mapsto (z, w - z^{m/n})$ carries Γ to another curve satisfying the hypotheses, of lower bidegree; and so we proceed until one of z, w is linear and the other constant.

As to analytic curves in affine space, almost nothing is known. The obvious analogue of the Theorem above is definitely false: for it is known that the unit disk in C can be properly analytically embedded in C^2 [H]; since the disk and the line are analytically inequivalent, no analytic change of coordinates in C^2 could unknot the disk to a line. It is, however, perfectly possible that every such disk is smoothly unknotted. Presently I am unable even to prove that an analytic line in C^2 is smoothly unknotted.