Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 29 (1983)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE TOPOLOGY OF REAL ALGEBRAIC SETS

Autor: Akbulut, Selman / King, Henry

Kapitel: §3. Blowing Down

DOI: https://doi.org/10.5169/seals-52981

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation


L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 16.10.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch



 $\mathcal{S}_{Alg}(V)$ is the set of distinct algebraic structures on V. Hence a natural problem is to compute $\mathcal{S}_{Alg}(V)$, or at least produce nontrivial elements of this set. For example if we take $M \subset V$ as in Proposition 2.10, then by Theorem 2.12 (V, M) is diffeomorphic to nonsingular algebraic sets (V', M'). Let |V| = |V'| denote the underlying smooth structures and let $V \stackrel{g}{\to} |V|$, $V' \stackrel{g'}{\to} |V|$ be the forgetful maps. Then (V, g) and (V', g') are distinct elements of $\mathcal{S}_{Alg}(|V|)$, otherwise M would be isotopic to a nonsingular algebraic subset of V.

An interesting question is whether algebraic structures on smooth manifolds satisfy the product structure theorem; that is, whether the natural map

$$\mathscr{S}_{Alg}(M) \times \mathbf{R}^n \to \mathscr{S}_{Alg}(M \times \mathbf{R}^n), \ (V, g) \mapsto (V \times \mathbf{R}^n, g \times id)$$

is surjection. The answer would be negative if one can find a smooth manifold M and $\theta \in H_*(M; \mathbb{Z}/2\mathbb{Z})$ such that M can not be diffeomorphic to a nonsingular algebraic set M' with $\theta \in H_*^A(M'; \mathbb{Z}/2\mathbb{Z})$. To see this, pick any smooth representative $N \xrightarrow{g} M$ of $\theta = g_*[N]$. By graphing g, we can assume $N \subset M \times \mathbb{R}^n$ for some n and g is induced by projection. By Theorem 2.12 we can find a diffeomorphism $\lambda : M \times \mathbb{R}^n \to V$ to a nonsingular algebraic set V with $\lambda(N)$ is an algebraic subset (one has to modify Theorem 2.12 to apply to this noncompact case). Then there can not exist a birational diffeomorphism $\mu : V \to M' \times \mathbb{R}^n$ where M' is a nonsingular algebraic set diffeomorphic to M, otherwise $\lambda(N) \xrightarrow{\mu} M' \times \mathbb{R}^n$ would represent $\theta \in H_*^A(M'; \mathbb{Z}/2\mathbb{Z})$.

§3. Blowing Down

Real algebraic sets obey some simple but useful topological properties: Proposition 3.1.

- (a) One point compactification an algebraic set is homeomorphic to an algebraic set.
- (b) Given algebraic sets $L \subset V$, then V L is homeomorphic to an algebraic set.
- (c) Given algebraic sets $L \subset V$ with V compact then V/L is homeomorphic to an algebraic set.

Proof:

- (a) Let $Z \subset \mathbb{R}^n$ be an algebraic set and assume that $Z \neq \mathbb{R}^n$ and $0 \notin Z$ (otherwise translate Z). Let $Z = f^{-1}(0)$ for some polynomial f(x); then define $F(x) = |x|^{2d} f\left(\frac{x}{|x|^2}\right)$, where d is the degree of f(x). Clearly F(x) is a polynomial and $F^{-1}(0)$ is the one point compactification of Z, since $x \mapsto \frac{x}{|x|^2}$ is the inversion through the unit sphere.
- (b) Let $V = f^{-1}(0)$, $L = g^{-1}(0)$ for some polynomials $f, g : \mathbb{R}^n \to \mathbb{R}$. Define $G(x, t) = |f(x)|^2 + |tg(x) 1|^2$, then $G^{-1}(0) \approx V L$.
- (c) By applying (a) we get the one point compactification of $G^{-1}(0)$ to be an algebraic set; if V is compact this set is homeomorphic to V/L.

This proposition implies that a set is homeomorphic to an algebraic set if and only if the one point compactification is homeomorphic to an algebraic set. Hence any noncompact algebraic set has a collar at infinity, since every algebraic set is locally cone-like [M]. Also we get that the reduced suspension $\Sigma^n X = X \times S^n/X \vee S^n$ of any algebraic set X is homeomorphic to an algebraic set.

There is a fancier version of the blowing down operation (c) (Proposition 3.3). First we need to discuss projectively closed algebraic sets. Let $p: \mathbf{R}^n \to \mathbf{R}$ be a polynomial. Another interpretation of this concept is the following: Let $\lambda: \mathbf{R}^n$ d. We call p(x) an overt polynomial if $p_d^{-1}(0)$ is either the empty set or $\{0\}$. We call an algebraic set $V = p^{-1}(0)$ a projectively closed algebraic set if p(x) is an overt polynomial. Another interpretation of this concept is the following: Let $\lambda: \mathbf{R}^n \to \mathbf{R}\mathbf{P}^n$ be the inclusion $\lambda(x_1, ..., x_n) = [1; x_1; ...; x_n]$ then $V = p^{-1}(0)$ is projectively closed if and only if λ is a projective algebraic subset of $\mathbf{R}\mathbf{P}^n$ in other words $\lambda(V)$ is Zariski closed in $\mathbf{R}\mathbf{P}^n$ (see also $[AK_2]$). Real algebraic sets along with maps can easily be made projectively closed by the following.

PROPOSITION 3.2. Let $f: Z \to W$ be an entire rational function between algebraic sets with Z nonsingular and compact. Then there is a projectively closed algebraic set $V \subset W \times \mathbb{R}^n$ a birational diffeomorphism g which makes the following commute

$$V \hookrightarrow W \times \mathbf{R}^n$$
 $g \uparrow \approx \downarrow^{\pi}$
 $Z \stackrel{f}{\rightarrow} W$

where π is the projection, n is some integer.

Proof: By taking the graph of f we can assume that $Z \subset W \times \mathbf{R}^m \subset \mathbf{R}^r$ for some r, and f is induced by projection. Also identify $\mathbf{R}^r \subset \mathbf{RP}^r$ via λ . Then let \overline{Z} be the Zariski closure of Z in \mathbf{RP}^r . We claim $\dim(\overline{Z}-Z) < \dim(Z)$. This is because if U is an irreducible component of \overline{Z} then $U \cap Z \neq \emptyset$, and therefore $U - Z = U \cap \mathbf{RP}^{r-1}$ is a proper algebraic subset of U where $\mathbf{RP}^{r-1} = \{[0; x_1; ...; x_r] \in \mathbf{RP}^r\}$. Since U is irreducible $\dim(U - Z) < \dim(U)$, also $\dim(U) = \dim(Z)$. Therefore $\dim(\overline{Z}-Z) < \dim(Z)$. So $\overline{Z} - Z = \operatorname{Sing}(\overline{Z})$. By resolution of singularities [H] (Theorem 1.1) there is a nonsingular algebraic set $V \subset \mathbf{RP}^r \times \prod_i \mathbf{RP}^{a_i}$ such that the projection induces birational diffeomorphism between V and Z. In particular $V \subset \mathbf{R}^r \times \prod_i \mathbf{RP}^{a_i}$.


$$\mathbf{RP}^r \times \prod_i \mathbf{RP}^{a_i} \hookrightarrow \mathbf{R}^{(r+1)^2 + \Sigma(a_i+1)^2}$$


is a projectively closed algebraic set. Hence V is projectively closed (check details).

Now assume that $L \subset W \subset \mathbf{R}^m$ be real algebraic sets, and $V \subset W \times \mathbf{R}^n$ be a projectively closed algebraic set. Let $q : \mathbf{R}^m \to \mathbf{R}$ be a polynomial with $q^{-1}(0) = L$. Define

$$D_q: W \times \mathbf{R}^n \to W \times \mathbf{R}^n$$

by $D_q(x, y) = (x, yq(x))$. D_q is a diffeomorphism on $(W-L) \times \mathbb{R}^n$ and $D_q(L \times \mathbb{R}^n) = L \times 0$. Therefore $D_q(V)$ is the quotient space of V by the equivalence relation $(x, y) \sim (x, 0)$ if $x \in L$. We call the operation $V \to D_q(V) \cup L$ (L is identified by $L \times 0$) blowing down V over L.





PROPOSITION 3.3. Given L, W, V as above, then $D_q(V) \cup L$ is an algebraic subset of $W \times \mathbb{R}^n$.

Proof: Let $p: \mathbb{R}^m \times \mathbb{R}^n \to \mathbb{R}$ be an overt polynomial of degree e with $V = p^{-1}(0)$ and let q be as above. Define a polynomial $r: \mathbb{R}^m \times \mathbb{R}^n \to \mathbb{R}$ by

$$r(x, y) = q(x)^e p\left(x, \frac{y}{q(x)}\right)$$

We claim $r^{-1}(0) = D_a(V) \cup L$. It is easy to see that

$$r^{-1}(0) \cap (W-L) \times \mathbf{R}^n = D_q(V) \cap (W-L) \times \mathbf{R}^n$$
,

so it suffices to show that $r^{-1}(0) \cap (L \times \mathbf{R}^n) = L \times 0$. We decompose $p(x, y) = p_e(x, y) + \alpha(x, y)$ where $p_e(x, y)$ is homogeneous of degree e and $\alpha(x, y)$ is a polynomial of degree less than e. Hence if $(x, y) \in r^{-1}(0) \cap (L \times \mathbf{R}^n)$ then r(x, y) = 0 and q(x) = 0, which implies $r(x, y) = p_e(0, y) = 0$. Then y = 0 since p is overt, so $(x, y) \in L \times 0$. Conversely if $(x, y) \in L \times 0$ then y = 0 and q(x) = 0. Hence $r(x, y) = p_e(0, 0) = 0$, i.e. $(x, y) \in r^{-1}(0) \cap (L \times \mathbf{R}^n)$.

There is a more useful version of Proposition 3.3 which says that after modifying D_q we can get $D_q(V) \cup L$ to be a projectively closed algebraic set (Proposition 3.1 of $[AK_6]$). This allows us to iterate this blowing down process.

§4. ISOLATED SINGULARITIES

The topology of real algebraic sets with isolated singularities is completely understood by the following Theorem.

Theorem 4.1 ([AK₂]). X is homeomorphic to an algebraic set with isolated singularities if and only if X is obtained by taking a smooth compact manifold W with boundary $\partial W = \bigcup_{i=1}^{r} M_i$, where each M_i bounds, then crushing some M_i 's to points and deleting the remaining M_i 's.

