

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 30 (1984)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: PARTITIONS INTO SQUARES
Autor: Grosswald, Emil
Kurzfassung
DOI: <https://doi.org/10.5169/seals-53828>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

PARTITIONS INTO SQUARES

by Emil GROSSWALD

ABSTRACT¹⁾. In following D. H. Lehmer (*Amer. Math. Monthly* 55 (1948), 476-481), the notation $P_k(n)$ stands for the number of partitions of the integer n into sums of k squares of positive integers in, say, non-decreasing order. D. H. Lehmer has determined the sets $S_{k,1}$ of integers n with $P_k(n) = 1$. Here the sets $S_{k,m}$ of integers satisfying $P_k(n) = m$ are determined. Next, the sets $S_{k,m}(x) \subset S_{k,m}$, with $n \leq x$ are described and exact or asymptotic formulae are obtained for their size.

1. INTRODUCTION. In 1948, D. H. Lehmer [8] studied the solutions of the equation $P_k(n) = 1$, where $P_k(n)$ is the number of *partitions* of the integer n into a sum of k squares of non-negative, non-decreasing integers. The number $P_k(n)$ of partitions differs from the number $r_k(n)$ of *representations* of n by k squares in that, for $r_k(n)$, we count representations as distinct if they differ by the order of the summands and the summands are squares of arbitrary (also negative) integers. If all summands of a partition counted by $P_k(n)$ are distinct and non-zero, then to it correspond $2^k k!$ representations counted by $r_k(n)$. If any one of the summands of a partition vanishes, or if not all are distinct, then to that partition correspond fewer than $2^k k!$ representations, but in any case, $P_k(n) \geq r_k(n)/2^k k!$.

In [8] Lehmer essentially solved the problem quoted above, except for $k = 3$. A contribution to the latter case was recently made by Bateman and the author (see [2]), but the problem is still not completely solved.

The purpose of the present paper is to discuss the more general question of the sets

$$S_{k,m} = \{n \mid P_k(n) = m\} \text{ and } S_{k,m}(x) = \{n \mid n \in S_{k,m}, n \leq x\},$$

as well as $|S_{k,m}(x)|$, the number of elements of $S_{k,m}(x)$.

¹⁾ Subject classification: 10 J 05.