

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 31 (1985)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: FLAT MANIFOLDS WITH Z/p^2 HOLOMOMY
Autor: Hiller, Howard
Kapitel: §2. Cohomology, restrictions and special classes
DOI: <https://doi.org/10.5169/seals-54570>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 02.09.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The splintering of these genera into isomorphism classes has been analyzed by Reiner [13]. One can, of course, replace R_1, R_2 by ideal classes in these rings and Λ_1 by $E(\alpha)$ (cf. section 1) where α is an ideal class in R_1 . There is an additional invariant lying in a quotient of the group of units of a certain finite ring and, if $p \equiv 1 \pmod{4}$ a certain quadratic residue character mod p can also appear as an invariant. The precise result is Theorem 7.3 of [13]. We will require only the observation [13, p. 494] that if $p = 2, 3$ there are no further invariants, i.e. each genus of an indecomposable is a single isomorphism class. In the case $p = 5$ already, although the class number of $\mathbf{Q}(e^{2\pi i/m})$ is one for $m = 5, 25$, the 21 genera of indecomposables split up into 40 isomorphism classes. Hence already the further isomorphism invariants mentioned above exert an influence.

§ 2. COHOMOLOGY, RESTRICTIONS AND SPECIAL CLASSES

If H is a finite group, M an H -lattice then: $H^i(H, M) \cong \bigoplus_p H^i(H, M_p)$, where p ranges over the primes dividing the order of H [3, p. 84]. Hence if M and M' are locally isomorphic, $H^i(H, M) \cong H^i(H, M')$; so the cohomology of an H -lattice depends only on its genus.

We recall the cohomology of a cyclic group $\mathbf{Z}/n = \langle \sigma \rangle$ [3, p. 58]. We write $N = 1 + \sigma + \dots + \sigma^{n-1}$ and $D = 1 - \sigma$. If M is a \mathbf{Z}/n -module, then

$$\begin{aligned} H^0(\mathbf{Z}/n, M) &= M^\sigma \\ H^{2i-1}(\mathbf{Z}/n, M) &= {}_N M / D \cdot M \\ H^{2i}(\mathbf{Z}/n, M) &= M^\sigma / N \cdot M \end{aligned}$$

for all $i \geq 1$, where M^σ denotes σ -invariants and ${}_N M = \{x \in M : Nx = 0\}$. From these remarks it is easy to compute the cohomology of the indecomposable \mathbf{Z}/p -lattices described in section 1.

(2.1) PROPOSITION. *The following table describes the cohomology of the indecomposable \mathbf{Z}/p -lattices:*

M	rank	H^0	H^1	H^2
1	1	\mathbf{Z}	0	\mathbf{Z}/p
α	$p - 1$	0	\mathbf{Z}/p	0
β	p	\mathbf{Z}	0	0

Similarly one can easily compute the cohomology and restriction of the first four \mathbf{Z}/p^2 -lattices of (1.3).

(2.2) PROPOSITION. *If $[M]_p$ denotes the restriction of the \mathbf{Z}/p^2 -lattice M to the subgroup of order p , we have the following table.*

M	rank	H^0	H^1	H^2	$[M]_p$
$M_1 = \mathbf{1}$	1	\mathbf{Z}	0	\mathbf{Z}/p^2	$\mathbf{1}$
$M_2 = R_1$	$p - 1$	0	\mathbf{Z}/p	0	$(p-1)\mathbf{1}$
$M_3 = R_2$	$p^2 - p$	0	\mathbf{Z}/p	0	$p\alpha$
$M_4 = \Lambda_1$	p	\mathbf{Z}	0	\mathbf{Z}/p	$p\mathbf{1}$

Furthermore, $\Lambda_2 = \mathbf{Z}[\mathbf{Z}/p^2]$, the regular representation of \mathbf{Z}/p^2 , satisfies $H^0 = \mathbf{Z}$, $H^1 = 0 = H^2$ and $[\Lambda_2]_\beta = p\beta$.

Proof. It suffices to observe that $M_2 = p^*(\alpha)$ and $M_4 = p^*(\beta)$, where $p: \mathbf{Z}/p^2 \rightarrow \mathbf{Z}/p$ is the natural projection, and M_3 fits into a short exact sequence:

$$0 \rightarrow M_4 \rightarrow \Lambda_2 \rightarrow M_3 \rightarrow 0.$$

The last remark follows from the freeness of Λ_2 .

To complete the table for the modules M_i , $i \geq 5$, we have the following lemma:

(2.3) LEMMA. *If L is a \mathbf{Z}/p -lattice, $\alpha \in L$, then the extension $M = (L, \alpha)$ defined by (1.2) satisfies:*

$$H^2(C; M) \cong \text{coker}(x_*: H^2(C; \varphi_2 \Lambda_2) \rightarrow H^2(C; L))$$

where C is either \mathbf{Z}/p^2 or \mathbf{Z}/p .

Proof. The diagram (1.2) induces

$$\begin{array}{ccccccc} \rightarrow & H^1(C, R_2) & \xrightarrow{\delta} & H^2(C, \varphi_2 \Lambda_2) & \rightarrow & 0 \\ & \downarrow & & \downarrow x_* & & & \\ \rightarrow & H^1(C, R_2) & \xrightarrow{\delta} & H^2(C, L) & \rightarrow & H^2(C, M) & \rightarrow 0 \end{array}$$

where the zeros follow from (2.2). An easy diagram-chase completes the proof.

(2.4) PROPOSITION. *The following table describes the cohomology of the indecomposable \mathbf{Z}/p^2 -lattices M_i , $i \geq 5$:*

M	rank	H^0	H^1	H^2
M_5	$p^2 - p + 1$	\mathbf{Z}	0	\mathbf{Z}/p
$M_6(0)$	p^2	\mathbf{Z}	0	0
$M_6(k)$	p^2	\mathbf{Z}	\mathbf{Z}/p	\mathbf{Z}/p
$M_7(k)$	$p^2 + 1$	$\mathbf{Z} \oplus \mathbf{Z}$	0	$\mathbf{Z}/p \oplus \mathbf{Z}/p$
$M_8(0)$	$p^2 - 1$	0	\mathbf{Z}/p^2	0
$M_8(k)$	$p^2 - 1$	0	$\mathbf{Z}/p \oplus \mathbf{Z}/p$	0
$M_9(k)$	p^2	\mathbf{Z}	\mathbf{Z}/p	\mathbf{Z}/p

Proof. Since

$$H^0(\mathbf{Z}/p^2; R_2) = 0, H^0(\mathbf{Z}/p^2; (L, x)) \cong H^0(\mathbf{Z}/p^2; L)$$

and these can be read off from (2.2). The groups H^2 are computed by (2.3). We work out one example in detail. Consider $M_6(k)$, $0 \leq k \leq p-1$, so that $L = \Lambda_1$. If we identify $\varphi_2 \Lambda_2$ with Λ_1 then the generator:

$$1 + x + \dots + x^{p-1} \in H^2(\mathbf{Z}/p^2, \varphi_2 \Lambda_2)$$

is sent by λ_*^k , $1 \leq k \leq p-1$, to

$$(1-x)^k (1+x+\dots+x^{p-1}) = (1-x)^{k-1} \cdot 0 = 0$$

in $H^2(\mathbf{Z}/p^2; \Lambda_1)$. If $k=0$, then the map is an isomorphism. Hence $H^2(\mathbf{Z}/p^2; M_6(0)) = 0$ and $H^2(\mathbf{Z}/p^2, M_6(k)) = \mathbf{Z}/p$, $k \geq 1$.

The groups $H^1(M_i)$ can be read off the long exact cohomology sequence of the bottom row of (1.2).

Remark. It follows from (2.2) that $M_6(0)$ is the genus of the regular representation.

We now record the restrictions of the modules M_i to the subgroup of order p .

(2.5) PROPOSITION. *The \mathbf{Z}/p -cohomology and the restrictions of M_i , $i \geq 5$, are given by:*

M	$H^2(\mathbf{Z}/p; [M]_p)$	$[M]_p$
M_5	0	$(p-1)\alpha + \beta$
$M_6(k)$	$k(\mathbf{Z}/p)$	$k\mathbf{1} + k\alpha + (p-k)\beta$
$M_7(k)$	$(k+1)(\mathbf{Z}/p)$	$(k+1)\mathbf{1} + k\alpha + (p-k)\beta$
$M_8(k)$	$k(\mathbf{Z}/p)$	$k\mathbf{1} + (k+1)\alpha + (p-k-1)\beta$
$M_9(k)$	$(k+1)(\mathbf{Z}/p)$	$(k+1)\mathbf{1} + (k+1)\alpha + (p-k-1)\beta$

Proof. One begins by computing $H^2(\mathbf{Z}/p, [M]_p)$, from (2.3). We work out an example again with $M = M_6(k)$. We will need these details later. The map

$$p(\mathbf{Z}/p) = H^2(\mathbf{Z}/p, \Lambda_1) \xrightarrow{\lambda_*^k} H^2(\mathbf{Z}/p; \Lambda_1) = p(\mathbf{Z}/p)$$

sends the generator x^j , $0 \leq j \leq p-1$, from the left-hand side to $x^j(1-x)^k$. The resulting matrix $C_{p,k}$ in $GL_p(\mathbf{Z}/p)$ can be described in the following way. If $p > k$, let $C_{p,k,j}$ denote a column p -vector whose entries are the coefficients of $(1-x)^k$ introduced “cyclically” starting in row j . For example:

$$C_{5,2,4} = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \\ 1 \\ -2 \end{bmatrix} \quad \text{We define } C_{p,k} \text{ to be the } p \times p$$

matrix whose j^{th} column is $C_{p,k,j}$. So, for example,

$$C_{5,2} = \begin{bmatrix} 1 & 0 & 0 & 1 & -2 \\ -2 & 1 & 0 & 0 & 1 \\ 1 & -2 & 1 & 0 & 0 \\ 0 & 1 & -2 & 1 & 0 \\ 0 & 0 & 1 & -2 & 1 \end{bmatrix}$$

It is a consequence of the identity $(1-x)^{k+1} = (1-x)(1-x)^k$ that

$$C_{p,k+1} = C_{p,1} C_{p,k}$$

so we get:

$$C_{p,k} = C_{p,1}^k = \begin{bmatrix} 1 & & & & -1 \\ -1 & 1 & & & 0 \\ & & -1 & 1 & \\ & & & & . \\ 0 & & & -1 & 1 \end{bmatrix}^k.$$

It is easy to see that $C_{p,1}(p(\mathbf{Z}/p)) = \{\bar{x} \in p(\mathbf{Z}/p) : \sum_{i=0}^{p-1} x_i = 0\}$. Hence $\text{rank } (C_{p,k}) = p - k$ and $\dim_{\mathbf{Z}/p} H^2(\mathbf{Z}/p; M_6(k)) = p - (p - k) = k$, as claimed. The other cases are similar.

Now from (2.1) we see that the \mathbf{Z}/p -dimension of H^2 is the multiplicity of $\mathbf{1}$ in $[M]_p$. The multiplicities of α and β are then determined by the bottom row of (1.2) restricted to the subgroup of order p .

Recall from Charlap [4] that a cohomology class $\alpha \in H^2(H, M)$ is called *special* if $i^*(\alpha) \neq 0$ for the inclusion $i: C \rightarrow H$ of any cyclic subgroup. The basic result is (see [4, p. 22]).

(2.6) PROPOSITION. *The extension Γ in (0, 1) corresponding to $\alpha \in H^2(H, M)$ is torsion-free (i.e. the fundamental group of a flat manifold) if and only if α is special.*

It remains to determine which indecomposables in (1.3) admit special classes. The result is:

(2.7) PROPOSITION. *There are $2p - 1$ genera of indecomposable \mathbf{Z}/p^2 -lattices that admit special classes. They are $M_1, M_4, M_7(k), 1 \leq k \leq p - 2$, and $M_9(k), 0 \leq k \leq p - 2$.*

Proof. From (2.2), (2.4) and (2.5) one sees that the given lattices along with $M_6(k), 1 \leq k \leq p - 2$, are the only possibilities. We must determine the restriction map

$$i^*(M): H^2(\mathbf{Z}/p^2; M) \rightarrow H^2(\mathbf{Z}/p; M)$$

in these cases. Clearly for $M = M_1$, $i^*(M)$ is the natural projection and for $M = M_4$, $i^*(M)$ is the diagonal embedding.

Now suppose $M = M_6(k)$. We have a commutative diagram of exact sequences from (1.2) and (2.3)

$$\begin{array}{ccccccc}
 H^2(\mathbf{Z}/p^2; \Lambda_1) & \rightarrow & H^2(\mathbf{Z}/p^2; M) & \rightarrow & 0 \\
 \downarrow \Delta = i*(\Lambda_1) & & \downarrow i*(M) & & \\
 H^2(\mathbf{Z}/p; \varphi_2 \Lambda_2) & \xrightarrow{\lambda_*^k} & H^2(\mathbf{Z}/p; \Lambda_1) & \rightarrow & H^2(\mathbf{Z}/p; M) & \rightarrow & 0
 \end{array}$$

where Δ is the diagonal map $\mathbf{Z}_p \rightarrow p(\mathbf{Z}/p)$. Hence to eliminate $M_6(k)$, it suffices to show $\text{Im}(\Delta) \subset \text{Im}(\lambda_*^k)$. Let e denote a column p -vector consisting of all 1's, according to the proof of (2.5) we must find an \bar{x}_k , $1 \leq k \leq p-1$ so that $C_{p,k} \cdot \bar{x}_k = C_{p,1}^k \cdot \bar{x}_k = e$. We do this inductively

on k . For example, $\bar{x}_1 = \begin{bmatrix} 1 \\ 2 \\ \vdots \\ p \end{bmatrix}$, as can easily be checked. Inductively we define

$$\bar{x}_k(i) = \begin{cases} 1 & i = 1 \\ \bar{x}_{k-1}(i-1) + \bar{x}_{k-1}(i) & i > 1. \end{cases}$$

Clearly $C_{p,1} \cdot \bar{x}_k = \bar{x}_{k-1}$, for all coordinates except possibly the first; we must show $\bar{x}_k(p) \equiv 0 \pmod{p}$. But a comparison of the \bar{x}_k 's with Pascal's triangle convinces one that

$$\bar{x}_k(p) = \binom{p-1+k}{p-1} \equiv \binom{k-1}{p-1} \binom{1}{0} \equiv 0 \pmod{p},$$

since $k-1 < p-1$.

We leave it for the reader to check that the restriction maps for $M_7(k)$ and $M_9(k)$ are non-trivial.

§ 3. $\mathbf{Z}/4$ -MANIFOLDS

In this section, we consider the case $p = 2$. For convenience, we change the notation slightly and write M_7 for $M_6(1)$ and M_i for $M_i(0)$, $i = 6, 8, 9$. According to (2.7), the indecomposable $\mathbf{Z}/4$ -lattices that carry special classes are M_1 , M_4 and M_9 . It is easy to see M_i is faithful if and only if