

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 31 (1985)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: FLAT MANIFOLDS WITH Z/p^2 HOLOMOMY
Autor: Hiller, Howard
Kapitel: §3. $Z/4$ -MANIFOLDS
DOI: <https://doi.org/10.5169/seals-54570>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 02.09.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

$$\begin{array}{ccccccc}
 H^2(\mathbf{Z}/p^2; \Lambda_1) & \rightarrow & H^2(\mathbf{Z}/p^2; M) & \rightarrow & 0 \\
 \downarrow \Delta = i*(\Lambda_1) & & \downarrow i*(M) & & \\
 H^2(\mathbf{Z}/p; \varphi_2 \Lambda_2) & \xrightarrow{\lambda_*^k} & H^2(\mathbf{Z}/p; \Lambda_1) & \rightarrow & H^2(\mathbf{Z}/p; M) & \rightarrow & 0
 \end{array}$$

where Δ is the diagonal map $\mathbf{Z}_p \rightarrow p(\mathbf{Z}/p)$. Hence to eliminate $M_6(k)$, it suffices to show $\text{Im}(\Delta) \subset \text{Im}(\lambda_*^k)$. Let e denote a column p -vector consisting of all 1's, according to the proof of (2.5) we must find an \bar{x}_k , $1 \leq k \leq p-1$ so that $C_{p,k} \cdot \bar{x}_k = C_{p,1}^k \cdot \bar{x}_k = e$. We do this inductively

on k . For example, $\bar{x}_1 = \begin{bmatrix} 1 \\ 2 \\ \vdots \\ p \end{bmatrix}$, as can easily be checked. Inductively we define

$$\bar{x}_k(i) = \begin{cases} 1 & i = 1 \\ \bar{x}_{k-1}(i-1) + \bar{x}_{k-1}(i) & i > 1. \end{cases}$$

Clearly $C_{p,1} \cdot \bar{x}_k = \bar{x}_{k-1}$, for all coordinates except possibly the first; we must show $\bar{x}_k(p) \equiv 0 \pmod{p}$. But a comparison of the \bar{x}_k 's with Pascal's triangle convinces one that

$$\bar{x}_k(p) = \binom{p-1+k}{p-1} \equiv \binom{k-1}{p-1} \binom{1}{0} \equiv 0 \pmod{p},$$

since $k-1 < p-1$.

We leave it for the reader to check that the restriction maps for $M_7(k)$ and $M_9(k)$ are non-trivial.

§ 3. $\mathbf{Z}/4$ -MANIFOLDS

In this section, we consider the case $p = 2$. For convenience, we change the notation slightly and write M_7 for $M_6(1)$ and M_i for $M_i(0)$, $i = 6, 8, 9$. According to (2.7), the indecomposable $\mathbf{Z}/4$ -lattices that carry special classes are M_1 , M_4 and M_9 . It is easy to see M_i is faithful if and only if

$i = 3, 5, 6, 7, 8, 9$. Hence if $M = \sum m_i M_i$ is an arbitrary $\mathbf{Z}/4$ -lattice then M is a faithful representation carrying a special class if and only if the multiplicities m_i satisfy the inequalities:

$$(3.0) \quad \begin{aligned} m_1 + m_4 + m_9 &> 0 \\ m_3 + m_5 + m_6 + m_7 + m_8 + m_9 &> 0. \end{aligned}$$

Since the multiplicities are a complete set of isomorphism invariants in the case $p = 2$ (see section 1) one can use the conditions (3.0) to show:

(3.1) THEOREM. If $L_n(m)$ denotes the number of isomorphism classes of n -dimensional \mathbf{Z}/m -lattices that carry special classes, then:

$$\begin{aligned} L_n(4) = \sum_{j=2}^{n-1} \left(a_j - \left[\frac{j}{2} \right] - 1 \right) + \sum_{j=[n]_2+2}^{n-2} (a_j - a_{j-1} - 1) \\ + \sum_{j=[n]_4}^{n-4} (a_j - a_{j-2} - a_{j-4} + a_{j-6}) \end{aligned}$$

where $[k]_p$ denotes the reduction of k modulo p , $[k]$ denote the largest integer $\leq k$ and the a_j 's are given by

$$P(t) = \sum_{j=0}^{\infty} a_j t^j = \frac{1}{(1-t)(1-t^2)^2(1-t^3)^2(1-t^4)^3}$$

In particular, the number of n -dimensional $\mathbf{Z}/4$ -manifolds is at least $L_n(4)$.

Proof. If $Q(t)$ is a power series, let $\text{coef}(n, Q(t))$ denote the coefficient of t^n in $Q(t)$. The number $L_n(4)$ counts the number of ways of writing

$$n = m_1 + m_2 + 2(m_3 + m_4) + 3(m_5 + m_8) + 4(m_6 + m_7 + m_9)$$

where the m_i 's satisfy (3.0). If $m_1 > 0$ there is a contribution:

$$\sum_{m_1=1}^{n-2} \text{coef}(n-m_1, P(t)) - \left(\left[\frac{n-m_1}{2} \right] + 1 \right)$$

where $\left[\frac{n-m_1}{2} \right] + 1$ is the number of ways of expressing $n - m_1$ as a combination of 1's (M_2) and 2's (M_4) (not permitted by (3.0)). Reindexing gives the first term for $L_n(4)$.

Similarly, if $m_1 = 0, m_4 > 0$ there is a contribution:

$$\sum_{m_4} \text{coef}(t^{n-2m_4}(1-t)P(t)) - 1$$

where 1 is subtracted to omit choosing m_2 alone. Finally, if $m_1 = m_4 = 0$, we have:

$$\sum_{m_9} \text{coeff}(t^{n-4m_9}, (1-t^2)(1-t^4)P(t)).$$

The coefficients of $(1-t)P(t)$ and $(1-t^2-t^4-t^6)P(t)$ are easily expressible in terms of the a_j 's and the result follows.

Remark. In order for a \mathbf{Z}/p -lattice to carry a special class, the multiplicity of the trivial representation must be non-zero. Topologically this is reflected in the fact that a \mathbf{Z}/p -manifold fibers over a circle. This is already false for a 4-dimensional $\mathbf{Z}/4$ -manifold as the following example shows.

Example. $L_4(4) = 6$. The multiplicities of the indecomposables in these 4-dimensional $\mathbf{Z}/4$ -lattices are given by:

notation of [2]	m_1	m_2	m_3	m_4	m_5	m_6	m_7	m_8	m_9
07/02/02	1				1				
12/01/04		1							1
12/01/02	1	1	1						
07/02/01	2			1					
12/01/03				1	1				
12/01/06									1

where the first column gives the label of these “ \mathbf{Z} -classes” from the table of the four-dimensional crystallographic groups in [2]. In fact, as these tables indicate, there is precisely one $\mathbf{Z}/4$ -manifold corresponding to each $\mathbf{Z}/4$ -lattice, hence there are exactly 6 4-dimensional $\mathbf{Z}/4$ -manifolds.

Remark. Recall that if $p < 23$, the field $\mathbf{Q}(e^{2\pi i/p})$ has class number one. This fact, along with the work of Charlap [4], shows that the number of n -dimensional \mathbf{Z}/p -manifolds is exactly $L_n(p)$, $p < 23$. This number is readily computable, as Charlap [4, p. 30] remarks, and the precise formula is:

$$(3.2) \quad L_n(p) = \sum_{j=p-1}^{n-1} \left(\left[\frac{j}{p-1} \right] - \left\langle \frac{j}{p} \right\rangle + 1 \right)$$

where $\langle k \rangle$ denotes the smallest integer $\geq k$. In particular, $L_p(p) = 1$, $L_n(p) = 0$, $p > n$, and when $p = 2$

$$(3.3) \quad L_n(2) = \left(\frac{n}{2}\right)^2 + \left(\frac{n}{2}\right) - 1 + \frac{[n]_2}{4}$$

using the notation of (3.1).

One can easily construct the following table of values of $L_n(p)$:

n	2	3	4	5	6
p					
2	1	3	5	8	11
3		1	2	3	4
5				1	2

Hence 14 of the 74 4-dimensional flat manifolds have cyclic holonomy ≤ 5 . (Furthermore, 26 have holonomy the Klein 4-group.) We describe analogous facts in dimension 5 below.

We let $SH^2(H, M)$ denote the set of special classes in $H^2(H, M)$. If H is a cyclic p -group and $i: \mathbf{Z}_p \hookrightarrow H$ is the inclusion of the subgroup of order p , then

$$SH^2(H, M) = H^2(H, M) - \ker(i^*) .$$

If N (resp. Z) denotes the normalizer (resp. the centralizer) of H in $\text{Aut}(M)$, there is an exact sequence (see [15, p. 50])

$$0 \rightarrow Z \rightarrow N \rightarrow \text{Aut}(H) .$$

We conjecture:

Conjecture. If $\mathbf{Z}[e^{2\pi i/p^k}]$ is a unique factorization domain for, $1 \leq k \leq n$, then N acts transitively on $SH^2(\mathbf{Z}/p^n; M)$ for any H -lattice M .

The case $n = 1$ of the Conjecture follows from Charlap [4]. Class number tables shows that the $n = 2$ case applies to $p = 2, 3, 5$, the $n = 3$ case to 2, 3 and the $n = 4$ case to $p = 2$. This conjecture implies that the lower bound of (3.1) is exact.

We mention that the multiplicities of the indecomposables in the 5-dimensional $\mathbf{Z}/4$ -lattices that admit special classes are given by:

	m_1	m_2	m_3	m_4	m_5	m_6	m_7	m_8	m_9
*	1								1
*		1					1		
*			1	1			1		
*			1	1				1	
*			1		1				
*			1		2				
*			1	2	1				
*			3					1	
*			2			1			
*			2	1	1				
*			3		1				
*				1	1				
*						1	1		
*							1		
*								1	
*			1						1

Those lattices that are starred clearly satisfy the Conjecture.

REFERENCES

- [1] BERMAN, S. and P. GUDIKOV. Indecomposable representations of finite groups over the ring of p -adic integers. *Amer. Math. Soc. Trans., Series 2*, vol. 50 (1966), 77-113.
- [2] BROWN, H., R. BÜLOW, J. NEUBÜSER, H. WONDRACTSCHEK and H. ZASSENHAUS. *Crystallographic groups of four-dimensional space*. John Wiley, New York, 1978.
- [3] BROWN, K. *Cohomology of Groups*. Springer Verlag, New York, 1983.
- [4] CHARLAP, L. Compact flat Riemannian manifolds I. *Ann. Math.* 81 (1965), 15-30.
- [5] CURTIS, C. W. and I. REINER. *Representation theory of finite groups and associative algebras*. Interscience, New York, 1966.
- [6] DIEDERICHSSEN, F. Über die Ausreduktion ganzzahliger Gruppen darstellungen bei arithmetischer Äquivalenz. *Abh. Hans. Univ.* 13 (1938), 357-412.
- [7] HELLER, A. and I. REINER. Representations of cyclic groups in rings of integers, I, II. *Ann. Math.* (2) 76 (1962), 73-92 and 77 (1963), 318-328.
- [8] HILLER, H. Minimal dimension of flat manifolds with abelian holonomy. *To appear*.