

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	31 (1985)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
 Artikel:	REPRESENTING $\Psi_2(p)$ ON A RIEMANN SURFACE OF LEAST GENUS
Autor:	Glover, Henry / Sjerve, Denis
Kapitel:	§1. Introduction
DOI:	https://doi.org/10.5169/seals-54572

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 25.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

REPRESENTING $PSl_2(p)$ ON A RIEMANN SURFACE OF LEAST GENUS

by Henry GLOVER and Denis SJERVE¹⁾

§ 1. INTRODUCTION

Given any finite group G there exists a closed Riemann surface S and an effective action $G \times S \rightarrow S$ by conformal automorphisms (here conformal means analytic). Therefore it makes sense to ask what is the least genus of such surfaces S . Recall that when the answer is that the genus equals zero (i.e. G acts on the two sphere) then G is from the list \mathbf{Z}/n , D_n , A_4 , S_4 or A_5 . The purpose of this paper is to determine this minimum genus for the simple groups $PSl_2(p)$, where $p \geq 5$ is a prime. Since given any finite group G and Riemann surface T there exists a regular branched covering $p: S \rightarrow T$ such that i) G is the group of branched covering transformations of p (i.e. $T = S/G$) and ii) G is the full group of automorphisms of S [Gr], it seems most interesting to realize G as the full group of automorphisms of a Riemann surface of least genus. In a sequel to this paper [GS] we will prove that this always happens when $p \not\equiv \pm 1 \pmod{8}$ or $\pmod{5}$ but *may* fail for these congruence equalities. When it does fail $PSl_2(p)$ will have index two in the full group of automorphisms. In addition, a particularly simple situation occurs when $p: S \rightarrow S/G$ has exactly three branch points. Our results always give this for $PSl_2(p)$. We conjecture analogous results for every finite simple group and we seek to relate these ideas to “moonshine” for simple groups [FLM]. In order to state our results we need some notation:

- (1) $PSl_2(p^k)$ is the projective special linear group of 2×2 matrices over the Galois field $GF(p^k)$.
- (2) $\Gamma = PSl_2(\mathbf{Z})$ is the classical modular group. Geometrically Γ is just the group of integral linear fractional transformations of the upper half plane H , that is transformations of the form $z \rightarrow \frac{az + b}{cz + d}$, where a, b, c, d

¹⁾ Research partially supported by N.S.E.R.C. grant 67-7218.

are integers so that $ad - bc = 1$. Algebraically Γ is the unimodular group $Sl_2(\mathbb{Z})$ modulo its center $= \{\pm I\}$.

A result of Newman [N] is that mod p reduction of entries gives an epimorphism $\Gamma \twoheadrightarrow PSl_2(p)$, and therefore an exact sequence $1 \rightarrow \Delta \rightarrow \Gamma \rightarrow PSl_2(p) \rightarrow 1$. Now Δ is a Fuchsian group and therefore $PSl_2(p)$ is acting conformally on the open Riemann surface H/Δ . By adding parabolic points we obtain a closed Riemann surface $\overline{H/\Delta}$ and a conformal action on $\overline{H/\Delta}$ by extension. According to [G] the genus of $\overline{H/\Delta}$ is

$$1 + \frac{|PSl_2(p)|}{2} \left(\frac{1}{6} - \frac{1}{p} \right) = 1 + \frac{p(p^2-1)}{4} \left(\frac{1}{6} - \frac{1}{p} \right)$$

where $|PSl_2(p)| = \frac{p(p^2-1)}{2}$ is the order of $PSl_2(p)$.

Definition. For any finite group G we let *genus* (G) denote the least genus of all Riemann surfaces S for which there exists an effective conformal action $G \times S \rightarrow S$. We note that *genus* (G) has also been called the symmetric genus of G in the literature.

Thus we certainly have $\text{genus}(PSl_2(p)) \leq 1 + \frac{p(p^2-1)}{4} \left(\frac{1}{6} - \frac{1}{p} \right)$. Putting $p = 5$ then gives $\text{genus}(PSl_2(5)) = 0$, and therefore we will tacitly assume in all that follows that $p \geq 7$.

For $p = 7, 11$ we get the inequalities $\text{genus}(PSl_2(7)) \leq 3$ and $\text{genus}(PSl_2(11)) \leq 26$. It will turn out that these inequalities are equalities (see the corollary of the introduction). The action of $PSl_2(7)$ on a surface of genus 3 is the action of the simple group of order 168 considered by Klein.

This inequality strongly suggests that $\text{genus}(PSl_2(p))$ can be calculated by realizing $PSl_2(p)$ as an epimorphic image of Γ , or some other Fuchsian group, and then minimizing over all such epimorphisms. For example Γ has the presentation:

$$\Gamma = \{S, T \mid S^2 = (ST)^3 = 1\},$$

where $S = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ and $T = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$.

Reducing coefficients mod p leads to a presentation of $PSl_2(p)$, namely

$$PSl_2(p) = \{A, B, C \mid A^2 = B^3 = C^p = ABC = 1, ETC\}$$

where we have made the substitutions $A = S$, $B = ST$ and $C = T^{-1}$. We have written the presentation in this manner so that it becomes clear that $PSl_2(p)$ is an epimorphic image of the triangle group

$$T(2, 3, p) = \{A, B, C \mid A^2 = B^3 = C^p = ABC = 1\}.$$

Recall that if r, s, t are integers ≥ 2 then $T(r, s, t)$ is the group of orientation preserving symmetries of the appropriate plane generated by rotations of $2\pi/r$, $2\pi/s$ and $2\pi/t$, respectively, about the vertices of a triangle having angles π/r , π/s and π/t respectively. The plane is spherical if $1/r + 1/s + 1/t > 1$, euclidean if $1/r + 1/s + 1/t = 1$, and hyperbolic if $1/r + 1/s + 1/t < 1$. See Magnus [M] for more details.

Using the above presentation of $PSl_2(p)$ leads to an exact sequence $1 \rightarrow \Delta \rightarrow T(2, 3, p) \rightarrow PSl_2(p) \rightarrow 1$ and an effective conformal action of $PSl_2(p)$ on the closed Riemann surface H/Δ . Again we have

$$\text{genus } (H/\Delta) = 1 + \frac{p(p^2-1)}{4} \left(\frac{1}{6} - \frac{1}{p} \right)$$

so there is no improvement. But now the idea is clear: find all triples (r, s, t) for which there is an exact sequence $1 \rightarrow \Delta \rightarrow T(r, s, t) \rightarrow PSl_2(p) \rightarrow 1$, compute the genus of H/Δ for any such extension, and then minimize over all possible triples. It turns out that this procedure gives genus $(PSl_2(p))$ because more branch points always gives a higher genus.

If $p \geq 13$ we make the definition $d = \min\{e \mid e \geq 7 \text{ and either } e \mid \frac{p-1}{2} \text{ or } e \mid \frac{p+1}{2}\}$. Then our results are:

THEOREM I. *Assume $p \geq 13$. Then there exists a short exact sequence $1 \rightarrow \Delta \rightarrow T(2, 3, d) \rightarrow PSl_2(p) \rightarrow 1$ and the genus of H/Δ is*

$$1 + \frac{p(p^2-1)}{4} \left(\frac{1}{6} - \frac{1}{d} \right).$$

THEOREM II.

- (a) *If $p \equiv \pm 1 \pmod{5}$ then there exists a short exact sequence $1 \rightarrow \Delta \rightarrow T(2, 5, 5) \rightarrow PSl_2(p) \rightarrow 1$ and the genus of H/Δ is $1 + \frac{p(p^2-1)}{40}$.*
- (b) *If $p \equiv \pm 1 \pmod{8}$ then there exists a short exact sequence $1 \rightarrow \Delta \rightarrow T(3, 3, 4) \rightarrow PSl_2(p) \rightarrow 1$ and the genus of H/Δ is $1 + \frac{p(p^2-1)}{48}$.*

(c) If $p \equiv \pm 1 \pmod{5}$ and $p \equiv \pm 1 \pmod{8}$ then there exists a short exact sequence $1 \rightarrow \Delta \rightarrow T(2, 4, 5) \rightarrow PSl_2(p) \rightarrow 1$ and the genus of H/Δ is $1 + \frac{p(p^2-1)}{80}$.

Then we will prove that genus $(PSl_2(p))$ is obtained by minimizing over all the possibilities above.

The result of this minimization is

COROLLARY. *The genus of $PSl_2(p)$ is given as follows:*

(a) $g = 1 + \frac{p(p^2-1)}{4} \left(\frac{1}{6} - \frac{1}{p} \right)$ if $p = 5, 7, 11$,

(b) $g = 1 + \frac{p(p^2-1)}{40}$ if $p \geq 13$, $p \equiv \pm 1 \pmod{5}$, $p \not\equiv \pm 1 \pmod{8}$
and $d \geq 15$,

(c) $g = 1 + \frac{p(p^2-1)}{48}$ if $p \geq 13$, $p \not\equiv \pm 1 \pmod{5}$, $p \equiv \pm 1 \pmod{8}$
and $d \geq 12$,

(d) $g = 1 + \frac{p(p^2-1)}{80}$ if $p \geq 13$, $p \equiv \pm 1 \pmod{5}$, $p \equiv \pm 1 \pmod{8}$
and $d \geq 9$,

(e) $g = 1 + \frac{p(p^2-1)}{4} \left(\frac{1}{6} - \frac{1}{d} \right)$ in all other cases.

In fact the least genus g always comes from the branched covering space action on the Riemann surface $S = H/\Delta$ associated to some extension $1 \rightarrow \Delta \rightarrow T(r, s, t) \rightarrow PSl_2(p) \rightarrow 1$, where

$$(r, s, t) = \begin{cases} (2, 3, p) & \text{if } p = 5, 7, 11, \\ (2, 5, 5) & \text{if } p \geq 13, \quad p \equiv \pm 1 \pmod{5}, \quad p \not\equiv \pm 1 \pmod{8} \quad \text{and} \quad d \geq 15, \\ (3, 3, 4) & \text{if } p \geq 13, \quad p \not\equiv \pm 1 \pmod{5}, \quad p \equiv \pm 1 \pmod{8} \quad \text{and} \quad d \geq 12, \\ (2, 4, 5) & \text{if } p \geq 13, \quad p \equiv \pm 1 \pmod{5}, \quad p \equiv \pm 1 \pmod{8} \quad \text{and} \quad d \geq 9, \\ (2, 3, d) & \text{in all other cases.} \end{cases}$$

It turns out that other triples (r, s, t) are not relevant for the determination of the minimal genus.

In most cases the answer is $(r, s, t) = (2, 3, d)$. For $p \leq 617$ the triple $(2, 5, 5)$ occurs once exactly, namely for $p = 509$, $(3, 3, 4)$ occurs exactly three

times, namely for $p = 103, 137$ and 569 and $(2, 4, 5)$ occurs exactly six times, for $p = 199, 239, 359, 439, 521$ and 599 .

If $S = H/\Delta$ is the surface of minimal genus for $PSl_2(p)$ coming from one of the extensions above then the orbit manifold $S/PSl_2(p)$ is the 2-sphere S^2 and the quotient map $S \rightarrow S^2$ is a branched covering with exactly 3 branch points. One of the most important steps in the proof of the main result of this paper is the converse, namely if S is a Riemann surface of least genus for the group $G = PSl_2(p)$ then $S/G = S^2$ and $S \rightarrow S^2$ is a branched covering with exactly 3 branch points (see section 3). Note that a related notion of genus, “the Cayley genus of a group” has been studied by others, among them Tucker [T]. Earlier results can be found in Hurwitz [H] and Burnside [B].

The remainder of this paper is organized as follows. In section 2 we describe various ways of generating $PSl_2(p)$ and then prove theorems I and II. Section 3 proves that if S is a Riemann surface of least genus for $PSl_2(p)$ then $S/PSl_2(p)$ is a 2-sphere S^2 and the branched covering $S \rightarrow S^2$ has exactly 3 branch points. The calculation of genus ($PSl_2(p)$) then follows from the results of section 2.

Finally we would like to thank Bomshik Chang for help with the group theory of $PSl_2(p)$. The first author would like to thank the University of British Columbia for its hospitality to him during the time this research was done.

§ 2. GENERATING TRIPLES FOR $PSl_2(p)$

Our goal in this section is to find triples (r, s, t) for which there are epimorphisms $T(r, s, t) \twoheadrightarrow PSl_2(p)$. In other words, given integers $r, s, t \geq 2$ are there matrices $A, B, C \in PSl_2(p)$ so that A, B, C generate $PSl_2(p)$ and $A^r = B^s = C^t = ABC = 1$? Throughout this section a standard reference for the group theory is Suzuki [S].

The spherical triangle groups are given in the following table

TABLE I

<i>triple</i>	<i>triangle group</i>	<i>order</i>
$(2, 2, n)$	dihedral	$2n$
$(2, 3, 3)$	tetrahedral (A_4)	12
$(2, 3, 4)$	octahedral (S_4)	24
$(2, 3, 5)$	icosahedral (A_5)	60