Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 31 (1985)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: HODGE DECOMPOSITION ON STRATIFIED LIE GROUPS
Autor: Duddy, John
Kapitel: 2. CR STRUCTURES AND THE HEISENBERG GROUP
DOI: https://doi.org/10.5169/seals-54573

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 06.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Let H be the kernel of Δ in $C_c^\infty(E_2)$. A Hodge decomposition for $C_c^\infty(E_2)$ is

$$C_c^\infty(E_2) = D_1(C_c^\infty(E_1)) \oplus D_2^*(C_c^\infty(E_3)) \oplus H .$$

Hodge studied the de Rham complex on a compact Riemannian manifold. The Riemannian metric induced the metrics on the bundles $\Lambda^p T^*(M)$ as well as the volume element.

In the next section we discuss abstract CR manifolds and look at the Heisenberg group in detail. We write down the $\bar{\partial}_b$ and $\Box_b$ operators explicitly and give Folland and Stein's inverse to $\Box_b$. In section 3 we introduce the stratified Lie groups and the associated homogeneous structures. We present the continuity theorems of Folland and Rothschild and Stein for convolution operators. In section 4 we prove the decomposition theorem in the general setting of stratified groups.

These results are an extension of the author's dissertation [4]. We wish to express our deep gratitude to M. Kuranishi. We would also like to thank D. Tartakoff for his help and encouragement.

2. CR STRUCTURES AND THE HEISENBERG GROUP

Let M be a C^∞ manifold of dimension $2n + 1$. The complexified tangent bundle of M, $\mathbf{C}T(M)$, is the bundle whose fiber is $\mathbf{C} \otimes_{\mathbf{R}} T_m(M)$ where $T_m(M)$ is the tangent space at $m \in M$. When there is no confusion we will drop reference to M in the notation for $T(M)$, $\mathbf{C}T(M)$, etc. So, $T(M) = T$ and $\mathbf{C}T(M) = \mathbf{C}T$, for example.

A CR structure on M is a subbundle $T_{1,0} \subset \mathbf{C}T$ such that (i) $T_{1,0} \cap \overline{T_{1,0}} = \{0\}$, (ii) $\operatorname{codim}(T_{1,0} \otimes \overline{T_{1,0}}) = 1$, (iii) if X and Y are smooth sections of $T_{1,0}$ then $[X, Y] = XY - YX$ is a section $T_{1,0}$. We set $T_{0,1} = \overline{T_{1,0}}$. If M has a CR structure it is called a CR manifold.

An example of a CR manifold is a real hypersurface M in a complex manifold M', $M \subset M'$. Define $T_{1,0}(M) = \mathbf{C}T(M) \cap T_{1,0}(M')$ where $T_{1,0}(M')$ is the holomorphic tangent bundle of M'.

If M is a CR manifold set $T^{1,0}$ (resp., $T^{0,1}$) to be the dual space to $T_{1,0}$ (resp., $T_{0,1}$). Let $\Lambda^{p,q}$ be the space of C^∞ sections of $\Lambda^p T^{1,0} \otimes \Lambda^q T^{0,1}$. Define the operator $\bar{\partial}_b : \Lambda^{p,q} \to \Lambda^{p,q+1}$ as follows: Let $\phi \in \Lambda^{p,q}$ and let $X_1, \ldots, X_p$ (resp., $Y_1, \ldots, Y_{q+1}$) be sections of $T_{1,0}$ (resp., $T_{0,1}$). Then

$$<\bar{\partial}_b\phi; (X_1 \wedge \dots \wedge X_p) \otimes (Y_1 \wedge \dots \wedge Y_{q+1})>$$
$$= (q+1)^{-1} \sum_{j=1}^{q+1} (-1)^{j+1} Y_j <\phi; (X_1 \wedge \dots \wedge X_p) \otimes (Y_1 \wedge \dots \hat{Y}_j \dots \wedge Y_{q+1})>$$
$$+ (q+1)^{-1} \sum_{i<j} (-1)^{i+j} <\phi; (X_1 \wedge \dots \wedge X_p) \otimes ([Y_i, Y_j] \wedge Y_1 \wedge \dots \hat{Y}_i \dots \hat{Y}_j \dots \wedge Y_{q+1})>.$$

The $\wedge$ symbol over a section means as usual that it is deleted from the expression. One can show that

i) $\bar{\partial}_b^2 = 0$,

ii) $\bar{\partial}_b(\phi \wedge \psi) = (\bar{\partial}_b\phi) \wedge \psi + (-1)^p \phi \wedge \bar{\partial}_b\psi$ for $\phi \in \Lambda^{0,p}$,

iii) $<\bar{\partial}_b f, Y> = Yf$ for $f \in \Lambda^{0,0}$ and Y a section of $T_{0,1}$.

See [6] for details.

The Heisenberg group, H, is a Lie group with a natural CR structure. The manifold is $\mathbf{C}^n \times \mathbf{R}$. Let (z, t), $(z', t') \in \mathbf{C}^n \times \mathbf{R} = H$. The group law is defined by

$$(z, t) \cdot (z', t') = \big(z+z', t+t'+2\,\mathrm{Im}(z \cdot z')\big)$$

where $z \cdot z' = \sum_{j=1}^{n} z_j \bar{z}'_j$. The identity element is $(0, 0)$ and $(z, t)^{-1} = (-z, -t)$.

Sometimes we will set $u = (z, t)$.

For $j = 1, \dots, n$ if we set $z_j = x_j + iy_j$, the mapping

$$(z, t) \to (x_1, \dots, x_n, y_1, \dots, y_n, t)$$

defines a C^∞ coordinate system on H. The left invariant vector fields (i.e., the elements of the Lie algebra) are $\mathbf{R}$-linear combinations of

$$X_j = \frac{\partial}{\partial x_j} + 2y_j \frac{\partial}{\partial t}, \quad Y_j = \frac{\partial}{\partial y_j} - 2x_j \frac{\partial}{\partial t}, \quad T = \frac{\partial}{\partial t}. \tag{2}$$

They satisfy the following commutation relations:

$$[X_j, T] = [Y_j, T] = [X_j, X_k] = [Y_j, Y_k] = 0, \tag{3}$$

$$[X_j, Y_k] = -4\delta_{jk} T. \tag{4}$$

Let $\mathbf{C}T$ be the complexified tangent bundle of H. Define

$$Z_j = \frac{1}{2}(X_j - iY_j) = \frac{\partial}{\partial z_j} + i\bar{z}_j \frac{\partial}{\partial t}. \tag{5}$$

Then Z_j, $\bar{Z}_j$, and T form a basis (over $\mathbf{R}$) of the space of left invariant complex tangent vector fields. In particular, they form a global frame of $\mathbf{C}T$. From (3), (4) and (5) we easily see that

$$(6) \qquad [Z_j, Z_k] = [\bar{Z}_j, \bar{Z}_k] = [Z_j, T] = [\bar{Z}_j, T] = 0$$
$$[Z_j, \bar{Z}_k] = -2i\delta_{jk}T .$$

Let $T_{1,0}$ (resp., $T_{0,1}$) be the subbundle of $\mathbf{C}T$ spanned by the Z_j's (resp., $\bar{Z}_j$'s). Then

$$\bar{T}_{1,0} = T_{0,1}$$
$$T_{1,0} \cap T_{0,1} = \{0\}$$
$$\operatorname{codim}(T_{1,0} \otimes T_{0,1}) = 1 .$$

Also, if V_1, V_2 are sections of $T^{1,0}$ we can write $V_i = \sum_{j=1}^{n} f_{ij}Z_j$, $i = 1, 2$ where the f_{ij} are C^∞ functions on H. Then by (6),

$$[V_1, V_2] = \sum_{k=1}^{n} \left(\sum_{j=1}^{n} (f_{1j}Z_j f_{2k} - f_{2j}Z_j f_{1k}) \right) Z_k .$$

So, the splitting of $\mathbf{C}T$ defines a CR structure on H.

Impose the left invariant Hermitian metric on $\mathbf{C}T$ which makes the Z's, $\bar{Z}$'s and T an orthonormal frame. Let ω^j and τ be dual to Z_j and T, respectively. Then ω^j, $\bar{\omega}^j$ and τ form an orthonormal frame for $\mathbf{C}T^*$. The volume element on H is

$$(7) \qquad du = 2^n dx_1 \wedge \dots \wedge dx_n \wedge dy_1 \wedge \dots \wedge dy_n \wedge dt .$$

Since H is nilpotent and, hence, unimodular, the volume element is both left and right invariant. One can also verify this directly.

Let $J = (j_1, \dots, j_q)$ be a multi-index with $1 \leqslant j_i \leqslant n$, $i = 1, \dots, q$. Define $|J| = q$ and $\bar{\omega}^J = \bar{\omega}^{j_1} \wedge \dots \wedge \bar{\omega}^{j_q}$. If $\phi \in \Lambda^{0,q}(H)$ we may write $\phi = \sum_{|J|=q} \phi_J \bar{\omega}^J$ where ϕ_J is a C^∞ function from H to $\mathbf{C}$. Let

$$\bar{\omega}^j \lrcorner\, \bar{\omega}^J = (-1)^k \bar{\omega}^{j_1} \wedge \dots \wedge \widehat{\bar{\omega}^{j_k}} \wedge \dots \wedge \bar{\omega}^{j_q} \quad \text{if} \quad j = j_k \quad \text{and} \quad \bar{\omega}^j \lrcorner\, \bar{\omega}^J = 0$$

otherwise. Folland and Stein prove that for $\phi \in \Lambda^{0,q}$

$$\text{i)} \quad \bar{\partial}_b \phi = \sum_{|J|=q} \sum_{j=1}^{n} \bar{Z}_j \phi_J \bar{\omega}^j \wedge \bar{\omega}^J ,$$

(8) ii) $\bar{\partial}_b^* \phi = -\sum_{|J|=q} \sum_{j=1}^{n} Z_j \phi_J \bar{\omega}^j \lrcorner\, \bar{\omega}^J$,

iii) $\Box_b \phi = \sum_{|J|=q} \left(-\frac{1}{2} \sum_{j=1}^{n} (Z_j \bar{Z}_j + \bar{Z}_j Z_j) + i(n-2q)T \right) \phi_J \bar{\omega}^J$.

Define the function

$$\Phi_\alpha(z, t) = (|z|^2 - it)^{-\frac{n+\alpha}{2}} (|z|^2 + it)^{-\frac{n-\alpha}{2}}.$$

Let $\phi \in \Lambda_c^{0,q}$, $q \neq 0, n$. For an appropriate constant, c_q, define

(9) $$K_q \phi(v) = c_q \sum_{|J|=q} \left(\int_H \phi_J(u) \Phi_{n-2q}(u^{-1}v) du \right) \bar{\omega}^J.$$

Folland and Stein prove that for the appropriate c_q

THEOREM 1. *Let* $\phi \in \Lambda_c^{0,q}$, $q \neq 0, n$. *Then* $\Box_b K_q \phi = K_q \Box_b \phi = \phi$.

In [4] we prove a stronger version of the following Hodge decomposition theorem.

THEOREM 2. *Let* $\phi \in \Lambda_c^{0,q}$, $q \neq 0, n$. *Then*

i) $H\phi = 0$ *where* H *is the orthogonal projection onto the kernel of* $\Box_b$.

ii) $\phi = \bar{\partial}_b \bar{\partial}_b^* K_q \phi + \bar{\partial}_b^* \bar{\partial}_b K_q \phi$.

We also prove

THEOREM 3. *If* $\phi \in \Lambda_c^{0,q}$, $q \neq 0, n$ *and if* $\bar{\partial}_b \phi = 0$ *then* $\psi = \bar{\partial}_b^* K_q \phi$ *satisfies* $\bar{\partial}_b \psi = \phi$.

These two theorems are special cases of theorems 6 and 7 proven in section 4.

3. DIFFERENTIAL COMPLEXES ON STRATIFIED GROUPS

We study a class of nilpotent Lie groups which we describe in terms of their Lie algebras. A graded Lie algebra, $\mathfrak{n}$, is a finite dimensional nilpotent algebra which has a direct sum decomposition, $\mathfrak{n} = \bigoplus_{i=1}^{r} \mathfrak{n}_i$ where the $\mathfrak{n}_i$ satisfy

i) $[\mathfrak{n}_i, \mathfrak{n}_j] \subseteq \mathfrak{n}_{i+j}$ if $i + j \leqslant r$,

ii) $[\mathfrak{n}_i, \mathfrak{n}_j] = 0$ if $i + j > r$.