Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 31 (1985)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: HODGE DECOMPOSITION ON STRATIFIED LIE GROUPS
Autor: Duddy, John
Kapitel: 4. The Hodge decomposition
DOI: https://doi.org/10.5169/seals-54573

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 22.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

THEOREM 4. (i) *Let* K *be a convolution operator of type* r *for* $r > 0$. *Then* K *extends from* $C_c^\infty(\mathbf{F}^m)$ *to a bounded operator from* $L^p(\mathbf{F}^m)$ *to* $L^q(\mathbf{F}^{m'})$ *where* $1 < p < Q/r$ *and* $q^{-1} = p^{-1} - r/Q$. (ii) *Let* K *be a convolution operator of type* 0. *Then* K *extends from* $C_c^\infty(\mathbf{F}^m)$ *to a bounded operator from* $S_k^p(\mathbf{F}^m)$ *to* $S_k^p(\mathbf{F}^{m'})$.

Finally, we mention the interaction between the homogeneous convolution operators and the left-invariant differential operators. Let $D: C^\infty(\mathbf{F}^{m'}) \to C^\infty(\mathbf{F}^{m''})$ be a left-invariant homogeneous differential operator of degree 1 and let K be a homogeneous convolution operator of type r, with $r \geqslant 1$. Then DK is a homogeneous convolution operator of type $r - 1$. Moreover, if $r > 1$ the kernel of DK is given by $Dk(x)$.

4. THE HODGE DECOMPOSITION

Consider the complex (1) where $E_i = \mathbf{R}^n \times \mathbf{F}^{m_i}$. Assume that each of the D_i is a first order, left-invariant operator, homogeneous of degree 1. So each entry of D_i is of the form $\sum_{j=1}^{d} a_j X_{1,j}$ where a_j is constant. Construct the Laplacian, Δ, with respect to the euclidian inner products on $\mathbf{F}^{m_i}$, $i = 1, 2, 3$. Assume there exists a homogeneous convolution operator of type 2, K, which inverts Δ. If $f \in C_c^\infty(\mathbf{F}^{m_2})$ then $f(x) = \Delta K f(x) = K\Delta f(x)$.

THEOREM 5. *Let* $f \in S_2^2(\mathbf{F}^{m_2})$. *As distributions,* $\Delta f = 0$ *if and only if* $f = 0$.

Proof. Obviously, if $f = 0$ then $\Delta f = 0$.

Assume $\Delta f = 0$. Let $\{f_j\}$ be a sequence in $C_c^\infty(\mathbf{F}^{m_2})$ such that $f_j \to f$ in $S_2^2(\mathbf{F}^{m_2})$. Then $f_j \to f$ in the sense of distributions. Moreover, $\Delta f_j \to \Delta f = 0$ in $L^2(\mathbf{F}^{m_2})$. Let $g \in C_c^\infty(\mathbf{F}^{m_2})$. Then

$$<f, g> = \lim_{j\to\infty} <f_j, g> = \lim_{j\to\infty} <f_j, \Delta K g> = \lim_{j\to\infty} <\Delta f_j, Kg> .$$

Because $g \in C_c^\infty(\mathbf{F}^{m_2})$ it is in L^p where $p = 2Q/(Q+4)$. Therefore, by Theorem 4(i), $Kg \in L^q$ where

$$q^{-1} = (Q+4)/2Q - 2/Q = 1/2, \text{ i.e., } Kg \in L^2(\mathbf{F}^{m_2}) .$$

For $Q \geqslant 5$, $1 < p < q < \infty$. So

$$| <f, g> | = \lim_{j\to\infty} | <\Delta f_j, Kg> | \leqslant \lim_{j\to\infty} \| \Delta f_j \|_{L^2(\mathbf{F}^{m_2})} \| Kg \|_{L^2(\mathbf{F}^{m_2})} = 0 .$$

So, as a distribution, $f = 0$. This proves the theorem.

We have shown that the only harmonic element in $S_2^2(\mathbf{F}^{m_2})$ is the zero element. Let $f \in S_2^2(\mathbf{F}^{m_2})$ and let $f_j \to f$ in $S_2^2(\mathbf{F}^{m_2})$ with $f_j \in C_c^\infty(\mathbf{F}^{m_2})$. Then

$$f = \lim_{j\to\infty} \Delta K f_j = \lim_{j\to\infty} D_1 D_1^* K f + \lim_{j\to\infty} D_2^* D_2 K f = D_1 D_1^* K f + D_2^* D_2 K f .$$

To complete the Hodge decomposition we must prove that

$$D_1 D_1^* K f \perp D_2^* D_2 K f .$$

We need the following notation. Let $D(R) = \{x \in N : |x| < R\}$ and $S(R) = \{x \in N : |x| = R\}$. Endow each set with the left-invariant metric induced by N. The metric gives rise to the corresponding volume elements which, in the case of $D(R)$, is the restriction of dx. Let $d\mu_R$ denote the volume element on $S(R)$. For $f, g \in C_c^\infty(D(R), \mathbf{F}^{m_i})$ define

$$(f, g)_{D(R), i} = \int_{D(R)} (f(x), g(x))_{i,x} dx$$

where $(\ ,\ )_{i,x}$ is the metric on $\mathbf{F}^{m_i}$. Similarly for $f, g \in C^\infty(S(R), \mathbf{F}^{m_i})$ define

$$(f, g)_{S(R), i} = \int_{S(R)} (f(x), g(x))_{i,x} d\mu_R(x) .$$

By restriction, any element $f \in C^\infty(N, \mathbf{F}^{m_i})$ gives rise to an element of $C^\infty(S(R), \mathbf{F}^{m_i})$ or $C^\infty(D(R), \mathbf{F}^{m_i})$. In our notation, we will not distinguish f from its restrictions.

We will be integrating by parts on $D(R)$ which will involve a boundary integral on $S(R)$. To that end we define the symbol of our differential operators. Define $\xi(x) = |x| - R$ and let $g \in C^1(N, \mathbf{F}^{m_i})$. Let $x \in S(R)$. Then $\xi(x) = 0$. The symbol of D_i at $x \in S(R)$ acting on $d\xi$ and on g is given by

$$\sigma(D_i, d\xi) g(x) = D_i(\xi g)(x) .$$

The integration by parts formula is

$$(D_i g, f)_{D(R), i+1} = (g, D_i^* f)_{D(R), i} + (\sigma(D_i, d\xi) g, f)_{S(R), i} . \tag{13}$$

THEOREM 6. *Assume* $f \in L^2(\mathbf{F}^{m_2}) \cap L^q(\mathbf{F}^{m_2})$ *where* $q = Q/Q + 2$. *Then* $f = D_1 D_1^* K f + D_2^* D_2 K f$ *and* $D_1 D_1^* K f \perp D_2^* D_2 K f$.

Proof. We have already seen that $f = D_1 D_1^* K f + D_2^* D_2 K f$. To prove the orthogonality we restrict our attention to $D(R)$ for R large.

For brevity let $h = (D_1 D_1^* Kf, D_2^* D_2 Kf)_2$. Also, let

$$h(R) = (D_1 D_1^* Kf, D_2^* D_2 Kf)_{D(R),2}.$$

Then $\lim_{R\to\infty} h(R) = h$. Note that $D_1 D_1^* K$ and $D_2^* D_2 K$ are type 0 operators. Since $f \in L^2(\mathbf{F}^{m_2})$ by theorem 4 (ii) we know that h is defined. Furthermore $h(R)$ is bounded for all R by $\| D_1 D_1^* Kf \|_{L^2(R^{m_2})} \| D_2^* D_2 Kf \|_{L^2(R^{m_2})}$.

We can compute $h(R)$ as follows:

$$h(R) = \big(D_1 D_1^* Kf, D_2^* D_2(Kf)\big)_{D(R),2} = (D_2 D_1 D_1^* Kf, D_2 Kf)_{D(R),3} + \big(D_1 D_1^* Kf, \sigma(D_2^*, d(|x|))D_2 Kf\big)_{S(R),2}$$

by (13). We now prove a sequence of lemmas.

LEMMA 1. *$h(R)$ is continuous.*

Proof. This follows from Lebesgue's dominated convergence theorem.

LEMMA 2. *Let $R \geqslant 1$, $x \in N$ and $|x| = R$. Then*

$$|\sigma(D_2^*, d|x|)g(x)| \leqslant C\,|g(x)|$$

where C is a constant independent of g.

Proof. Recall that $X_1, \ldots, X_d$ is our orthonormal basis for $\mathfrak{n}_1$ where $d = \dim(\mathfrak{n}_1)$. The entries of D_2^* are linear combinations of the X_i, $i = 1, \ldots, d$ with coefficients in $\mathbf{F}$. Let D_{ij} be the i,j entry, $1 \leqslant i \leqslant m_2$, $1 \leqslant j \leqslant m_3$. Then $D_{ij} = \sum_{k=1}^{d} C_{ij}^k X_k$. Thus, for $x \in S(R)$

$$\begin{aligned}
|\sigma(D_2^*, d|x|)g(x)| &\leqslant C \sum_{i=1}^{m_2} \Big| \sum_{j=1}^{m_3} D_{ij}\big((|x|-R)g_j(x)\big) \Big| \\
&\leqslant C \sum_{i,j,k} | C_{ij}^k X_k\big((|x|-R)g_j(x)\big) | \\
&\leqslant C \sum_{jk} | (X_k|x|)g_j(x) | \qquad (\text{since } |x| = R) \\
&\leqslant C\big(\max_k (X_k|x|)\big)\, |g_j(x)|.
\end{aligned}$$

We must show $X_k|x|$ is bounded. But $|x|$ is C^∞ away from the origin and homogeneous of degree 1. So $X_k|x|$ is homogeneous of degree 0. Thus, it is determined by its values on $\{|x| = 1\}$. It is C^∞ on this set and, therefore, bounded. This proves the lemma.

Since $h(R) \to h$ as $R \to \infty$ we have

LEMMA 3. *For* $\varepsilon > 0$, $\lim_{r\to\infty} \frac{1}{2\varepsilon}\int_{r-\varepsilon}^{r+\varepsilon} h(R)dR = h$.

We continue with the proof of our theorem. By the preceding lemma it suffices for us to prove that $\lim_{r\to\infty} \frac{1}{2\varepsilon}\int_{r-\varepsilon}^{r+\varepsilon} h(R)dR = 0$. But,

$$\frac{1}{2\varepsilon}\int_{r-\varepsilon}^{r+\varepsilon} h(R)dR \tag{14}$$

$$= \frac{1}{2\varepsilon}\int_{r-\varepsilon\leqslant|x|\leqslant r+\varepsilon} (D_1D_1^*Kf, \sigma(D_2^*, d|x|)D_2Kf)_x \,\|\, d\,|\,x\,|\,\|\, dx$$

because $d\mu_R dR = \|\, d\,|\,x\,|\,\|\, dx$. We claim that $\|\, d\,|\,x\,|\,\|$ is bounded. Let ω^{ij} be dual to X_{ij} where X_{ij} is our orthonormal basis. Then $d\,|\,x\,| = \sum_{i,j}(X_{ij}|x|)\omega^{ij}$. Since $|\,x\,|$ is homogeneous of degree 1 and X_{ij} is homogeneous of degree i we have $X_{ij}\,|\,x\,|$ is homogeneous of degree $1-i$. Hence, for $|\,x\,| \geqslant 1$ each $X_{ij}\,|\,x\,|$ is bounded. So $\|\, d\,|\,x\,|\,\|$ is bounded.

By assumption, $f \in L^q(\mathbf{F}^{m_2})$, $q = Q/Q+2$ and we know that D_2K is type 1. By Theorem 4(i) we know that $D_2Kf \in L^2(\mathbf{F}^{m_2})$. Thus, by Lemma 2 $|\,\chi_r\sigma(D_2^*, d|x|)D_2Kf\,| \leqslant C\,|\,\chi_rD_2Kf\,|$ where χ_r is the characteristic function of $\{r-\varepsilon \leqslant |\,x\,| \leqslant r+\varepsilon\}$. We conclude that $\chi_r\sigma(D_2^*, d|x|)D_2Kf \in L^2(\mathbf{F}^{m_2})$. By the Schwarz inequality and the fact that $\|\, d\,|\,x\,|\,\|$ is bounded, we get, from (14)

$$\frac{1}{2\varepsilon}\int_{r-\varepsilon}^{r+\varepsilon} h(R)dR \leqslant \frac{c}{2\varepsilon}\,\|\,\chi_rD_1D_1^*Kf\,\|_{L^2(\mathbf{F}^{m_2})}\,\|\,\chi_rD_2Kf\,\|_{L^2(\mathbf{F}^{m_3})}\,.$$

As $r\to\infty$, both $\|\,\chi_rD_1D_1^*Kf\,\|_{L^2(\mathbf{F}^{m_2})}$ and $\|\,\chi_rD_2Kf\,\|_{L^2(\mathbf{F}^{m_3})}$ tend to 0. So $h = \lim_{r\to\infty}\frac{1}{2\varepsilon}\int_{r-\varepsilon}^{r+\varepsilon} h(R)dR = 0$. This proves the theorem.

This theorem together with Theorem 5 proves the Hodge decomposition. A similar argument gives the solution to the problem of finding g such that $D_1g = f$ for a given f.

THEOREM 7. *Let* $f \in L^2(\mathbf{F}^{m_2}) \cap L^q(\mathbf{F}^{m_2})$ *with* $q = Q/Q+2$. *Suppose* $D_2f = 0$. *Then there exists* $g \in L^2(\mathbf{F}^{m_1})$ *such that* $D_1g = f$.

Proof. We have $f = D_1D_1^*Kf + D_2^*D_2Kf$. It suffices to prove $(f, D_2^*D_2Kf) = 0$ because this implies $D_2^*D_2Kf = 0$ since

$$D_1D_1^*Kf \perp D_2^*D_2Kf\,.$$

We may set $g = D_1^* Kf$. Using the same notation as in the preceding theorem we have

$$(f, D_2^* D_2 Kf)_2 = \lim_{R \to \infty} (f, D_2^* D_2 Kf)_{D(R),\,2}$$

$$= \lim_{R \to \infty} \left((D_2 f, D_2 Kf)_{D(R),\,3} + \left(f, \sigma(D_2^*, d|x|) D_2 Kf\right)_{S(R),\,2}\right)$$

$$= \lim_{R \to \infty} \left(f, \sigma(D_2^*, d|x|) D_2 Kf\right)_{S(R),\,2} \qquad (\text{since } D_2 f = 0).$$

The same argument as in Theorem 6 proves that the limit is zero. This proves the theorem.

REFERENCES

[1] ATIYAH, M. F. and R. BOTT. A Lefschetz fixed point formula for elliptic complexes, I, II. *Ann. of Math. 86* (1967), 347-407; *88* (1968), 451-491.

[2] DADOK, J. and R. HARVEY. The fundamental solution for the Kohn-Laplacian $\square_b$ on the sphere in $\mathbf{C}^n$. *Math. Ann. 244* (1979), 89-104.

[3] DE RHAM, G. *Variétés Différentiables.* Paris: Hermann & Cie, 1955.

[4] DUDDY, J. A. *The boundary Laplacian on the Heisenberg group and the sphere.* Ph. D. dissertation, Columbia University, 1978.

[5] DOLBEAULT, P. Sur la cohomologie des variétés analytiques complexes. *C. R. Acad. Sci. Paris 236* (1953), 175-177.

[6] FOLLAND, G. B. and J. J. KOHN. *The Neumann Problem for the Cauchy-Riemann Complex.* Princeton, N.J.: Princeton University Press, 1972.

[7] FOLLAND, G. B. and E. M. STEIN. Parametrices and estimates for the $\bar{\partial}_b$ complex on strongly pseudoconvex boundaries. *Bull. Amer. Math. Soc. 80* (1974), 253-258.

[8] FOLLAND, G. B. Estimates for the $\bar{\partial}_b$ complex and analysis on the Heisenberg group. *Comm. Pure Appl. Math. 27* (1974), 429-522.

[9] —— Subelliptic estimates and function spaces on nilpotent Lie groups. *Ark. Math. 13* (1975), 161-207.

[10] GREENFIELD, S. J. Cauchy-Riemann equations on several variables. *Ann. Scuola Norm. Pisa 22* (1968), 275-314.

[11] HARVEY, R. and J. POLKING. Fundamental solutions in complex analysis I and II. *Duke Math. J. 46* (1979), 253-340.

[12] HODGE, W. V. D. *The Theory and Applications of Harmonic Integrals.* New York: Cambridge University Press, 1941 (2d ed., 1952).

[13] KODAIRA, K. Harmonic fields in Riemannian manifolds (generalized potential theory). *Ann. of Math. 50* (1949), 587-665.

[14] KOHN, J. J. and H. ROSSI. On the extension of holomorphic functions from the boundary of a complex manifold. *Ann. of Math. 81* (1965), 451-472.

[15] LEWY, H. An example of a smooth linear partial differential equation without solution. *Ann. of Math. 66* (1957), 155-158.