

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 31 (1985)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE TRACE AS AN ALGEBRA HOMOMORPHISM
Autor: Osborn, Howard
Kapitel: 1. Endomorphism algebras
DOI: <https://doi.org/10.5169/seals-54566>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 28.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

1. ENDOMORPHISM ALGEBRAS

In this section V will be an arbitrary module over a commutative ring R with unit, and for each $p \geq 0$ $\wedge^p V$ will be its p^{th} exterior power and $\text{End } \wedge^p V$ will be the R -module of endomorphisms $\wedge^p V \rightarrow \wedge^p V$; $\Pi_p \text{End } \wedge^p V$ will be the direct product of the R -modules $\text{End } \wedge^p V$. We shall define three distinct products in $\Pi_p \text{End } \wedge^p V$; the first two products are standard, and they will be used to define the third product. If $\wedge^p V$ itself vanishes for sufficiently large $p \geq 0$ the direct product $\Pi_p \text{End } \wedge^p V$ and the direct sum $\coprod_p \text{End } \wedge^p V$ agree; although this special condition will be satisfied in later sections the definitions in this section will be formulated in complete generality for the direct product $\Pi_p \text{End } \wedge^p V$.

Elements of $\Pi_p \text{End } \wedge^p V$ will be indicated by boldface capital letters $\mathbf{A}, \mathbf{B}, \dots$, the p^{th} components being $A_p, B_p, \dots \in \text{End } \wedge^p V$ for each $p \geq 0$. The simplest product in $\Pi_p \text{End } \wedge^p V$ is induced by compositions: the p^{th} component of the *composition product* $\mathbf{A}\mathbf{B} \in \Pi_p \text{End } \wedge^p V$ is the usual composition $A_p B_p \in \text{End } \wedge^p V$ of the endomorphisms A_p and B_p of $\wedge^p V$, where $A_p B_q = 0$ for $p \neq q$. Trivially $\Pi_p \text{End } \wedge^p V$ is an associative R -algebra with respect to the composition product, and there is a two-sided unit element \mathbf{I} whose p^{th} component is the identity endomorphism $I_p \in \text{End } \wedge^p V$ for each $p \geq 0$.

There is another reasonably familiar product in $\Pi_p \text{End } \wedge^p V$, the product of $A_p \in \text{End } \wedge^p V$ and $B_q \in \text{End } \wedge^q V$ providing an element

$$A_p \cdot B_q \in \text{End } \wedge^{p+q} V$$

for each $p \geq 0$ and each $q \geq 0$. Since elements of $\text{End } \wedge^{p+q} V$ are uniquely defined in terms of the behavior on exterior products $x_1 \wedge \dots \wedge x_{p+q} \in \wedge^{p+q} V$, it suffices to require that

$$(A_p \cdot B_q)(x_1 \wedge \dots \wedge x_{p+q}) = \sum_{\pi} \varepsilon_{\pi} A_p(x_{\pi_1} \wedge \dots \wedge x_{\pi_p}) \wedge B_q(x_{\pi_{p+1}} \wedge \dots \wedge x_{\pi_{p+q}})$$

where the sum is computed over all permutations π of $\{1, \dots, p+q\}$ such that both $\pi_1 < \dots < \pi_p$ and $\pi_{p+1} < \dots < \pi_{p+q}$, and where ε_{π} is the parity ± 1 of the permutation π . Such "shuffle products" $A_p \cdot B_q \in \text{End } \wedge^{p+q} V$ provide a unique *shuffle product* $\mathbf{A} \cdot \mathbf{B} \in \Pi_r \text{End } \wedge^r V$ of any two elements \mathbf{A} and \mathbf{B} in $\Pi_r \text{End } \wedge^r V$.

One easily verifies that the shuffle product is associative and strictly commutative; specifically, $A_p \cdot B_q = B_q \cdot A_p \in \text{End } \wedge^{p+q} V$ with no plus-or-

minus signs. For example, for $p = 1$ and $q = 1$ one has

$$\begin{aligned} (A_1 \cdot B_1)(x_1 \wedge x_2) &= A_1 x_1 \wedge B_1 x_2 - A_1 x_2 \wedge B_1 x_1 \\ &= -B_1 x_2 \wedge A_1 x_1 + B_1 x_1 \wedge A_1 x_2 = (B_1 \cdot A_1)(-x_2 \wedge x_1) \\ &= (B_1 \cdot A_1)(x_1 \wedge x_2), \end{aligned}$$

hence $A_1 \cdot B_1 = B_1 \cdot A_1 \in \text{End } \wedge^2 V$. The algebra $\Pi_p \text{End } \wedge^p V$ has a unique (two-sided) unit element with respect to the shuffle product, whose only nonzero component is the identity endomorphism I_0 of $\wedge^0 V$.

For any endomorphism A of V itself and any $p \geq 0$ there is a well-defined element $A_p \in \text{End } \wedge^p V$ such that

$$A_p(x_1 \wedge \dots \wedge x_p) = Ax_1 \wedge \dots \wedge Ax_p$$

for any $x_1 \wedge \dots \wedge x_p \in \wedge^p V$; in particular $A_1 = A$. Observe that the p -fold shuffle product $A^{\cdot p} = A \cdot \dots \cdot A$ is defined by

$$A^{\cdot p}(x_1 \wedge \dots \wedge x_p) = \sum_{\pi} \varepsilon_{\pi} Ax_{\pi_1} \wedge \dots \wedge Ax_{\pi_p},$$

the summation extending overall $p!$ permutations π of $\{1, \dots, p\}$. Since $\varepsilon_{\pi} Ax_{\pi_1} \wedge \dots \wedge Ax_{\pi_p} = Ax_1 \wedge \dots \wedge Ax_p$ for each permutation π it follows that $A^{\cdot p} = p! A_p$. For this reason A_p can reasonably be written $\frac{1}{p!} A^{\cdot p}$, without

requiring the ground ring to contain the element $\frac{1}{p!}$. Thus the direct product

of the elements $A_p \left(= \frac{1}{p!} A^{\cdot p} \right)$ over all $p \geq 0$ is essentially an exponential $e^{\cdot A} \in \Pi_p \text{End } \wedge^p V$. One easily verifies that $e^{\cdot A} \cdot e^{\cdot (-A)} = I_0 = e^{\cdot (-A)} \cdot e^{\cdot A}$, where $I_0 \in \text{End } \wedge^0 V$ represents the unit element in $\Pi_p \text{End } \wedge^p V$ with respect to the shuffle product.

For each $p \geq 0$ the p -fold shuffle product $I^{\cdot p}$ of the identity endomorphism $I \in \text{End } V$ satisfies $\frac{1}{p!} I^{\cdot p} = I_p$, where I_p is the identity endomorphism in $\text{End } \wedge^p V$. Hence $e^{\cdot I}$ is precisely the two-sided unit element \mathbf{I} of $\Pi_p \text{End } \wedge^p V$ with respect to the composition product. Since

$$e^{\cdot I} \cdot e^{\cdot (-I)} = I_0 = e^{\cdot (-I)} \cdot e^{\cdot I},$$

where $I_0 \in \text{End } \wedge^0 V$ represents the unit element with respect to the shuffle product, one can therefore define an invertible map α of $\Pi_p \text{End } \wedge^p V$ into itself by letting $\alpha \mathbf{A} \in \Pi_p \text{End } \wedge^p V$ be the shuffle product $e^{\cdot I} \cdot \mathbf{A}$ for any $\mathbf{A} \in \Pi_p \text{End } \wedge^p V$; the inverse α^{-1} of α is given by $\alpha^{-1} \mathbf{A} = e^{\cdot (-I)} \cdot \mathbf{A}$.

1.1 *Definition*: The *third product* of any two elements \mathbf{A} and \mathbf{B} of $\Pi_p \text{End } \wedge^p V$ is given by $\mathbf{A} \times \mathbf{B} = \alpha^{-1}((\alpha\mathbf{A})(\alpha\mathbf{B})) \in \Pi_p \text{End } \wedge^p V$, where $(\alpha\mathbf{A})(\alpha\mathbf{B})$ is the composition product of the shuffle products $\alpha\mathbf{A} = e \cdot I \cdot \mathbf{A}$ and $\alpha\mathbf{B} = e \cdot I \cdot \mathbf{B}$.

Since the composition product is associative the third product is trivially associative. Furthermore, if $I_0 \in \text{End } \wedge^0 V$ represents the unit element in $\Pi_p \text{End } \wedge^p V$ with respect to the shuffle product one has

$$I_0 \times \mathbf{A} = \alpha^{-1}((\alpha I_0)(\alpha\mathbf{A})) = \alpha^{-1}((e \cdot I)(\alpha\mathbf{A})) = \alpha^{-1}(\mathbf{I}(\alpha\mathbf{A})) = \alpha^{-1}(\alpha\mathbf{A}) = \mathbf{A}$$

and similarly $\mathbf{A} \times I_0 = \mathbf{A}$ for any $\mathbf{A} \in \Pi_p \text{End } \wedge^p V$; that is, I_0 is also the unit element of $\Pi_p \text{End } \wedge^p V$ with respect to the third product. The rationale for introducing the third product appears in the next section.

2. THE TRACE

We now specialize the arbitrary R -module V of the preceding section.

2.1 *Definition*: A module V over a commutative ring R with unit is *traceable* of rank $n > 0$ if and only if $\text{End } \wedge^n V$ is a free R -module of rank one.

If $\wedge^n V$ is itself free of rank one then V is clearly traceable of rank n . However, $\text{End } \wedge^n V$ can be free of rank one with no such condition on $\wedge^n V$. For example, let X be any paracompact hausdorff space, let R be the ring $C(X)$ of continuous real-valued functions on X , and let V be the $C(X)$ -module of continuous sections of a real n -plane bundle ξ over X ; then V is traceable of rank n . However $\wedge^n V$ is itself free of rank one if and only if ξ is orientable.

Flanders [1] showed for any module V over a commutative ring with unit that if $\wedge^n V$ is free of rank one then $\wedge^p V = 0$ for every $p > n$; a similar argument shows that if V is traceable of rank $n > 0$ then $\text{End } \wedge^p V = 0$ for every $p > n$. Thus if V is traceable of rank $n > 0$ there is no distinction between the direct product $\Pi_p \text{End } \wedge^p V$ and the direct sum $\amalg_p \text{End } \wedge^p V$. Consequently the third product of Definition 1.1 can be regarded as a product in $\amalg_p \text{End } \wedge^p V$ whenever V is traceable.

If V is traceable of rank n then every element of $\text{End } \wedge^n V$ is scalar multiplication by a unique element of the commutative ground ring R with unit. For example, for any $\mathbf{A} \in \amalg_p \text{End } \wedge^p V$ and each $p = 0, \dots, n$ let