

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 32 (1986)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ON CONSECUTIVE VALUES OF THE LIOUVILLE FUNCTION  
**Autor:** Hildebrand, Adolf  
**Kapitel:** 4. Proof of the Theorem, conclusion  
**DOI:** <https://doi.org/10.5169/seals-55088>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 10.12.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

then

$$\lambda\left(\frac{N}{5} \pm 1\right) = \frac{\lambda(N \pm 5)}{\lambda(5)} = 1 = -\lambda(N) = \lambda\left(\frac{N}{5}\right),$$

and  $N/5 = 4n/5 \in I_n$  is good. We may therefore suppose that at least one of values  $\lambda(N+5)$  and  $\lambda(N-5)$  equals 1.

For definiteness we shall assume  $\lambda(N+5) = 1$ ; the other case is treated in exactly the same way.

If  $\lambda(N+3) = 1$  or  $\lambda(N+6) = 1$ , then  $N+4 \in I_n$  or  $N+5 \in I_n$  is good. But in the remaining case

$$\lambda(N+3) = \lambda(N+6) = -1$$

we have

$$\lambda\left(\frac{N}{3}\right) = \lambda\left(\frac{N}{3} + 1\right) = \lambda\left(\frac{N}{3} + 2\right) = 1,$$

so that  $(N+3)/3 \in I_n$  is good.

Thus (3) implies the existence of a good integer in the interval (4), as we had to show.

#### 4. PROOF OF THE THEOREM, CONCLUSION

So far we have proved that (1) has infinitely many solutions in the cases  $\varepsilon_1 = \varepsilon_2 = \varepsilon_3 = 1$  and  $\varepsilon_1 = \varepsilon_2 = \varepsilon_3 = -1$ . But this obviously implies that for each of the triples  $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (1, 1, -1), (-1, -1, 1), (1, -1, -1)$  and  $(-1, 1, 1)$  there are also infinitely many solutions to (1). It remains therefore to consider the triples  $(1, -1, 1)$  and  $(-1, 1, -1)$ . Since the arguments in both cases are the same (with  $+1$  and  $-1$  interchanged), we shall confine ourselves to the case  $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (1, -1, 1)$ . Accordingly, we call  $n \geq 2$  good, whenever

$$\lambda(n+1) = \lambda(n-1) = 1, \quad \lambda(n) = -1.$$

We have to show that there are infinitely many such  $n$ .

Suppose, to get a contradiction, that there are only finitely many good integers, all of them  $\leq N_0$ , say. Suppose further that

$$(5) \quad \lambda(n) = 1(m_0 \leq n \leq n_0)$$

holds for some integers  $n_0 > m_0 \geq 2N_0$ . We shall show that then

$$(6) \quad \lambda(n) = 1(m_i \leq n \leq n_i)$$

holds for all  $i \geq 0$ , where  $m_i$  and  $n_i$  are defined inductively by

$$(7) \quad m_{i+1} = \left\lceil \frac{3m_i + 1}{2} \right\rceil, \quad n_{i+1} = \left\lceil \frac{3n_i}{2} \right\rceil (i \geq 0).$$

This will easily lead to the desired contradiction.

By our assumption (5), (6) holds for  $i = 0$ . Assume now that (6) does not hold for all  $i \geq 0$ , and let  $i \geq 0$  be minimal such that (6) holds for  $i$  and fails for  $i + 1$ . Thus, for some  $n \in [m_{i+1}, n_{i+1}]$ , which we shall fix, we have  $\lambda(n) = -1$ . Write

$$(8) \quad 2n = 3n' + \theta (\theta \in \{0, 1, -1\}).$$

From (7) we get

$$3m_i \leq 2m_{i+1} \leq 2n \leq 2n_{i+1} \leq 3n_i,$$

so that

$$m_i \leq n' \leq n_i,$$

and hence by (6) (which we assumed to hold for  $i$ )

$$\lambda(3n') = -\lambda(n') = -1.$$

Since, by our assumption  $\lambda(n) = -1$ ,

$$\lambda(2n) = -\lambda(n) = 1,$$

we cannot have  $\theta = 0$  in (8). The arguments in the cases  $\theta = \pm 1$  being identical, we shall henceforth assume that (8) holds with  $\theta = 1$ .

We must have

$$\lambda(2(n-1)) = \lambda(3n'-1) = -1,$$

since otherwise  $3n'$  would be good and

$$3n' \geq 3m_i \geq 3m_0 > N_0,$$

in contradiction to our assumption. Also, since

$$m_i \leq n' + 1 = \frac{2}{3}(n+1) \leq \left\lceil \frac{2}{3}(n_{i+1}+1) \right\rceil = \left\lceil \frac{2}{3} \left( \left\lceil \frac{3n_i}{2} \right\rceil + 1 \right) \right\rceil \leq n_i,$$

we have by (6)

$$\lambda(2(n+1)) = \lambda(3(n'+1)) = -\lambda(n'+1) = -1.$$

These two identities imply

$$\lambda(n \pm 1) = -\lambda(2(n \pm 1)) = 1,$$

and since  $\lambda(n) = -1$ , we conclude that  $n(>N_0)$  is good and therefore arrive at a contradiction.

Thus (5) (with  $n_0 > m_0 \geq 2N_0$ ) implies (6) for all  $i \geq 0$ . To derive from this the desired contradiction, we suppose first that (5) holds for some  $n_0 > m_0 \geq 2N_0$  satisfying

$$(9) \quad n_0 - m_0 \geq 3.$$

In other words, we suppose (for the moment) that there exist four consecutive integers  $n \geq 2N_0$ , for which  $\lambda(n) = 1$ . Putting  $d_i = n_i - m_i$ , we have, by the recursion formulae (7),

$$d_{i+1} \geq \frac{3}{2} d_i - 1 = \frac{3}{2} d_i \left(1 - \frac{2}{3d_i}\right) \quad (i \geq 0).$$

Taking into account (9), we obtain by induction in turn

$$d_i \geq 3 \quad (i \geq 0),$$

$$d_i \geq 3 \left(\frac{7}{6}\right)^i \quad (i \geq 0),$$

and finally

$$d_i \geq \left(\frac{3}{2}\right)^i \prod_{j=0}^i \left(1 - \frac{2}{3d_j}\right) \geq C \left(\frac{3}{2}\right)^i \quad (i \geq 0),$$

where

$$C = \prod_{j \geq 0} \left(1 - \frac{2}{9} \left(\frac{6}{7}\right)^j\right) > 0.$$

Since on the other hand by (7)

$$d_i \leq n_i \leq \left(\frac{3}{2}\right)^i n_0 \quad (i \geq 0),$$

we see from (6), that there are arbitrary large values of  $x$ , such that  $\lambda(n)$  is constant in the interval  $[x(1-\varepsilon), x]$ , where  $\varepsilon = C/n_0$ . But this is impossible since, for  $x$  sufficiently large, every such interval contains integers  $n$  and  $n'$  of the form

$$n = 4^a 9^b, \quad n' = 2 \cdot 4^c 9^d (a, b, c, d \in \mathbf{N}),$$

for which

$$\lambda(n) = 1, \quad \lambda(n') = -1.$$

We therefore have obtained the desired contradiction under the assumption that there exist four consecutive integers  $n \geq 2N_0$ , for which  $\lambda(n) = 1$ . By the part of the theorem already proved, there exist at least three such integers. Therefore (5) holds for some  $m_0 > 2N_0$  with  $n_0 = m_0 + 2$ , and we may now assume that

$$\lambda(m_0 - 1) = \lambda(m_0 + 3) = -1.$$

If  $m_0$  is odd, then this implies

$$\lambda\left(\frac{m_0 - 1}{2}\right) = \lambda\left(\frac{m_0 + 3}{2}\right) = 1, \quad \lambda\left(\frac{m_0 + 1}{2}\right) = -1,$$

so that  $(m_0 + 1)/2 > N_0$  is good, in contradiction to our assumption. But if  $m_0$  is even, then defining  $m_1$  and  $n_1$  by (7), (6) holds for  $i = 1$ , and we have

$$m_1 \geq 2N_0, \quad n_1 - m_1 = \frac{3(m_0 + 2)}{2} - \frac{3m_0}{2} = 3.$$

Thus we are back in the case already treated.

By contradiction, we therefore conclude that (1) has infinitely many solutions for  $(\varepsilon_1, \varepsilon_2, \varepsilon_3) = (1, -1, 1)$ , and the proof of the theorem is complete.

## 5. CONCLUDING REMARKS

In the foregoing proof, the relevant property of the Liouville function was that  $\lambda(n)$  is completely multiplicative and assumes only the values  $\pm 1$ . Besides this, we used only the fact that  $\lambda(2) = \lambda(3) = \lambda(5) = -1$  and (in the proof of the lemma)

$$\lambda(14) = \lambda(16) = 1, \quad \lambda(29) = \lambda(31) = -1.$$

The proof, as it stands, works for any completely multiplicative function  $f(n) = \pm 1$  with these properties. By suitably modifying the proof, it is possible to cover other classes of multiplicative functions as well.