

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 32 (1986)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: INTRODUCTION TO MICROLOCAL ANALYSIS
Autor: Kashiwara, Masaki
Kapitel: §1. Systems of Differential Equations (See [O], [Bj])
DOI: <https://doi.org/10.5169/seals-55089>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

0.5. In the situation of § 0.2, $u(x) = c_0(x)f(x)^s + \dots$ satisfies $\text{supp } \text{sp}(u(x)) = \{\pm df(x)\}$. Therefore $P_m(x, df)$ must be zero if $P(x, \partial)u(x) = 0$. In fact otherwise the bijectivity of $P: \mathcal{C}_M \rightarrow \mathcal{C}_M$ implies $\text{sp}(u) = 0$.

0.6. Such a method of studying functions or differential equations locally on the cotangent bundle is called microlocal analysis. After Sato's discovery of microfunctions, microlocal analysis was studied intensively in Sato-Kawai-Kashiwara [SKK].

Also L. Hörmander [H] worked in the C^∞ -case. Since then, microlocal analysis has been one of the most fundamental tools in the theory of linear partial differential equations.

§ 1. SYSTEMS OF DIFFERENTIAL EQUATIONS (See [O], [Bj])

1.1. Let X be a complex manifold. A system of linear differential equations can be written in the form

$$(1.1.1) \quad \sum_{j=1}^{N_0} P_{ij}(x, \partial)u_j = 0, \quad i = 1, 2, \dots, N_1.$$

Here u_1, \dots, u_{N_0} denote unknown functions and the $P_{ij}(x, \partial)$ are differential operators on X . The holomorphic function solutions of (1.1.1) are simply the kernel of the homomorphism

$$(1.1.2) \quad P: \mathcal{O}_X^{N_0} \rightarrow \mathcal{O}_X^{N_1}$$

which assigns (v_1, \dots, v_{N_1}) to (u_1, \dots, u_{N_0}) , where $v_i = \sum_j P_{ij}(x, \partial)u_j$.

Let us denote by \mathcal{D}_X the ring of differential operators with holomorphic coefficients. Then

$$(1.1.3) \quad P: \mathcal{D}_X^{N_1} \rightarrow \mathcal{D}_X^{N_0}$$

given by (Q_1, \dots, Q_{N_1}) to $(\Sigma Q_i P_{i1}, \dots, \Sigma Q_i P_{iN_0})$ is a left \mathcal{D}_X -linear homomorphism. If we denote by \mathcal{M} the cokernel of (1.1.3), then \mathcal{M} becomes a left \mathcal{D}_X -module and $\mathcal{H}om_{\mathcal{D}_X}(\mathcal{M}, \mathcal{O}_X)$ is the kernel of (1.1.2). This means that the set of holomorphic solutions to $Pu = 0$ depends only on \mathcal{M} .

For this reason we mean by a system of linear differential equations a left \mathcal{D}_X -module.

1.2. Let us take a local coordinate system (x_1, \dots, x_n) of X . Then any differential operator P can be written in the form

$$(1.2.1) \quad P(x, \partial) = \sum_{\alpha \in \mathbb{N}^n} a_\alpha(x) \partial^\alpha$$

where $\partial^\alpha = \partial^{|\alpha|}/\partial x_1^{\alpha_1} \dots \partial x_n^{\alpha_n}$, $|\alpha| = \alpha_1 + \dots + \alpha_n$ and the $a_\alpha(x)$ are holomorphic functions. For $j \in \mathbb{N}$, we set

$$P_j(x, \xi) = \sum_{|\alpha|=j} a_\alpha(x) \xi^\alpha,$$

where $\xi^\alpha = \xi_1^{\alpha_1} \dots \xi_n^{\alpha_n}$, and we call $\{P_j(x, \xi)\}$ the *total symbol* of P . The largest m such that $P_m \neq 0$ is called the *order* of P and P_m is called the *principal symbol* of P and denoted by $\sigma(P)$.

Let us denote by T^*X the cotangent bundle of X , and let

$$(x_1, \dots, x_n; \xi_1, \dots, \xi_n)$$

be the associated coordinates of T^*X . It is a classical result that if we consider $\sigma(P)$ as a function on T^*X , then this does not depend on our choice of the local coordinate system (x_1, \dots, x_n) .

1.3. Let M be a real analytic manifold, and X its complexification, e.g., $M = \mathbf{R}^n \subset X = \mathbf{C}^n$. Let P be a differential operator on X . When $\sigma(P)(x, \xi) \neq 0$ for $(x, \xi) \in \mathbf{R}^n \times (\mathbf{R}^n \setminus \{0\})$, P is called an elliptic differential operator. In this case, we have the following result.

PROPOSITION 1.3.1. *If u is a hyperfunction (or distribution) on M and Pu is real analytic, then u is real analytic. More precisely if we denote by \mathcal{A} the sheaf of real analytic functions on M and by \mathcal{B} (resp. $\mathcal{D}\mathcal{b}$) the sheaf of hyperfunctions (resp. distributions) on M , then $P: \mathcal{B}/\mathcal{A} \rightarrow \mathcal{B}/\mathcal{A}$ (resp. $P: \mathcal{D}\mathcal{b}/\mathcal{A} \rightarrow \mathcal{D}\mathcal{b}/\mathcal{A}$) is a sheaf isomorphism.*

This suggests that if $\sigma(P)(x, \xi) \neq 0$, we can consider the inverse P^{-1} in a certain sense. Since (x, ξ) is a point of the cotangent bundle, P^{-1} is attached to the cotangent bundle.

In fact, as we shall see in the sequel, we can construct a sheaf of rings \mathcal{E}_X on T^*X such that $\mathcal{D}_X \subset \pi_* \mathcal{E}_X$, where π is the canonical projection $T^*X \rightarrow X$. Moreover if $P \in \mathcal{D}_X$ satisfies $\sigma(P)(x, \xi) \neq 0$ at a point $(x, \xi) \in T^*X$, then P^{-1} exists as a section of \mathcal{E}_X on a neighborhood of (x, ξ) .

This can be compared to the analogous phenomena for polynomial rings, as shown in the following table.