

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 32 (1986)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: INTRODUCTION TO MICROLOCAL ANALYSIS
Autor: Kashiwara, Masaki
Kapitel: §2. Micro-differential Operators (See [SKK], [Bj], [S], [K2])
DOI: <https://doi.org/10.5169/seals-55089>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

$\mathbb{C}[x_1, \dots, x_n]$	\mathcal{D}_X
\mathbb{C}^n	T^*X
the sheaf $\mathcal{O}_{\mathbb{C}^n}$ of holomorphic functions	\mathcal{E}_X

§ 2. MICRO-DIFFERENTIAL OPERATORS (See [SKK], [Bj], [S], [K2])

2.1. Let X be an n -dimensional complex manifold and let $\pi_X: T^*X \rightarrow X$ be the cotangent bundle of X . Let us take a local coordinate system (x_1, \dots, x_n) of X and the associated coordinates $(x_1, \dots, x_n; \xi_1, \dots, \xi_n)$ of T^*X . For a differential operator P , let $\{P_j(x, \xi)\}$ be the total symbol of P as in § 1.2. We sometimes write $P = \sum P_j(x, \partial)$.

Let $Q = \sum Q_j(x, \partial)$ be another differential operator. Set $S = P + Q$ and $R = PQ$. Then the total symbols $\{S_j\}$ and $\{R_j\}$ of R and S are given explicitly by

$$(2.1.1) \quad S_j = P_j + Q_j$$

$$(2.1.2) \quad R_l = \sum_{\substack{l=j+k-|\alpha| \\ \alpha \in \mathbb{N}^n}} \frac{1}{\alpha!} (\partial_\xi^\alpha P_j)(\partial_x^\alpha Q_k)$$

where $\partial_\xi^\alpha = (\partial/\partial\xi_1)^{\alpha_1} \dots (\partial/\partial\xi_n)^{\alpha_n}$ and $\partial_x^\alpha = (\partial/\partial x_1)^{\alpha_1} \dots (\partial/\partial x_n)^{\alpha_n}$.

The total symbol $\{P_j(x, \xi)\}$ of a differential operator behaves as follows under coordinate transformations. Let (x_1, \dots, x_n) and $(\tilde{x}_1, \dots, \tilde{x}_n)$ be two local coordinate systems. Let (ξ_1, \dots, ξ_n) and $(\tilde{\xi}_1, \dots, \tilde{\xi}_n)$ be related by

$$\xi_k = \sum_j \tilde{\xi}_j \cdot \frac{\partial \tilde{x}_j}{\partial x_k}$$

i.e. $(x_1, \dots, x_n; \xi_1, \dots, \xi_n)$ and $(\tilde{x}_1, \dots, \tilde{x}_n; \tilde{\xi}_1, \dots, \tilde{\xi}_n)$ are the associated local coordinate systems of the cotangent bundle T^*X . Let P be a differential operator on X and let $\{P_j(x, \xi)\}$ and $\{\tilde{P}_j(\tilde{x}, \tilde{\xi})\}$ be the total symbols of P with respect to the local coordinate systems (x_1, \dots, x_n) and $(\tilde{x}_1, \dots, \tilde{x}_n)$, respectively. Then one has

$$(2.1.3) \quad \begin{aligned} & \tilde{P}_l(\tilde{x}, \tilde{\xi}) \\ &= \sum_{v, \alpha_1, \dots, \alpha_v} \frac{1}{v! \alpha_1! \dots \alpha_v!} \langle \tilde{\xi}, \partial_x^{\alpha_1} \tilde{x} \rangle \dots \langle \tilde{\xi}, \partial_x^{\alpha_v} \tilde{x} \rangle \partial_\xi^{\alpha_1 + \dots + \alpha_v} P_j(x, \xi). \end{aligned}$$

Here the indices run over $j \in \mathbf{Z}$, $v \in \mathbf{N}$, $\alpha_1, \dots, \alpha_v \in \mathbf{N}^n$ such that $|\alpha_1|, \dots, |\alpha_v| \geq 2$ and $l = j + v - |\alpha_1| - \dots - |\alpha_v|$. For $\beta \in \mathbf{N}^n$, $\langle \tilde{\xi}, \partial_x^\beta \tilde{x} \rangle$ denotes $\sum_j \tilde{\xi}_j \partial_x^\beta \tilde{x}_j$.

2.2. The total symbol $\{P_j(x, \xi)\}$ of a differential operator is a polynomial in ξ . We shall define microdifferential operators by admitting P_j to be holomorphic in ξ .

For $\lambda \in \mathbf{C}$, let $\mathcal{O}_{T^*X}(\lambda)$ be the sheaf of homogeneous holomorphic functions of degree λ on T^*X , i.e., holomorphic functions $f(x, \xi)$ satisfying

$$(\sum \xi_j \partial/\partial \xi_j - \lambda) f(x, \xi) = 0.$$

Definition 2.2.1. For $\lambda \in \mathbf{C}$ we define the sheaf $\mathcal{E}_X(\lambda)$ on T^*X by

$$\Omega \mapsto \{(P_{\lambda-j}(x, \xi))_{j \in \mathbf{N}} ; P_{\lambda-j} \in \Gamma(\Omega; \mathcal{O}_{T^*X}(\lambda-j)) \text{ and satisfies the following conditions (2.2.1)}\}$$

(2.2.1) for any compact subset K of Ω , there exists a $C_K > 0$ such that

$$\sup_K |P_{\lambda-j}| \leq C_K^{-j} (j!) \quad \text{for all } j > 0.$$

Remark. The growth condition (2.2.1) can be explained as follows. For a differential operator $P = \sum P_j(x, \partial)$, we have

$$P(x, \partial) (\langle x, \xi \rangle + p)^\mu = \sum P_j(x, \xi) \frac{\Gamma(\mu)}{\Gamma(\mu-j+1)} (\langle x, \xi \rangle + p)^{\mu-j}.$$

For $P = (P_{\lambda-j}(x, \xi)) \in \mathcal{E}(\lambda)$ we set, by analogy

$$P(\langle x, \xi \rangle + p)^\mu = \sum_j P_{\lambda-j}(x, \xi) \frac{\Gamma(\mu)}{\Gamma(\mu-\lambda+j+1)} (\langle x, \xi \rangle + p)^{\mu-\lambda+j}.$$

Then the growth condition (2.2.1) is simply the condition that the right hand side converges when $0 < |\langle x, \xi \rangle + p| \ll 1$.

Now, we have the following

PROPOSITION 2.2.2 ([SKK], Chap. II, § 1, [Bj] Chap. IV, § 1).

- (0) $\mathcal{E}_X(\lambda)$ contains $\mathcal{E}_X(\lambda-m)$ as a subsheaf for $m \in \mathbf{N}$.
- (1) Patching by rule (2.1.3) under coordinate transformations, $\mathcal{E}_X(\lambda)$ becomes a sheaf defined globally on T^*X .

- (2) By rule (2.1.1), $\mathcal{E}_X(\lambda)$ is a sheaf of \mathbf{C} -vector space on T^*X .
(3) By rule (2.1.2), we can define the “product” homomorphism:

$$\mathcal{E}_X(\lambda) \underset{\mathbf{C}}{\otimes} \mathcal{E}_X(\mu) \rightarrow \mathcal{E}_X(\lambda + \mu),$$

which satisfies the associative law.

- (4) In particular, $\mathcal{E}_X(0)$ and $\mathcal{E}_X = \bigcup_{m \in \mathbf{Z}} \mathcal{E}_X(m)$ become sheaves of (non commutative) rings on T^*X , with a unit.

The unit is given by $(P_j(x, \xi))$ with $P_j = 1$ for $j = 0$ and $P_j = 0$ for $j \neq 0$.

We define the homomorphism

$$\sigma_\lambda: \mathcal{E}_X(\lambda) \rightarrow \mathcal{O}_{T^*X}(\lambda)$$

by

$$(P_{\lambda-j}) \mapsto P_\lambda.$$

Then, σ_λ is a well-defined homomorphism on T^*X (i.e. compatible with coordinate transformation) and we have an exact sequence

$$0 \rightarrow \mathcal{E}_X(\lambda - 1) \rightarrow \mathcal{E}_X(\lambda) \xrightarrow{\sigma_\lambda} \mathcal{O}_{T^*X}(\lambda) \rightarrow 0.$$

Now we have the following proposition, which says that the ring \mathcal{E}_X is a kind of localization of \mathcal{D}_X .

PROPOSITION 2.2.3.

- (1) For $P \in \mathcal{E}(\lambda)$ and $Q \in \mathcal{E}(\mu)$, we have $\sigma_{\lambda+\mu}(PQ) = \sigma_\lambda(P)\sigma_\mu(Q)$.
(2) ([SKK] Chap. II, Thm. 2.1.1) If $P \in \mathcal{E}(\lambda)$ satisfies $\sigma_\lambda(P)(q) \neq 0$ at $q \in T^*X$, then there exists $Q \in \mathcal{E}(-\lambda)$ such that $PQ = QP = 1$.

The relations between \mathcal{E}_X and \mathcal{D}_X are summarized in the following theorem.

THEOREM 2.2.4 ([SKK], Chap. II, § 3).

- (i) \mathcal{E}_X contains $\pi^{-1}\mathcal{D}_X$ as a subring and is flat over $\pi^{-1}\mathcal{D}_X$.
(ii) $\mathcal{E}_X|_{T_X^*X} \simeq \mathcal{D}_X$, where T_X^*X is the zero section of T^*X .
(iii) For a coherent \mathcal{D}_X -module \mathcal{M} , the characteristic variety of \mathcal{M} coincides with the support of $\mathcal{E}_X \underset{\pi_X^{-1}\mathcal{D}_X}{\otimes} \mathcal{M}$.