

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 32 (1986)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: INTRODUCTION TO MICROLOCAL ANALYSIS
Autor: Kashiwara, Masaki
Kapitel: §4. Variants of E (See [SKK], [Bj], [S])
DOI: <https://doi.org/10.5169/seals-55089>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

§ 3. THE ALGEBRAIC PROPERTIES OF \mathcal{E} (See [SKK], [Bj])

3.1. In the preceding section, we introduced the notion of micro-differential operators. The ring \mathcal{E} of micro-differential operators has nice algebraic properties similar to those of the ring of holomorphic functions.

Let us recall some definitions of finiteness properties.

Definition 3.1.1. Let \mathcal{A} be a sheaf of rings on a topological space S .

- (1) An \mathcal{A} -module \mathcal{M} is called *of finite type* (resp. *of finite presentation*) if for any point $x \in X$ there exists a neighborhood U and an exact sequence $0 \leftarrow \mathcal{M}|_U \leftarrow \mathcal{A}^p|_U$ (resp. $0 \leftarrow \mathcal{M}|_U \leftarrow \mathcal{A}^p|_U \leftarrow \mathcal{A}^q|_U$).
- (2) \mathcal{M} is called *pseudo-coherent*, if any submodule of finite type defined on an open subset is of finite presentation. If \mathcal{M} is pseudo-coherent and of finite type, then \mathcal{M} is called *coherent*.
- (3) \mathcal{M} is called *Noetherian* if \mathcal{M} satisfies the following properties:
 - (a) \mathcal{M} is coherent.
 - (b) For any $x \in X$, \mathcal{M}_x is a Noetherian \mathcal{A}_x -module (i.e. any increasing sequence of \mathcal{A}_x -submodules is stationary).
 - (c) For any open subset U , any increasing sequence of coherent $(\mathcal{A}|_U)$ -submodules of $\mathcal{M}|_U$ is locally stationary.

As for the sheaf of holomorphic functions, we have

THEOREM 3.1.1 ([SKK] Chap. II, Thm. 3.4.1, Prop. 3.2.7). *Let \mathring{T}^*X denote the complement of the zero section in T^*X .*

- (1) \mathcal{E}_X and $\mathcal{E}_X(0)$ are Noetherian rings on T^*X .
- (2) \mathcal{E}_X is flat over $\pi^{-1}\mathcal{D}_X$.
- (3) $\mathcal{E}_X(\lambda)|_{\mathring{T}^*X}$ is a Noetherian $\mathcal{E}_X(0)|_{\mathring{T}^*X}$ -module.
- (4) For $p \in T^*X$, $\mathcal{E}_X(0)_p$ is a local ring with the residual field \mathbf{C} .
- (5) A coherent \mathcal{E}_X -module is pseudo-coherent over $\mathcal{E}_X(0)$.

§ 4. VARIANTS OF \mathcal{E} (See [SKK], [Bj], [S])

4.1. We have defined the sheaf of rings \mathcal{E} . However we can introduce other sheaves of rings, similar to \mathcal{E} , which makes the theory transparent.

4.2. The sheaf $\hat{\mathcal{E}} = \lim_{\leftarrow} \mathcal{E}/\mathcal{E}(-m)$ is called the sheaf of *formal micro-differential operators*. This is nothing but the sheaf similar to \mathcal{E} , obtained by dropping the growth condition (2.2.1).

4.3. We can define the sheaf \mathcal{E}^∞ of micro-differential operators of infinite order ([SKK]). For an open $\Omega \subset \mathbf{C}^n$, we set

$$\Gamma(\Omega; \mathcal{E}^\infty) = \{(p_j)_{j \in \mathbf{Z}} ; p_j \in \Gamma(\Omega; \mathcal{O}_{T^*X}(j))\}$$

satisfying the following conditions (4.3.1) and (4.3.2).

(4.3.1) For any compact set $K \subset \Omega$, there is a $C_K > 0$ such that $\sup_K |p_j| \leq C_K^{-j} (-j)! \text{ for } j < 0$.

(4.3.2) For any compact set $K \subset \Omega$ and any $\varepsilon > 0$, there exists a $C_{K,\varepsilon} > 0$ such that

$$\sup_K |p_j| \leq C_{K,\varepsilon} \frac{\varepsilon^j}{j!} \quad \text{for } j \geq 1.$$

4.4. We can also define the sheaf $\mathcal{E}^\mathbf{R}$ on T^*X by $\mathcal{H}^n(\mu_\Delta(\mathcal{O}_{X \times X}^{(0,n)}))$. (See [KS] Chap. II, [SKK]). Here $n = \dim X$, $\mathcal{O}_{X \times X}^{(0,n)}$ is the sheaf of holomorphic forms on $X \times X$ which are n -forms with respect to the second variable, and μ_Δ is the micro-localization with respect to the diagonal set of $X \times X$ (See [SKK] Chap. II for the details).

4.5. We have $\mathcal{E}_X \subset \mathcal{E}_X^\infty \subset \mathcal{E}_X^\mathbf{R}$, $\mathcal{E}_X \subset \hat{\mathcal{E}}_X$. Moreover, \mathcal{E}_X^∞ , $\mathcal{E}_X^\mathbf{R}$ and $\hat{\mathcal{E}}_X$ are faithfully flat over \mathcal{E}_X . The sheaf $\hat{\mathcal{E}}_X$ is Noetherian. The sheaf $\mathcal{E}_X^\mathbf{R}$ contains $\mathcal{E}_X(\lambda)$'s compatible with the multiplication.

4.6. If we denote by γ the projection map $T^*X \rightarrow T^*X/\mathbf{C}^*$, then $R^j \gamma_* \mathcal{E}^\mathbf{R} = 0$ for $j \neq 0$ and $\mathcal{E}^\infty = \gamma^{-1} \gamma_* \mathcal{E}^\mathbf{R}$.

4.7. In [SKK], \mathcal{E} , $\hat{\mathcal{E}}$, and \mathcal{E}^∞ are denoted by \mathcal{P}^f , $\hat{\mathcal{P}}$ and \mathcal{P} .

4.8. To explain the differences between \mathcal{E} , \mathcal{E}^∞ , $\mathcal{E}^\mathbf{R}$ and $\hat{\mathcal{E}}$, we shall take the following example. Let X be a complex manifold and Y a hypersurface of X . We shall take local coordinates (x_1, \dots, x_n) of X such that Y is given by $x_1 = 0$. The \mathcal{D}_X -module $\mathcal{D}_X/\mathcal{D}_X x_1 + \sum_{j>1} \mathcal{D}_X \partial_j$ is denoted by $\mathcal{B}_{Y|X}$. Set

$$\begin{aligned}\mathcal{C}_{Y|X} &= \mathcal{E}_X \underset{\pi^{-1}\mathcal{D}_X}{\otimes} \mathcal{B}_{Y|X}, \hat{\mathcal{C}}_{Y|X} = \hat{\mathcal{E}}_X \underset{\mathcal{E}_X}{\otimes} \mathcal{C}_{Y|X}, \\ \mathcal{C}_{Y|X}^\infty &= \mathcal{E}_X^\infty \underset{\mathcal{E}_X}{\otimes} \mathcal{C}_{Y|X} \quad \text{and} \quad \mathcal{C}_{Y|X}^R = \mathcal{E}_X^R \underset{\mathcal{E}_X}{\otimes} \mathcal{C}_{Y|X}.\end{aligned}$$

Then we have, setting $p = (0, dx_1)$, $x_0 = 0$

$$\begin{aligned}\mathcal{C}_{Y|X,p} &= \{a + b \log x_1; a \in \mathcal{O}_{X,x_0}[1/x_1], b \in \mathcal{O}_{X,x_0}\}/\mathcal{O}_{X,x_0} \\ &\cong (\mathcal{O}_{X,x_0}[1/x_1]/\mathcal{O}_{X,x_0}) \oplus \mathcal{O}_{X,x_0} \\ \hat{\mathcal{C}}_{Y|X,p} &= \{a + b \log x_1; a \in \mathcal{O}_{X,x_0}[1/x_1], b \in \hat{\mathcal{O}}_{X|Y,x_0}\}/\mathcal{O}_{X,x_0} \\ &\cong (\mathcal{O}_{X,x_0}[1/x_1]/\mathcal{O}_{X,x_0}) \oplus \hat{\mathcal{O}}_{X|Y,x_0}.\end{aligned}$$

Here $\hat{\mathcal{O}}_{X|Y} = \lim_{\leftarrow} \mathcal{O}_X/x_1^m \mathcal{O}_X$ is the sheaf of formal power series in the x_1 -direction.

$$\mathcal{C}_{Y|X,p}^\infty = \{a + b \log x_1; a \in (j_* j^{-1} \mathcal{O}_X)_{x_0}, b \in \mathcal{O}_{X,x_0}\}/\mathcal{O}_{X,x_0}$$

where j is the open embedding $X \setminus Y \hookrightarrow X$.

$$\mathcal{C}_{Y|X,p}^R = \varinjlim_U \mathcal{O}(U)/\mathcal{O}_{X,x_0}.$$

Here U ranges over the set of open subsets of the form

$$\{x \in X; |x| < \varepsilon, \operatorname{Re} x_1 < \varepsilon \operatorname{Im} x_1\}.$$

4.8. If we use \mathcal{E}_X^∞ , the structure of \mathcal{E} -modules becomes simpler. We just mention two theorems in this direction.

THEOREM 4.8.1 ([KK] Thm. 5.2.1). *Let \mathcal{M} be a holonomic \mathcal{E}_X -module. Then there exists a (unique) regular holonomic \mathcal{E}_X -module \mathcal{M}_{reg} such that*

$$\mathcal{E}_X^\infty \underset{\mathcal{E}_X}{\otimes} \mathcal{M} \cong \mathcal{E}_X^\infty \underset{\mathcal{E}_X}{\otimes} \mathcal{M}_{\text{reg}}.$$

THEOREM 4.8.2 ([SKK] Chap. II, Thm. 5.3.1). *Let X and Y be complex manifolds and let T_Y^*Y be the zero section of T^*Y . If \mathcal{M} is an $\mathcal{E}_{X \times Y}$ -module whose support is contained in $T^*X \times T_Y^*Y$, then there exists a (locally) coherent \mathcal{E}_X -module \mathcal{L} such that*

$$\mathcal{E}_{X \times Y}^\infty \underset{\mathcal{E}_{X \times Y}}{\otimes} \mathcal{M} \cong \mathcal{E}_{X \times Y}^\infty \underset{\mathcal{E}_{X \times Y}}{\otimes} (\mathcal{L} \hat{\otimes} \mathcal{O}_Y).$$

Here $\hat{\otimes}$ denotes the exterior tensor product. (See § 8).