

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 32 (1986)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: INTRODUCTION TO MICROLOCAL ANALYSIS
Autor: Kashiwara, Masaki
Kapitel: §7. Quantized Contact Transformations
DOI: <https://doi.org/10.5169/seals-55089>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

An analytic subset V of T^*X is called *involutive* if $f|_V = g|_V = 0$ implies $\{f, g\}|_V = 0$.

The following theorem exhibits a phenomenon which has no analogue in the commutative case.

THEOREM 6.3.2 ([G]). *Let \mathcal{M} be a coherent \mathcal{E}_X -module defined on an open subset Ω of T^*X and let \mathcal{L} be a $\mathcal{E}_X(0)|_\Omega$ -module which is a union of coherent $\mathcal{E}_X(0)$ -modules. Then $V = \{p \in \Omega; \mathcal{L} \text{ is not coherent over } \mathcal{E}_X(0) \text{ on any neighborhood of } p\}$ is an involutive analytic subset of Ω .*

COROLLARY 6.3.3 ([SKK] Chap. II, Theorem 5.3.2, [M]). *For any coherent \mathcal{E}_X -module \mathcal{M} , $\text{Supp } \mathcal{M}$ is involutive.*

Since any involutive subset has codimension less than or equal to $\dim X$, we have

COROLLARY 6.3.4. *The support of a coherent \mathcal{E}_X -module has codimension $\leq \dim X$.*

After some algebraic calculation, this implies

THEOREM 6.3.5 ([SKK] Chap. II, Theorem 5.3.5). *For any point $p \in T^*X$, $\mathcal{E}_{X,p}$ has a global cohomological dimension $\dim X$.*

6.4. An analytic subset Λ of T^*X is called *Lagrangean* if Λ is involutive and $\dim \Lambda = \dim X$. A coherent \mathcal{E}_X -module is called *holonomic* if its support is Lagrangean.

§ 7. QUANTIZED CONTACT TRANSFORMATIONS

7.1. In the previous section, we saw that the symplectic structure of T^*X is closely related to micro-differential operators via the relation of commutator and Poisson bracket. In this section, we shall explain another relation.

Definition 7.2.1. Let X and Y be complex manifolds of the same dimension. A morphism φ from an open subset U of T^*X to T^*Y is called a *homogeneous symplectic transformation* if $\varphi^*\theta_Y = \theta_X$.

We can easily see the following

(7.2.1) If φ is a homogeneous symplectic transformation, then φ is a local isomorphism and is compatible with the action of \mathbf{C}^* .

(7.2.2) Assume $Y = \mathbf{C}^n$ and let $(y_1, \dots, y_n; \eta_1, \dots, \eta_n)$ be the coordinates of T^*Y , so that $\theta_Y = \sum \eta_j dy_j$.

Set $p_j = \eta_j \circ \varphi$ and $q_j = y_j \circ \varphi$. Then we have

(7.2.3.1) $\{p_j, p_k\} = \{q_j, q_k\} = 0, \{p_j, q_k\} = \delta_{j,k}$ for $j, k = 1, \dots, n$.

(7.2.3.2) p_j is homogeneous of degree 1 and q_j is homogeneous of degree 0 with respect to the fiber coordinates.

(7.2.4) Conversely assume that functions $\{q_1, \dots, q_n, p_1, \dots, p_n\}$ on $U \subset T^*X$ satisfy (7.2.3.1) and (7.2.3.2). Then the map $\varphi: U \rightarrow T^*Y$, given by

$$U \ni x \mapsto (q_1(x), \dots, q_n(x); p_1(x), \dots, p_n(x)) \in T^*Y,$$

is a homogeneous symplectic transformation. We call $(q_1, \dots, q_n; p_1, \dots, p_n)$ a *homogeneous symplectic coordinate system*.

THEOREM 7.2.2 ([SKK] Chap. II § 3.2, [K2] § 2.4, [Bj] Chap. 4 § 6). *Let $\varphi: T^*X \supset U \rightarrow T^*Y$ be a homogeneous symplectic transformation, let p_X be a point of U and set $p_Y = \varphi(p_X)$. Then we have*

- (a) *There exists an open neighborhood U' of p_X and a \mathbf{C} -algebra isomorphism $\Phi: \varphi^{-1}\mathcal{E}_Y|_{U'} \xrightarrow{\sim} \mathcal{E}_X|_{U'}$ (we call (φ, Φ) a quantized contact transformation).*
- (b) *If $\Phi: \varphi^{-1}\mathcal{E}_Y \rightarrow \mathcal{E}_X|_U$ is a \mathbf{C} -algebra homomorphism then for any m, Φ gives an isomorphism $\varphi^{-1}\mathcal{E}_Y(m) \xrightarrow{\sim} \mathcal{E}_X(m)|_U$. Moreover the following diagram commutes:*

$$\begin{array}{ccc} \varphi^{-1}\mathcal{E}_Y(m) & \xrightarrow{\Phi} & \mathcal{E}_X(m)|_U \\ \downarrow \sigma_m & & \downarrow \sigma_m \\ \varphi^{-1}\mathcal{O}_{T^*Y}(m) & \xrightarrow{\Phi^*} & \mathcal{O}_{T^*X}(m)|_U \end{array}$$

- (c) *Let Φ and Φ' be two \mathbf{C} -algebra homomorphisms $\varphi^{-1}\mathcal{E}_Y \rightarrow \mathcal{E}_X|_U$.*

Then there exist $\lambda \in \mathbf{C}$, a neighborhood U' of p_X and $P \in \Gamma(U; \mathcal{E}_X(\lambda))$ such that $\sigma_\lambda(P)$ is invertible and

$$\Phi'(Q) = P\Phi(Q)P^{-1} \quad \text{for } Q \in \varphi^{-1}\mathcal{E}_Y|_{U'}.$$

Moreover λ is unique and P is unique up to constant multiple.

(d) Let $Y = \mathbf{C}^n$ and let U be an open subset of T^*X .

If $P_j \in \Gamma(U; \mathcal{E}_X(1))$ and $Q_j \in \Gamma(U; \mathcal{E}_X(0))$ ($1 \leq j \leq n$) satisfy

$$(7.2.5) \quad [P_j, P_k] = [Q_j, Q_k] = 0 \\ [P_j, Q_k] = \delta_{jk}$$

then there exists a unique quantized contact transformation (φ, Φ) such that

$$\varphi(p) = (\sigma_0(Q_1)(p), \dots, \sigma_0(Q_n)(p), \sigma_1(P_1)(p), \dots, \sigma_1(P_n)(p)),$$

and $\Phi(y_j) = Q_j, \Phi(\partial_{y_j}) = P_j$.

We call $\{Q_1, \dots, Q_n, P_1, \dots, P_n\}$ quantized canonical coordinates.

7.3. We shall give several examples of quantized contact transformations.

Example 7.3.1. If $P(\partial)$ is a constant coefficient micro-differential operator of order 1, then

$$(x_1 + [P, x_1], x_2 + [P, x_2], \dots, x_n + [P, x_n], \partial_{x_1}, \dots, \partial_{x_n})$$

gives quantized canonical coordinates.

Example 7.3.2. More generally if P is a micro-differential operator of order 1 and $\exp tH_{\sigma_1(P)}$ exists, then $\exp tP$ gives a quantized contact transformation Φ_t , by solving the equation $\frac{d}{dt} \Phi_t(Q) = [P, \Phi_t(Q)]$ with the initial condition $\Phi_t(Q) = Q$ for $t = 0$.

Example 7.3.3. (Paraboloidal transformation [K2] p. 36). Set $X = \mathbf{C}^{1+n} = \{(t, x) \in \mathbf{C} \times \mathbf{C}^n\}$,

$$\Omega = \{(t, x; \tau, \xi) \in T^*X; \tau \neq 0\}, G = \mathrm{Sp}(n; \mathbf{C})$$

$$= \{g \in \mathrm{GL}(2n; \mathbf{C}); {}^t g J g = J\} \quad \text{with} \quad J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

For $g = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in G$, let Ψ_g be the quantized contact transformation given by

$$\partial_x \mapsto \alpha \partial_x - \beta x \partial_t$$

$$x \mapsto \gamma \partial_x \partial_t^{-1} + \delta x$$

$$\partial_t \mapsto \partial_t$$

$$\begin{aligned} t \mapsto t + \frac{1}{2} \{ & \langle \partial_x, {}^t \gamma \alpha \partial_x \rangle \partial_t^{-2} + \langle \partial_x, {}^t \gamma \beta x \rangle \partial_t^{-1} \\ & + \langle {}^t \gamma \beta x, \partial_x \rangle \partial_t^{-1} + \langle x, {}^t \delta \beta x \rangle \}. \end{aligned}$$

Then we have $\Psi_{g_1} \Psi_{g_2} = \Psi_{g_1 g_2}$.

§ 8. FUNCTORIAL PROPERTIES OF MICRO-DIFFERENTIAL MODULES (See [SKK])

8.1. External Tensor Product.

Let X and Y be complex manifolds and let p_1 and p_2 be the projections $T^*(X \times Y) \rightarrow T^*X$ and $T^*(X \times Y) \rightarrow T^*Y$, respectively. Then $\mathcal{E}_{X \times Y}$ contains $p_1^{-1}\mathcal{E}_X \otimes p_2^{-1}\mathcal{E}_Y$ as a subring. For an \mathcal{E}_X -module \mathcal{M} and an \mathcal{E}_Y -module \mathcal{N} , we define the $\mathcal{E}_{X \times Y}$ -module $\mathcal{M} \hat{\otimes} \mathcal{N}$ by

$$(8.1.1) \quad \mathcal{M} \hat{\otimes} \mathcal{N} = \mathcal{E}_{X \times Y} \underset{\substack{p_1^{-1}\mathcal{E}_X \otimes p_2^{-1}\mathcal{E}_Y \\ \mathbf{C}}}{\otimes} (p_1^{-1}\mathcal{M} \underset{\mathbf{C}}{\otimes} p_2^{-1}\mathcal{N}).$$

Then one can easily see

PROPOSITION 8.1.1.

- (i) $\mathcal{M} \hat{\otimes} \mathcal{N}$ is an exact functor in \mathcal{M} and in \mathcal{N} and $\text{Supp}(\mathcal{M} \hat{\otimes} \mathcal{N}) = \text{Supp } \mathcal{M} \times \text{Supp } \mathcal{N}$.
- (ii) If \mathcal{M} is \mathcal{E}_X -coherent and \mathcal{N} is \mathcal{E}_Y -coherent, then $\mathcal{M} \hat{\otimes} \mathcal{N}$ is $\mathcal{E}_{X \times Y}$ -coherent.

8.2. For a complex submanifold Y of a complex manifold X of codimension l , the sheaf $\lim_{\substack{\rightarrow \\ m}} \mathcal{E}xt_{\mathcal{O}_X}^l(\mathcal{O}_X/\mathcal{J}^m, \mathcal{O}_X)$ has a natural structure of \mathcal{D}_X -module,

which is denoted by $\mathcal{B}_{Y|X}$. Here \mathcal{J} is the defining ideal of Y . The homomorphism $\mathcal{O}_Y \rightarrow \mathcal{E}xt_{\mathcal{O}_X}^l(\mathcal{O}_Y, \Omega_X^l) \rightarrow \Omega_X^l \otimes_{\mathcal{O}_X} \mathcal{B}_{Y|X}$ gives the canonical section $c(Y, X)$ of $\Omega_X^l \otimes_{\mathcal{O}_X} \mathcal{B}_{Y|X}$. If we take local coordinates (x_1, \dots, x_n) of X such that Y is defined by $x_1 = \dots = x_l = 0$, then we have