

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	32 (1986)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
 Artikel:	INTRODUCTION TO MICROLOCAL ANALYSIS
Autor:	Kashiwara, Masaki
Kapitel:	§10. Structure of Regular Holonomic E-Modules (See [SKK], [KK])
DOI:	https://doi.org/10.5169/seals-55089

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

9.4. Noting that any nowhere dense closed analytic subset of a Lagrangean variety is never involutive, Theorem 9.2.3 implies the following theorem.

THEOREM 9.4.1. *Let \mathcal{M} be a holonomic \mathcal{E}_X -module. Then the following conditions are equivalent.*

- (i) *There exists a Lagrangean subvariety Λ such that \mathcal{M} has regular singularities along Λ .*
- (ii) *For any involutive subvariety Λ which contains $\text{Supp } \mathcal{M}$, \mathcal{M} has regular singularities along Λ .*
- (iii) *There exists an open dense subset Ω of $\text{Supp } \mathcal{M}$ such that \mathcal{M} has regular singularities along $\text{Supp } \mathcal{M}$ on Ω .*

If these equivalent conditions are satisfied, we say that \mathcal{M} is a *regular holonomic \mathcal{E}_X -module*.

The following properties are almost immediate.

THEOREM 9.4.2.

- (i) *Let $0 \rightarrow \mathcal{M}' \rightarrow \mathcal{M} \rightarrow \mathcal{M}'' \rightarrow 0$ be an exact sequence of three coherent \mathcal{E}_X -modules. If two of them are regular holonomic then so is the third.*
- (ii) *If \mathcal{M} is regular holonomic, its dual \mathcal{M}^* is also regular holonomic.*

We just mention another analytic property of regular holonomic modules, which generalizes the fact that a formal solution of an ordinary differential equation with regular singularity converges.

THEOREM 9.4.3 ([KK] Theorem 6.1.3). *If \mathcal{M} and \mathcal{N} are regular holonomic \mathcal{E}_X -modules, then $\mathcal{E}\text{xt}_{\mathcal{E}_X}^j(\mathcal{M}, \mathcal{N}) \rightarrow \mathcal{E}\text{xt}_{\mathcal{E}_X}^j(\mathcal{M}, \hat{\mathcal{E}}_X \otimes_{\mathcal{E}_X} \mathcal{N})$ and $\mathcal{E}\text{xt}_{\mathcal{E}_X}^j(\mathcal{M}, \mathcal{N}) \rightarrow \mathcal{E}\text{xt}_{\mathcal{E}_X}^j(\mathcal{M}, \mathcal{E}_X^\infty \otimes_{\mathcal{E}_X} \mathcal{N})$ are isomorphisms.*

§ 10. STRUCTURE OF REGULAR HOLONOMIC \mathcal{E} -MODULES (See [SKK], [KK])

10.1. Let Λ be a Lagrangean submanifold of T^*X . We define \mathcal{J}_Λ and \mathcal{E}_Λ as in § 9.2.

Then $\mathcal{E}_\Lambda(-1) = \mathcal{E}_\Lambda \cdot \mathcal{E}(-1)$ is a two-sided ideal of \mathcal{E}_Λ and $\mathcal{E}_\Lambda/\mathcal{E}_\Lambda(-1)$ is a sheaf of rings which contains $\mathcal{O}_\Lambda(0) = \mathcal{E}(0)/\mathcal{J}_\Lambda(-1)$, the sheaf of homogeneous functions on Λ .

Let us take an invertible \mathcal{O}_Λ -module \mathcal{L} such that $\mathcal{L}^{\otimes 2} \cong \omega_\Lambda \otimes_{\mathcal{O}_X} \omega_X^{\otimes -1}$.

Such an \mathcal{L} exists at least locally. For $P = P_1(x, \partial) + P_0(x, \partial) + \dots \in \mathcal{J}$ we define, for $\phi \in \mathcal{O}_\Lambda$ and an invertible section s of \mathcal{L} ,

$$L(P)(\phi s) = \left\{ H_{P_1}(\phi) + \frac{1}{2} \phi \frac{L_{H_{P_1}}(s^{\otimes 2} \otimes dx)}{s^{\otimes 2} \otimes dx} + \left(P_0 - \frac{1}{2} \sum_{j=1}^n \frac{\partial^2 P_1}{\partial x_i \partial \xi_i} \right) \phi \right\} s.$$

Here $dx = dx_1 \wedge \dots \wedge dx_n \in \omega_X$ and $s^{\otimes 2} \otimes dx$ is regarded as a section of ω_Λ . The Lie derivative $L_{H_{P_1}}$ of H_{P_1} operates on ω_Λ as the first order differential operators so that $L_{H_{P_1}}(s^{\otimes 2} \otimes dx)$ is a section of ω_Λ and $L_{H_{P_1}}(s^{\otimes 2} \otimes dx)/s^{\otimes 2} \otimes dx$ is a function on Λ .

We thus obtain $L: \mathcal{J}_\Lambda \rightarrow \mathcal{E}nd_{\mathbf{C}}(\mathcal{L})$. Then this does not depend on the choice of local coordinate system and moreover it extends to the ring homomorphism $L: \mathcal{E}_\Lambda \rightarrow \mathcal{E}nd_{\mathbf{C}}(\mathcal{L})$. Since the image is contained in the differential endomorphism of \mathcal{L} , we obtain the ring homomorphism $L: \mathcal{E}_\Lambda \rightarrow \mathcal{L} \otimes_{\mathcal{O}_\Lambda} \mathcal{D}_\Lambda \otimes_{\mathcal{O}_\Lambda} \mathcal{L}^{\otimes -1}$.

PROPOSITION 10.1.1. *By $L, \mathcal{E}_\Lambda/\mathcal{E}_\Lambda(-1)$ coincides with the subsheaf of $\mathcal{L} \otimes_{\mathcal{O}_\Lambda} \mathcal{D}_\Lambda \otimes_{\mathcal{O}_\Lambda} \mathcal{L}^{\otimes -1}$ consisting of differential endomorphisms of \mathcal{L} homogeneous of degree 0.*

If we take

$$\mathcal{J}_\Lambda \in \mathfrak{g} = \mathfrak{g}_1(x, \partial) + \mathfrak{g}_0(x, \partial) + \dots$$

such that $d\mathfrak{g}_1 \equiv -\theta_X \bmod I_\Lambda \Omega^1$ and

$$\frac{1}{2} \sum \frac{\partial^2 \mathfrak{g}_1}{\partial x_i \partial \xi_i} \equiv \mathfrak{g}_0(x, \xi) \bmod \mathcal{J}_\Lambda$$

then $L(\mathfrak{g})$ gives the Euler operator of \mathcal{L} . Such a \mathfrak{g} is unique modulo $\mathcal{J}_\Lambda^2(-1) = \mathcal{E}_\Lambda(-1) \cap \mathcal{E}_X(1)$.

10.2. Let \mathcal{M} be a regular holonomic \mathcal{E}_X -module whose support is Λ . Let \mathcal{M}_0 be a coherent sub- \mathcal{E}_Λ -module of M which generates \mathcal{M} . Such an \mathcal{M}_0 is called a *saturated lattice* of \mathcal{M} . Then $\bar{\mathcal{M}} = \mathcal{M}_0/\mathcal{E}_\Lambda(-1)\mathcal{M}_0$ is an $\mathcal{E}_\Lambda/\mathcal{E}_\Lambda(-1)$ -module, which is coherent over $\mathcal{O}_\Lambda(0)$.

Since a coherent sheaf with integrable connection is locally free, we have

LEMMA 10.2.1. *$\bar{\mathcal{M}}$ is a locally free $\mathcal{O}_\Lambda(0)$ -module of finite rank.*

Since \mathfrak{g} belongs to the center of $\mathcal{E}_\Lambda/\mathcal{E}_\Lambda(-1)$, \mathfrak{g} can be considered as an endomorphism of $\mathcal{H}om_{\mathcal{E}_\Lambda/\mathcal{E}_\Lambda(-1)}(\bar{\mathcal{M}}, \mathcal{L})$, which is a locally constant sheaf on Λ . Its eigenvalues are called the *order* of \mathcal{M} with respect to \mathcal{M}_0 .

10.3. Let us take a section $G \subset \mathbf{C}$ of $\mathbf{C} \rightarrow \mathbf{C}/\mathbf{Z}$. Then there exists a unique saturated lattice \mathcal{M}_0 such that the orders of \mathcal{M} with respect to \mathcal{M}_0 are contained in G (See [K4]). Then

$$\mathcal{F} = \mathcal{H}om_{\mathcal{E}_\Lambda/\mathcal{E}_\Lambda(-1)}(\bar{\mathcal{M}}, \mathcal{L})$$

and

$$M = \exp 2\pi i \mathfrak{g} \in \mathcal{A}ut(\mathcal{F})$$

does not depend on the choice of G .

THEOREM 10.3.1 ([KK] Chapter I, § 3). *Assume that there exists an invertible \mathcal{O}_Λ -module \mathcal{L} such that $\mathcal{L}^{\otimes 2} = \omega_\Lambda \otimes \omega_X^{\otimes -1}$. Then the category of regular holonomic \mathcal{E}_X -modules with support in Λ is equivalent to the category of (\mathcal{F}, M) 's where \mathcal{F} is a locally constant \mathbf{C}_Λ -module and $M \in \mathcal{A}ut_{\mathbf{C}}(\mathcal{F})$.*

10.4. If $u \in \mathcal{M}$, then the solution to $L(P)\varphi = 0$ for $P \in \mathcal{E}_\Lambda$ with $Pu = 0$ is called a principal symbol of u and denoted by $\sigma(u)$. The homogeneous degree of $\sigma(u)$ is called the order of u . In the terminology of § 10.2, the principal symbol is a section of $\mathcal{H}om_{\mathcal{E}_\Lambda/\mathcal{E}_\Lambda(-1)}(\mathcal{E}_\Lambda u/\mathcal{E}_\Lambda(-1)u, \mathcal{L})$ and the order is the eigenvalue of \mathfrak{g} in $\mathcal{H}om_{\mathcal{E}_\Lambda/\mathcal{E}_\Lambda(-1)}(\mathcal{E}_\Lambda u/\mathcal{E}_\Lambda(-1)u, \mathcal{L})$.

10.4. When the characteristic variety is not smooth, we don't know much about the structure of holonomic systems. In this direction, we have

THEOREM 10.4.1 ([K-K] Theorem 1.2.2). *Let Z be a closed analytic subset of an open subset Ω of T^*X , $n = \dim X$, and let \mathcal{M} and \mathcal{N} be holonomic $\mathcal{E}_X|_\Omega$ -modules.*

(i) *If $\dim Z \leq n-1$, then*

$$\Gamma(\Omega; \mathcal{H}om_{\mathcal{E}_X}(\mathcal{M}, \mathcal{N})) \rightarrow \Gamma(\Omega \setminus Z, \mathcal{H}om_{\mathcal{E}_X}(\mathcal{M}, \mathcal{N}))$$

is injective.

(ii) *If $\dim Z \leq n-2$, then*

$$\Gamma(\Omega; \mathcal{H}om_{\mathcal{E}_X}(\mathcal{M}, \mathcal{N})) \rightarrow \Gamma(\Omega \setminus Z; \mathcal{H}om_{\mathcal{E}_X}(\mathcal{M}, \mathcal{N}))$$

is an isomorphism.

In particular if $\text{Supp } \mathcal{M} \subset \Lambda_1 \cup \Lambda_2$ and if $\dim(\Lambda_1 \cap \Lambda_2) \leq n-2$, then \mathcal{M} is a direct sum of two holonomic \mathcal{E}_X -modules supported on Λ_1 and Λ_2 , respectively.

Here is another type of theorem.

THEOREM 10.4.3 ([SKKO]). Let $\mathcal{M} = \mathcal{E}u = \mathcal{E}/\mathcal{J}$ be a holonomic \mathcal{E} -module defined on a neighborhood of $p \in T^*X$. Assume $\text{Supp } \mathcal{M} = \Lambda_1 \cup \Lambda_2$ and

- (i) Λ_1, Λ_2 and $\Lambda_1 \cap \Lambda_2$ are non-singular and $\dim \Lambda_1 = \dim \Lambda_2 = n$, $\dim(\Lambda_1 \cap \Lambda_2) = n-1$.
- (ii) $T_{p'} \Lambda_1 \cap T_{p'} \Lambda_2 = T_{p'}(\Lambda_1 \cap \Lambda_2)$ for any p' in a neighborhood of p in $\Lambda_1 \cap \Lambda_2$.
- (iii) The symbol ideal of \mathcal{J} coincides with the ideal of functions vanishing on $\Lambda_1 \cup \Lambda_2$.

Setting $k = \text{ord}_{\Lambda_1} u - \text{ord}_{\Lambda_2} u - 1/2$, we have

- (a) \mathcal{M} has a non-zero quotient supported on $\Lambda_1 \Leftrightarrow \mathcal{M}$ has a non-zero submodule supported on $\Lambda_2 \Leftrightarrow k \in \mathbf{Z}$.
- (b) \mathcal{M}_p is a simple \mathcal{E}_p -module $\Leftrightarrow k \notin \mathbf{Z}$.

Sketch of the proof. By a quantized contact transformation, we can transform p, Λ_1, Λ_2 and \mathcal{J} as follows:

$$\begin{aligned} p &= (0, dx_1) \\ \Lambda_1 &= \{(x, \xi); x_1 = \xi_2 = \dots = \xi_n = 0\} \\ \Lambda_2 &= \{(x, \xi); x_1 = x_2 = \xi_3 = \dots = \xi_n = 0\} \\ \mathcal{J} &= \mathcal{E}(x_1 \partial_1 - \lambda) + \mathcal{E}(x_2 \partial_2 - \mu) + \sum_{j>2} \mathcal{E} \partial_j \end{aligned}$$

In this case, we can easily check the theorem.

§ 11. APPLICATION TO THE b -FUNCTION (see [SKKO])

11.1. As one of the most successful application of microlocal analysis, we shall sketch here how to calculate the b -function of a function under certain conditions.