Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 32 (1986)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: TREES, TAIL WAGGING AND GROUP PRESENTATIONS
Autor: Armstrong, M. A.
Kapitel: 1. Graphs
DOI: https://doi.org/10.5169/seals-55090

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 03.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

L'Enseignement Mathématique, t. 32 (1986), p. 261-270

TREES, TAIL WAGGING AND GROUP PRESENTATIONS

by M. A. ARMSTRONG

The Bass-Serre theorem gives a presentation for a group of automorphisms of a tree. Like all good theorems it has attracted considerable attention and there are now several proofs available [4], [3], [1]. Our goal is a natural elementary proof which makes maximal use of the geometry of the tree.

1. GRAPHS

A *graph* X consists of two sets E (directed edges) and V (vertices) and two functions

$$E \to E, \qquad e \mapsto \bar{e}$$

$$E \to V \times V, \qquad e \mapsto (i(e), t(e))$$

which satisfy $\bar{\bar{e}} = e$, $\bar{e} \neq e$ and $i(\bar{e}) = t(e)$ for each $e \in E$. The vertices $i(e)$, $t(e)$ are the initial and terminal vertices of the directed edge e, and $\bar{e}$ is the reverse of e. Henceforth we refer to directed edges simply as edges.

A path in X joining vertex u to vertex v is an ordered string of edges $e_1 e_2 \dots e_n$ such that $i(e_1) = u$, $i(e_{k+1}) = t(e_k)$ for $1 \leqslant k \leqslant n-1$, and $t(e_n) = v$. If $v = u$ we have a circuit. A path of the form $e\bar{e}$ is a *round trip* and a circuit which does not contain any round trips will be called a loop. If any two distinct vertices may be joined by a path then the graph is connected. A *tree* is a connected graph which does not contain any loops.

Let X be a tree. A path in X is a *geodesic* if it does not contain any round trips. Given distinct vertices u, v of X there is a *unique* geodesic $\overrightarrow{uv}$ which joins u to v.

An action of a group G on a graph X is an action of G on E and on V such that $g\bar{e} = \overline{ge}$, $i(ge) = gi(e)$, $t(ge) = gt(e)$ and $ge \neq \bar{e}$ for each $e \in E$. Because group elements are not allowed to reverse edges we have a

quotient graph X/G. When G acts on X we shall often say that G is a *group of automorphisms of* X.

We adopt the usual notation whereby G_x denotes the stabilizer of a vertex x. If $g \in G$ happens to fix x we write g_x for the element g thought of as a member of G_x. Of course G_e denotes the stabilizer of the edge e. If x is a vertex of e then G_e is a subgroup of G_x.

Suppose G acts on a tree X. If $g \in G$ fixes the vertices u, v then it must fix the whole geodesic $\overrightarrow{uv}$, since otherwise the image of $\overrightarrow{uv}$ under g would be a second geodesic from u to v.

2. Lifting edges

Let G be a group of automorphisms of a tree X. Choose a maximal tree M in X/G and lift it [4, Proposition I.14] to a subtree T of X. The vertices of T form a set of representatives for the action of G on the vertices of X. For each pair of edges f, $\bar{f}$ from $X/G - M$ select one, say f, and lift it to an edge e of X which has its initial vertex x in T. Exactly one vertex z of T lies in the same orbit as $t(e)$ and we choose an element γ_f from G that maps z onto $t(e)$. We can now lift $\bar{f}$ to $(\gamma_f)^{-1}\bar{e}$. This has its initial vertex z in T and $\gamma_{\bar{f}} = (\gamma_f)^{-1}$ sends the vertex x of T to its terminal vertex (Figure 1). Finally we extend the correspondence $f \to \gamma_f$ over the edges of M by setting $\gamma_f = 1$ (the identity element of G) whenever $f \in M$.

The *Bass-Serre theorem* [4, Theorem I.13] gives the following presentation for G.

(a) *Generators.* The elements of all the G_w where w is a vertex of T and the γ_f where f is an edge of X/G.

(b) *Relations.* The internal relations of each stabilizer G_w together with

$\gamma_f = 1$ if f is an edge of M,

$\gamma_{\bar{f}} = (\gamma_f)^{-1}$ and

$\gamma_{\bar{f}}\, g_x \gamma_f = (\gamma_{\bar{f}}\, g\, \gamma_f)_z$ where e is the chosen lift of f and $g \in G_e$.

(If f is an edge of M then $z = t(e)$ and the final relation reduces to $g_x = g_z$ whenever $g \in G_e$).