Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 32 (1986)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: TREES, TAIL WAGGING AND GROUP PRESENTATIONS
Autor: Armstrong, M. A.
Kapitel: 5. Nearest fixed points
DOI: https://doi.org/10.5169/seals-55090

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 03.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

$$
\begin{aligned}
& b_{x_1} \lambda_{f_1} b_{y_2} \lambda_{f_2'} \dots b_{y_s} \lambda_{f_s'} b_v R \\
= {} & a_{x_1} \lambda_{f_1} \lambda_{\bar{f}_1} a_{x_1}^{-1} b_{x_1} \lambda_{f_1} b_{y_2} \lambda_{f_2'} \dots b_{y_s} \lambda_{f_s'} b_v R \\
= {} & a_{x_1} \lambda_{f_1} \lambda_{\bar{f}_1} c_{x_1} \lambda_{f_1} b_{y_2} \lambda_{f_2'} \dots b_{y_s} \lambda_{f_s'} b_v R \\
= {} & a_{x_1} \lambda_{f_1} (\gamma_{\bar{f}_1} c \gamma_{f_1})_{z_1} b_{y_2} \lambda_{f_2'} \dots b_{y_s} \lambda_{f_s'} b_v R \\
= {} & a_{x_1} \lambda_{f_1} (\gamma_{\bar{f}_1} c \gamma_{f_1})_{x_2} b_{y_2} \lambda_{f_2'} \dots b_{y_s} \lambda_{f_s'} b_v R \\
= {} & a_{x_1} \lambda_{f_1} a'_{x_2} b_{y_2} \lambda_{f_2'} \dots b_{y_s} \lambda_{f_s'} b_v R
\end{aligned}
$$

where $a'_{x_2} = (\gamma_{\bar{f}_1} c \gamma_{f_1})_{x_2}$. If x_2 happens to equal y_2 then we simplify this further to

$$a_{x_1} \lambda_{f_1} a''_{x_2} \lambda_{f_2} b_{y_3} \lambda_{f_3'} \dots b_{y_s} \lambda_{f_s'} b_v R$$

where a''_{x_2} is the product $a'_{x_2} b_{y_2}$ in G_{x_2}. We now compare a_{x_2} with a'_{x_2} if $x_2 \neq y_2$, noting that $\gamma_{f_2} = 1$ in this case, or with a''_{x_2} if $x_2 = y_2$, and repeat the process. Eventually we obtain

$$b_{x_1} \lambda_{f_1} b_{y_2} \lambda_{f_2'} \dots b_{y_s} \lambda_{f_s'} b_v R = a_{x_1} \lambda_{f_1} a_{x_2} \lambda_{f_2} \dots a_{x_r} \lambda_{f_r} a''_v R .$$

As $g = a_{x_1} \gamma_{f_1} \dots a_{x_r} \gamma_{f_r} a_v = a_{x_1} \gamma_{f_1} \dots a_{x_r} \gamma_{f_r} a''_v$ we see that $a''_v = a_v$. This completes the proof that ψ is well defined.

5. Nearest fixed points

To show ψ is a homomorphism we shall verify

$$\psi(hg) = \psi(h)\psi(g)$$

under the assumption that h either leaves some vertex of T fixed or is one of the elements γ_f. This is sufficient because the elements of the G_w (w a vertex of T) together with the γ_f (f an edge of $X/G-M$) form a set of generators for G.

Suppose h fixes the vertex w of T. Walk along the geodesic $\overrightarrow{vw}$ and let x be the first vertex we meet which is left fixed by h. Then $\overrightarrow{vx}$ is contained in T, and $\overrightarrow{vx}$ followed by $h(\overrightarrow{xv})$ is the geodesic from v to hv. This quits T for the first time at x and we see that

$$\psi(h) = h_x R .$$



FIGURE 3

Using the geodesic from v to gv we have $\psi(g) = a_{x_1}\lambda_{f_1} \dots a_{x_r}\lambda_{f_r} a_v R$ in the usual way. Therefore

$$\psi(h)\psi(g) = h_x a_{x_1}\lambda_{f_1} \dots a_{x_r}\lambda_{f_r} a_v R .$$

In order to compute $\psi(hg)$ we need the geodesic from v to $(hg)v$. We can construct this as follows, take $\overrightarrow{v\,hv}$ followed by the image of $\overrightarrow{v\,gv}$ under h and remove any round trips.

If $\overrightarrow{v\,gv}$ does not contain all of $\overrightarrow{vx}$ (Figure 3) then $\overrightarrow{v(hg)v}$ leaves T for the first time at x. A tail wag of $\overrightarrow{v(hg)v}$ using h_x^{-1} leads us to a path which has the same tail as $\overrightarrow{v\,gv}$, then the process continues as for g. Thus

$$\psi(hg) = h_x a_{x_1}\lambda_{f_1} \dots a_{x_r}\lambda_{f_r} a_v R = \psi(h)\psi(g) .$$

Otherwise $\overrightarrow{v\,gv}$ contains all of $\overrightarrow{vx}$ (Figure 4) and we split the argument into three cases.



FIGURE 4

(a) $\overrightarrow{v\,gv}$ stays in T for at least one more edge after x. Then $\overrightarrow{v(hg)v}$ must leave T at x. As above, a first choice of h_x^{-1} leads to a path with the same tail as $\overrightarrow{v\,gv}$.

(b) $\overrightarrow{v\,gv}$ and $\overrightarrow{v(hg)v}$ both leave T at x. Then $x_1 = x$ and we write a_x instead of a_{x_1}. A first tail wag of $\overrightarrow{v(hg)v}$ using $\gamma_{\bar{f}_1}(h_x\, a_x)^{-1}$ produces the same path as a first tail wag of $\overrightarrow{v\,gv}$ using $\gamma_{\bar{f}_1}\, a_x^{-1}$. Thus

$$\psi(hg) = h_x\, a_x\, \lambda_{f_1}\, a_{x_2}\, \lambda_{f_2} \dots a_{x_r}\, \lambda_{f_r}\, a_v R = \psi(h)\psi(g)\,.$$

(c) $\overrightarrow{v\,gv}$ leaves T at x, but $\overrightarrow{v(hg)v}$ stays in T for at least one more edge after x. Then $x_1 = x$, $\gamma_{f_1} = 1$ and we may as well equate a_{x_1} with h_x^{-1}. A first tail wag of $\overrightarrow{v\,gv}$ using h_x gives a path with the same tail as $\overrightarrow{v(hg)v}$. Thus

$$\begin{aligned}\psi(hg) &= a_{x_2}\, \lambda_{f_2} \dots a_{x_r}\, \lambda_{f_r}\, a_v R \\ &= h_x\, h_x^{-1}\, a_{x_2}\, \lambda_{f_2} \dots a_{x_r}\, \lambda_{f_r}\, a_v R \\ &= \psi(h)\psi(g)\,.\end{aligned}$$

Suppose finally that $h = \gamma_f$ for some edge f of $X/G-M$. As usual e is the chosen lift of f into X with $x = i(e) \in T$ and $z = t(\gamma_{\bar{f}}\, e)$. Let $y = i(\gamma_{\bar{f}}\, e)$. The geodesic from v to $\gamma_f v$ is made up of $\overrightarrow{vx}$ followed by e followed by $\gamma_f(\overrightarrow{zv})$. This leaves T for the first time at x and a single tail wag using $\gamma_{\bar{f}}$ produces $\overrightarrow{zv}$. Therefore

$$\psi(\gamma_f) = \lambda_f R\,.$$

To obtain the geodesic from v to $(\gamma_f g)v$ we follow $\overrightarrow{v\gamma_f v}$ by $\gamma_f(\overrightarrow{v\,gv})$ and then remove any round trips (Figure 5). If $\overrightarrow{v\,gv}$ does not contain $\overrightarrow{vy}$, then $\overrightarrow{v(\gamma_f g)v}$ leaves T for the first time at x and a single tail wag using $\gamma_{\bar{f}}$



FIGURE 5

produces a path with the same tail as $\overrightarrow{v\,gv}$. The process then continues as for g and

$$\psi(\gamma_f g) = \lambda_f\, a_{x_1}\, \lambda_{f_1} \dots a_{x_r}\, \lambda_{f_r}\, a_v R = \psi(\gamma_f)\psi(g)\,.$$

Otherwise $\overrightarrow{v\,gv}$ contains $\overrightarrow{vy}$. Then $x_1 = z$, $\gamma_{f_1} = \gamma_{\bar{f}}$ and we may as well take $a_{x_1} = 1$. A first tail wag of $\overrightarrow{v\,gv}$ using γ_f leaves a path with the same tail as $\overrightarrow{v(\gamma_f g)v}$. Thus

$$\begin{aligned}\psi(\gamma_f g) &= a_{x_2}\, \lambda_{f_2} \dots a_{x_r}\, \lambda_{f_r}\, a_v R \\ &= \lambda_f\, \lambda_{\bar{f}}\, a_{x_2}\, \lambda_{f_2} \dots a_{x_r}\, \lambda_{f_r}\, a_v R \\ &= \psi(\gamma_f)\psi(g)\,.\end{aligned}$$

This completes the proof that ψ is a homomorphism.

Our construction of ψ ensures that if $\psi(g) = R$ then $g = 1$. So ψ is injective. The cosets $h_w R$ (w a vertex of T and $h(w) = w$) and $\lambda_f R$ (f an edge of X/G) together generate $[(*\,G_w)*F]/R$. Now $\psi(h) = h_x R$ where x is the nearest fixed point of h to v. But h fixes all of $\overrightarrow{xw}$ so

$$\psi(h) = h_x R = h_w R\,.$$

Also

$$\psi(\gamma_f) = \lambda_f R\,.$$

Therefore the image of ψ is all of $[(*\,G_w)*F]/R$ and we have shown that ψ is an isomorphism.

The author would like to thank the members of the Mathematics Department of the University of Geneva for their hospitality during the preparation of this article.

REFERENCES

[1] DICKS, W. *Groups Trees and Projective Modules.* Lect. Notes in Math. 790, Springer-Verlag 1980.

[2] HAUSMANN, J.-C. Sur l'usage de critères pour reconnaître un groupe libre, un produit amalgamé ou une *HNN*-extension. *L'Enseignement Mathématique* 27 (1981), 221-242.