

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 32 (1986)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ON THE JONES POLYNOMIAL Swiss Seminar in Berne  
**Autor:** de la Harpe, Pierre / Kervaire, Michel / Weber, Claude  
**Kapitel:** §8. L. Kauffman's approach to V. Jones' one-variable polynomial  
**DOI:** <https://doi.org/10.5169/seals-55091>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 19.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

of  $K$ . Hence, for a knot, a link with a single component, the exponent of  $m$  in  $P_K(l, m)$  is even and therefore  $\Delta_K(t) = P_K(i, i(t^{1/2} - t^{-1/2}))$  is indeed a Laurent polynomial in  $t$ .

To obtain the one-variable Jones polynomial we use the substitution  $l = it, m = i(t^{1/2} - t^{-1/2})$ . Explicitly,

$$V_K(t) = P_K(it, i(t^{1/2} - t^{-1/2}))$$

Then we have

PROPERTY 7.6.  $V_K(t)$  satisfies the skein invariance

$$tV(K_+) - t^{-1}V(K_-) + (t^{1/2} - t^{-1/2})V(K_0) = 0,$$

which (together with  $V(\bigcirc) = 1$ ) characterizes Jones one-variable polynomial, with the sign conventions used in reference [Jo<sub>3</sub>].

Whereas  $P_K(l, m)$  determines  $\Delta_K(t)$  and  $V_K(t)$ , it is known that there are no other relations between these polynomials. More precisely:

(1) The Alexander polynomial  $\Delta_K(t)$  does not determine Jones polynomial  $V_K(t)$  because the trivial knot  $\bigcirc$  and Conway's eleven crossing knot  $11_{471}$  have  $\Delta(t) = 1$ , but  $V_K(t) \neq 1$  for  $K = 11_{471}$ .

(2)  $V_K(t)$  does not determine  $\Delta_K(t)$ : The knots  $4_1$  and  $11_{388}$  have the same  $V(t)$  but different  $\Delta(t)$ .

(3)  $V_K(t)$  and  $\Delta_K(t)$  together do not determine  $P_K(l, m)$ : The knot  $11_{388}$  and its mirror image have the same  $V(t)$  and  $\Delta(t)$  but different  $P(l, m)$ .


For more details on these questions, see [L.-M.].

We now turn to L. Kauffman's definition of the one-variable Jones polynomial  $V_K(t)$  directly from the link diagram.

## § 8. L. KAUFFMAN'S APPROACH TO V. JONES' ONE-VARIABLE POLYNOMIAL


The importance of Kauffman's approach [Ka<sub>3</sub>] is that it gives a new way to define and compute Jones polynomial  $V_K(t)$ . It is by using this definition that Kauffman and Murasugi prove their theorems about alternating links (see § 10 and 11).

Let  $L$  be an *unoriented* link diagram. Look at a double point; with no string orientation, they all look the same, up to a local homeomorphism:




Locally, the plane  $\mathbf{R}^2$  is divided into four regions.


Look at the quarter turn the “over” line must make, in the positive sense, in order to coincide with the “under” line. Call “*A*” the two regions which are swept by the over line during the trip. Call “*B*” the other two.



*Definition.* A marker for a double point is a choice of “*A*” or “*B*” for this double point. It is symbolised like that:




Marker *A*




Marker *B*

Now, if a marker is chosen, one can split the link diagram by connecting the two opposite regions whose name has been elected. Here are the pictures.



splitting if marker  
A is chosen



splitting if marker  
B is chosen

*Definition.* A state  $S$  for  $L$  is a choice of a marker at every double point of  $L$ .

Suppose now that a state  $S$  for  $L$  is given. Make the correct splitting at every double point of  $L$ . The underlying knot projection  $\Gamma$  is transformed into a bunch  $\Gamma_S$  of disjoint simple closed curves in  $S^2$ . Let  $|S|$  be the number of curves in  $\Gamma_S$ .

Write  $a(S)$  for the number of markers  $A$  in the state  $S$  and write  $b(S)$  for the number of  $B$ 's.

If  $c(L)$  denotes the number of crossings (double points) of  $L$ , one clearly has  $2^{c(L)}$  states.

$L$  being given, Kauffman defines a polynomial  $\langle L \rangle \in \mathbf{Z}[A, B, d]$  in the following way:

$$\langle L \rangle = \sum_S A^{a(S)} B^{b(S)} d^{|S|-1}$$

the summation being taken over the  $2^{c(L)}$  states.

*Notations.* Write " $\bigcirc$ " for an unoriented, connected, simple closed curve in  $\mathbf{R}^2$  and write  $\bigcirc \amalg L$  for a disjoint union of such a diagram and an unoriented link diagram  $L$ .

*Property 1.*  $\langle \bigcirc \rangle = 1$ .

*Property 2.*  $\langle \bigcirc \amalg L \rangle = d \langle L \rangle$  if  $L$  is non empty.

*Property 3.* Let  $L$  be an unoriented link diagram. Select a crossing  $\times$  and write  $L_A$  for the diagram obtained from  $L$  by connecting the two regions  $A$  at  $\times$ , and write  $L_B$  for the diagram obtained by connecting the two  $B$ 's. Then:

$$\langle L \rangle = A \langle L_A \rangle + B \langle L_B \rangle .$$

PROPOSITION 8.1.  $\langle \rangle$  is the unique function from the set of unoriented link diagrams to  $\mathbb{Z}[A, B, d]$  which satisfies properties 1, 2 and 3.

The proof is straightforward.

PROPOSITION 8.2. If one sets  $B = A^{-1}$  and  $d = -(A^2 + A^{-2})$ , one gets a function into  $\mathbb{Z}[A^{\pm 1}]$  which is invariant under Reidemeister moves (ii) and (iii).

*Notations.* Following Kauffman, we shall still write  $\langle \rangle$  for the function into  $\mathbb{Z}[A^{\pm 1}]$ . From now on, only this function will be used.

We now recall briefly Kauffman's proof of proposition 8.2.

First of all, we shall use Kauffman's schematic way of writing property 3:

$$\langle \times \rangle = A \langle \rangle \langle \rangle + B \langle \asymp \rangle$$

Invariance under move (ii):

$$\begin{aligned} \langle \overbrace{\times} \rangle &= A \langle \overline{\times} \rangle + B \langle \times \rangle \\ &= A [B \langle \overline{\asymp} \rangle + A \langle \asymp \rangle] \\ &\quad + B [B \langle \asymp \rangle + A \langle \rangle \langle \rangle] \\ &= (ABd + A^2 + B^2) \langle \asymp \rangle + AB \langle \rangle \langle \rangle \\ &= \langle \rangle \langle \rangle, \end{aligned}$$

since we have set  $B = A^{-1}$  and  $d = -(A^2 + A^{-2})$ .

Invariance under move (iii):

$$\begin{aligned} \langle \overline{\times} \rangle &= B \langle \overline{\times} \rangle + A \langle \overline{\times} \rangle \\ &= B \langle \overline{\asymp} \rangle + A \langle \overline{\asymp} \rangle \end{aligned}$$

by invariance under move (ii)

$$= \langle \text{diagram} \rangle .$$


Q.E.D.

This seems to be as far as one can get without orienting link diagrams, because  $\langle \rangle$  is *not* invariant under Reidemeister move (i).

To remedy this state of affairs, Kauffman proceeds like this:

Let  $L$  be an *oriented* link diagram.

Recall now that, up to a rotation in  $\mathbf{R}^2$ , there are two types of double points:



Sign +1

Sign -1

*Definition.* The writhe number  $w(L)$  is the sum of the signs of the double points of  $L$ .

This number is also called twist number. It was known to Tait and much used by Little. See § 9 of these notes.

Kauffman's polynomial  $f_L(A) \in \mathbf{Z}[A^{\pm 1}]$  is then defined in the following way:

$$f_L(A) = (-A)^{-3w(L)} \langle L \rangle .$$

**PROPOSITION 8.3.** *The polynomial  $f$  is invariant under Reidemeister moves (i), (ii) and (iii).*

*Proof of proposition 8.3.* The writhe number is unchanged by the moves (ii) and (iii). Hence proposition (8.2) implies the invariance of  $f$  under the moves (ii) and (iii).

We now prove the invariance under move (i).

Let  $\hat{L}$  be a link diagram with a portion looking like this:



and let  $L$  be the link diagram obtained from  $\hat{L}$  by removing the loop. It is immediate that

$$\hat{L}_A = \left. \vphantom{\hat{L}_A} \right\} \quad \text{and} \quad \hat{L}_B = \left. \vphantom{\hat{L}_B} \right\} \circ$$

If we apply property 3 for  $\langle \rangle$  we get

$$\langle \hat{L} \rangle = A \langle L \rangle + A^{-1} \langle L \amalg \bigcirc \rangle .$$

By property 2

$$\langle \hat{L} \rangle = A \langle L \rangle + A^{-1}(-A^2 - A^{-2}) \langle L \rangle .$$

So  $\langle \hat{L} \rangle = (A - A^{-1}(A^2 + A^{-2})) \langle L \rangle = (-A)^{-3} \langle L \rangle .$

Now, for any orientation of the string, the sign of the double point is  $-1$ .

Hence  $w(\hat{L}) = w(L) - 1$ .

Going back to the definition,

$$\begin{aligned} f_{\hat{L}} &= (-A)^{-3w(\hat{L})} \langle \hat{L} \rangle = (-A)^{-3w(L)+3} \langle \hat{L} \rangle \\ &= (-A)^{-3w(L)+3} (-A)^{-3} \langle L \rangle \\ &= (-A)^{-3w(L)} \langle L \rangle = f_L . \end{aligned}$$

The proof for the other loop is similar.

Q.E.D.

From proposition 8.3 we deduce that Kauffman's polynomial induces a map  $f: \mathcal{L} \rightarrow \mathbf{Z}[A^{\pm 1}]$ .

**THEOREM 8.4.** *The map  $f: \mathcal{L} \rightarrow \mathbf{Z}[A^{\pm 1}]$  satisfies:*

1.  $f(\bigcirc) \equiv 1$ .
2. If  $L_+, L_-$  and  $L_0$  are skein related (see § 3), then:

$$A^4 f_{L_+} - A^{-4} f_{L_-} = (A^{-2} - A^2) f_{L_0} .$$

From the universality of Jones polynomial, we obtain:

COROLLARY 8.5. *Let  $K$  be an oriented link in  $\mathbf{R}^3$  and let  $L$  be an oriented diagram of  $K$ . Then:*

$$V_K(t) = f_L(t^{1/4}).$$

Recall that we use Jones definition in the Bulletin AMS [Jo<sub>3</sub>] for  $V_K$ . If we were to use Jones definition in the Notices AMS [Jo<sub>4</sub>], we would set  $A = t^{-1/4}$ .

*Proof of theorem 8.4.* The proof of 1. is straightforward from the definition. For 2., using Kauffman's notations one has:

$$\langle \times \rangle = A \langle \smile \rangle + A^{-1} \langle \rangle \langle \rangle$$

and

$$\langle \smile \rangle = A^{-1} \langle \times \rangle + A \langle \rangle \langle \rangle$$

Hence:

$$A \langle \times \rangle - A^{-1} \langle \smile \rangle = (A^2 - A^{-2}) \langle \smile \rangle$$

If we orient the strings and put the writhe number in the picture, we get the formula 2. Q.E.D.

Using L. Kauffman's definition of Jones polynomial, the following properties are easily proved (enjoyable exercise left to the reader):

I. If  $K_1$  and  $K_2$  are two oriented links in  $S^3$ , let  $K_1 \amalg K_2$  denote their distant union (one in each hemisphere). Then:

$$V_{K_1 \amalg K_2} = \mu V_{K_1} \cdot V_{K_2}$$

where  $\mu = -(t^{1/2} + t^{-1/2})$ .

II. Let  $K_1 \# K_2$  denote any connected sum of  $K_1$  and  $K_2$  as in § 7 prop. 4. Then:

$$V_{K_1 \# K_2} = V_{K_1} \cdot V_{K_2}.$$

III. Let  $K^\times$  denote the mirror image of  $K$ . Then:

$$V_{K^\times}(t) = V_K(t^{-1}).$$

The first three formulas are rather straightforward from the definitions.

IV. (Jones reversing result). Let  $K$  be an oriented link in  $S^3$  and let  $\gamma$  be a component of  $K$ . Let  $\lambda$  be the linking coefficient of  $\gamma$  with what is left of  $K$  when we remove  $\gamma$ . (We suppose that this is not empty!) Let  $\hat{K}$  be the oriented link obtained from  $K$  by changing the orientation of  $\gamma$ , while keeping the others fixed. Then:

$$V_{\hat{K}}(t) = t^{3\lambda}V_K(t).$$

*Proof.* Of course, we have  $\langle K \rangle = \langle \hat{K} \rangle$ , because, for the polynomial  $\langle \rangle$ , orientations do not matter.

Now:  $w(K) = w(\gamma) + 2\lambda$ .

So:  $w(\hat{K}) = w(\gamma) - 2\lambda$ .

Hence:  $w(\hat{K}) = w(K) - 4\lambda$ .

We substitute and get:

$$\begin{aligned} f_{\hat{K}}(A) &= (-A)^{-3w(\hat{K})} \langle \hat{K} \rangle = (-A)^{-3w(K) + 12\lambda} \langle K \rangle \\ &= (-A)^{12\lambda} (-A)^{-3w(K)} \langle K \rangle = (-A)^{12\lambda} f_K(A) = A^{12\lambda} f_K(A). \end{aligned}$$

As one substitutes  $t^{1/4}$  for  $A$  to get Jones 1-variable polynomial, the result follows.

To finish this paragraph, we illustrate quickly Kauffman's definition by computing Jones one variable polynomial for the right-handed trefoil  $T_+$ . (Compare § 3.)

There are 8 states associated to the standard knot diagram. One readily sees that

$$\langle T_+ \rangle = A^3d + 3A^2Bd^0 + 3AB^2d + B^3d^2.$$

Substituting  $d = -(A^2 + A^{-2})$  and  $B = A^{-1}$  one gets

$$\langle T_+ \rangle = -A^5 - A^{-3} + A^{-7}.$$

As  $w(T_+) = 3$ , one gets

$$f_{T_+}(A) = (-A)^{-9} \langle T_+ \rangle = A^{-4} + A^{-12} - A^{-16}.$$

Substituting  $t = A^{1/4}$  one finally obtains

$$V_{T_+}(t) = t^{-1} + t^{-3} - t^{-4} = t^{-4}(-1 + t + t^3).$$

Now, if one uses our computation in § 3

$$P(T_+) = -2a_- a_+^{-1} - a_-^2 a_+^{-2} + a_+^{-2} a_0^2$$

and substitutes  $a_+ = l, a_- = l^{-1}, a_0 = m$  one gets

$$P_{T_+}(l, m) = (-2l^{-2} - l^{-4})m^0 + l^{-2}m^2.$$

The last substitution  $l = it; m = i(t^{1/2} - t^{-1/2})$  gives (with relief!) the same result for Jones one variable polynomial. (Bulletin AMS definition.)

### § 9. TAIT CONJECTURES

Tait was primarily interested in the classification of knots (i.e. one component links). He organized the job in two steps.

*Step 1.* Classify generic immersions of the circle in  $S^2$  (not  $\mathbf{R}^2$ !) modulo homeomorphisms (possibly orientation reversing) of  $S^2$ . This was mostly done by the Rev. T. P. Kirkman (around 1880).

In this process, one has to remember that one is looking at knots in  $\mathbf{R}^3$  and that one is trying to list knots according to their “knottiness”, i.e. their minimal crossing number. So, Tait first reduced the number of double points of a generic immersion by making one “local 180° rotation”.

*Examples.*

