

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 32 (1986)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SOME CHARACTERIZATIONS OF COXETER GROUPS
Autor: Deodhar, Vinay V.
Kapitel: §1. Introduction
DOI: <https://doi.org/10.5169/seals-55081>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

SOME CHARACTERIZATIONS OF COXETER GROUPS

by Vinay V. DEODHAR ¹⁾

ABSTRACT: The aim of this note is to compile together various characterizations of Coxeter groups. Some of these are well-known, some are not-so-well-known and some are entirely new. The motivation behind the new ones is explained in the introduction.

§ 1. INTRODUCTION

Coxeter groups originally arose as group of symmetries of various geometrical objects. These became the center of activity in Lie Theory because of fundamental work of E. Cartan, H. Weyl and others regarding the structure of semi-simple Lie algebras. A little later on, Coxeter gave a complete classification of finite groups generated by reflections which included the Weyl groups, the dihedral groups and two sporadic groups (H_3 and H_4). In doing so, he gave a presentation of these groups which then led to other families of groups which have similar presentations. These include the affine Weyl groups and this is motivation enough to develop the theory of general Coxeter groups. Such a study was initiated around late 60's and an important characterization in terms of the so-called exchange condition was given (cf. [B], [S]). In recent years, this theory has been further developed and a lot of important work has been done. A major part of this work is in connection with the Bruhat ordering and its role in various different contexts in Lie theory ([K-L]). Another object under investigation is the so-called root-system of a general Coxeter group. It is seen that a number of properties of Coxeter groups can be derived using these root-systems ([D]). In an attempt to understand the role of the above two concepts in this theory, the author found out that these two

¹⁾ Partially supported by N.S.F. grant No. 82-00752.

properties characterize Coxeter groups. It therefore seems worthwhile to compile together various characterizations of Coxeter groups. This is done in § 2. A part of it is of expository nature though our proofs for the well-known characterizations are somewhat more direct.

The author wishes to thank J. Tits and J. E. Humphreys for encouragement for writing this note up. Thanks are also due to T. Springer for his comments on a preliminary version of this note.

§ 2. MAIN THEOREM

Let W be a group generated by a set S of involutory generators (i.e. order $s = 2 \forall s \in S$). One then has the notion of the length $l(w)$ of an element $w \in W$ viz. the least integer k such that $w = s_1 \dots s_k$ with $s_i \in S$. Further, such an expression is called a reduced expression. We then have the following:

MAIN THEOREM. Let W, S be as above. Then the following conditions are equivalent:

1) *Coxeter condition*: If \tilde{W} is the free group generated by a copy \tilde{S} of S subject to relations $(\tilde{s})^2 = \text{id} \forall s \in S$ and $\eta: \tilde{W} \rightarrow W$ is the canonical map, then $\text{Ker } \eta$ is generated as a normal subgroup by elements of the type:

$$\{(\tilde{s}_1 \tilde{s}_2)^{m_{s_1, s_2}}, s_1 \neq s_2 \in S, m_{s_1, s_2} \geq 2\} \text{ i.e. } \langle S \mid s^2 = \text{id} \forall s \in S, (s_1 s_2)^{m_{s_1, s_2}} = \text{id}$$

for some pairs $s_1 \neq s_2$ in $S \rangle$ is a presentation of W . (Note that the above relations *may not* involve *all* pairs $s_1 \neq s_2$).

2) *Root-system condition*: There exists a representation \dot{V} of W over \mathbf{R} , a W -invariant set Φ of non-zero vectors in V which is symmetric (i.e. $\Phi = -\Phi$) and a subset $\{e_s \mid s \in S\}$ of Φ such that the following conditions are satisfied.

(i) Every $\phi \in \Phi$ can be written as $\sum_{s \in S} a_s e_s$ with either all $a_s \geq 0$ or all

$a_s \leq 0$, but not in both ways.

(Accordingly, we write $\phi > 0$ or $\phi < 0$.)

(ii) $e_s \in \Phi$, $s(e_s) < 0$ and $s(\phi) > 0$ for all $\phi > 0$, $\phi \neq e_s$.

(iii) If $w \in W$, $s, s' \in S$ are such that $w(e_{s'}) = e_s$. Then $ws'w^{-1} = s$.