Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	32 (1986)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	THE GEOMETRY OF THE HOPF FIBRATIONS
Autor:	Gluck, Herman / Ziller, Wolfgang
Kapitel:	5. Normed Division Algebras and the Cayley Numbers
DOI:	https://doi.org/10.5169/seals-55085

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

## Download PDF: 18.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

isometry of the base space  $HP^{n-1}$  in its canonical metric. It is easy to check that when n = 2, every orientation preserving isometry of the base  $HP^1 = S^4(1/2)$  can be produced this way, while no orientation reversing one can (since the group is connected). We remark without proof that all isometries of  $HP^{n-1}$ , n > 2, can be produced this way, and that they are all orientation preserving.

# 5. NORMED DIVISION ALGEBRAS AND THE CAYLEY NUMBERS

In order to describe the Hopf fibration  $H: S^7 \hookrightarrow S^{15} \to S^8$  in the next section, we first review here some facts about normed division algebras and the arithmetic of Cayley numbers. More can be found in two excellent references, [Cu] and [H-L, pp. 140-145].

A normed division algebra B is a finite dimensional algebra over the reals R, with multiplicative unit 1, and equipped with an inner product <, > whose associated norm | | satisfies

$$|xy| = |x| |y|$$
 for all  $x, y \in B$ .

By Hurwitz' Theorem ([Hu 1], 1898), a proof of which we will outline here, every normed division algebra is isomorphic to either the reals R, the complex numbers C, the quaternions H or the Cayley numbers Ca. Actually, what Hurwitz proved is that normed division algebras can only occur in dimensions 1, 2, 4 and 8. He stated the corresponding uniqueness result without proof. In [Hu 2], published in 1923 after his death, Hurwitz credits E. Robert [Ro] with writing out the details of the uniqueness argument in a 1912 Zurich thesis.

Now let B denote a given normed division algebra. Let Re B denote the one-dimensional linear subspace spanned by the identity 1, and Im B the orthogonal complement of Re B. Then each  $x \in B$  has a unique orthogonal decomposition,

 $x = x_1 + x'$ ,  $x_1 \in \operatorname{Re} B$  and  $x' \in \operatorname{Im} B$ ,

into its real and imaginary parts. Conjugation in B is defined by:

$$\bar{x} = x_1 - x' \, .$$

Here are some basic facts about arithmetic in any normed division algebra B:

- 1)  $\langle xw, yw \rangle = \langle x, y \rangle |w|^2 = \langle wx, wy \rangle$ . Thus right or left multiplication by a unit vector w is an isometry of B.
- 2) Every nonzero  $x \in B$  has a unique left and right inverse:

$$x^{-1} = \bar{x}/|x|^2$$
.

3) Given x and y in B with  $x \neq 0$ , the equations

xw = y and wx = y

can each be solved uniquely, with

$$w = x^{-1}y$$
 and  $w = yx^{-1}$ 

respectively.

 $\overline{xy} = \overline{y} \, \overline{x}$ .

- 5) If x is imaginary (that is,  $x \in \text{Im } B$ ), then  $x^2 = -|x|^2$ .
- 6) Orthogonal imaginaries anti-commute. That is,

 $x, y \in \text{Im } B$  and  $\langle x, y \rangle = 0$  imply xy = -yx.

7) The Moufang identities, the first two of which say that left and right multiplication by xyx can be performed successively:

$$(xyx)z = x(y(xz))$$
$$z(xyx) = ((zx)y)x$$
$$x(yz)x = (xy)(zx).$$

Given three elements  $x, y, z \in B$ , their associator is defined by

$$[x, y, z] = (xy)z - x(yz).$$

The following weak form of associativity always holds in a normed division algebra: the trilinear form [x, y, z] is *alternating*, i.e., it vanishes whenever two of its arguments are equal. Such an algebra is said to be *alternative*.

The Cayley-Dickson process generalizes the way in which the complex numbers are built up from the reals, and begins with the following

PROPOSITION 5.1. (see [Cu] or [H-L]). Let A be a subalgebra (containing 1) of the normed division algebra B. Let  $\varepsilon$  be an element of B orthogonal to A with  $|\varepsilon| = 1$ . Then

i) At is orthogonal to A, and

ii)  $(a+b\varepsilon)(c+d\varepsilon) = (ac-\overline{d}b) + (da+b\overline{c})\varepsilon$  for all a, b, c, d in A.

HOPF FIBRATIONS

The proof makes use of the commutation rules given in Fact 6 above. We note for future use that the above proposition implies that any subalgebra of Ca generated by two elements must be isomorphic to R, Cor H, and hence must be associative.

Suppose now that we start with a normed division algebra A and define a product on  $A \oplus A$  by

 $(a, b) (c, d) = (ac - \bar{d}b, da + b\bar{c}).$ 

The new algebra  $B = A \oplus A$  is said to be obtained from A via the Cayley-Dickson process. In particular,

$$C = R \oplus R$$
,  $H = C \oplus C$ ,  $Ca = H \oplus H$ 

via the Cayley-Dickson process.

PROPOSITION 5.2. (Jacobson [Ja], 1958). Suppose  $B = A \oplus A$  is obtained from A by the Cayley-Dickson process. Then

1) B is commutative  $\Leftrightarrow A = R$ .

2) B is associative  $\Leftrightarrow$  A is commutative.

3) B is alternative  $\Leftrightarrow$  A is associative.

See [Cu] or [H-L] for details.

From this proposition, we have:

 $C = R \oplus R$  is commutative;

 $H = C \oplus C$  is associative, but not commutative;

 $Ca = H \oplus H$  is alternative, but not associative;

 $Ca \oplus Ca$  is not alternative, hence not a normed division algebra.

THEOREM 5.3. (Hurwitz [Hu 1]). The only normed division algebras are R, C, H and Ca.

One can check directly that R, C, H and Ca are normed, though the calculation for Ca is somewhat lengthy. An alternative argument can be found in [Cu]. That there are no other normed division algebras follows from Propositions 5.1 and 5.2.

We end this section with the following description of all possible automorphisms of the Cayley numbers.

PROPOSITION 5.4. Suppose  $e_1$ ,  $e_2$  and  $e_3$  are orthonormal imaginary Cayley numbers with  $e_3$  orthogonal to  $e_1 e_2$ . Then there exists a unique automorphism of Ca sending  $i = (i, 0) \mapsto e_1$ ,  $j = (j, 0) \mapsto e_2$  and  $\varepsilon = (0, 1) \mapsto e_3$ .

This follows from three applications of Proposition 5.1.

From Proposition 5.4, one concludes that the group of all automorphisms of the Cayley numbers (a Lie group known as  $G_2$ ) is 14-dimensional.

6. The Hopf Fibration  $S^7 \hookrightarrow S^{15} \to S^8$ 

Choose orthonormal coordinates in  $R^{16}$  and identify it with Cayley 2-space  $Ca^2$ . In  $Ca^2$  consider subsets of the form

$$L_m = \{(u, mu) : u \in Ca\} \quad \text{for each} \quad m \in Ca,$$
$$L_\infty = \{(0, v) : v \in Ca\}.$$

They are 8-dimensional real linear subspaces of  $R^{16}$ , but not Cayley subspaces of  $Ca^2$  because they are not closed under Cayley multiplication. This is the effect of the nonassociativity of the Cayley numbers. Nevertheless, we call  $L_m$  and  $L_{\infty}$  Cayley lines for simplicity.

We need to check that these Cayley lines fill out  $Ca^2$ , with any two meeting only at the origin. Given  $(u, v) \in Ca^2$ , if u = 0 then this point is on the Cayley line  $L_{\infty}$ . If  $u \neq 0$ , let  $m = v u^{-1}$ . Then  $m u = (v u^{-1}) u = v$ by Fact 3 of the preceding section. Hence the point (u, v) lies on the Cayley line  $L_m$ . Thus the Cayley lines fill out  $Ca^2$ .

Clearly  $L_{\infty}$  meets each other Cayley line only at the origin. And if the point (u, v), with  $u \neq 0$ , lies on the Cayley lines  $L_m$  and  $L_n$ , then v = m u = n u. Hence m = n. Thus any two Cayley lines meet only at the origin.

The unit 7-spheres on these Cayley lines then define for us the Hopf fibration  $S^7 \hookrightarrow S^{15} \to S^8$ . Note that the base space is clearly homeomorphic to an 8-sphere, since there is one Cayley line for each Cayley number m, and one for the number  $\infty$ .

In a similar fashion, if we start with any k-dimensional normed division algebra K, we obtain a Hopf fibration

$$S^{k-1} \hookrightarrow S^{2k-1} \to S^k$$
.

Note by Hurwitz's theorem that K is isomorphic to R, C, H or Ca, so there are really no new cases.