Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 33 (1987)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SOME EXAMPLES OF GROUP ALGEBRAS WITHOUT FILTRED MULTIPLICATIVE BASIS

Autor: Paris, Luis

Kapitel: §3. The group of quaternion units

DOI: https://doi.org/10.5169/seals-87900

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 17.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

since

$$(1-a)(1-b) = (1-b)(1-a) \bmod M^3 .$$

We know that $(1-a)^2, (1-b)^2, (1-a)(1-b)$ forms a K-basis of M^2/M^3. Hence $u \cdot v \neq 0$ and $v \cdot u \neq 0 \bmod M^3$. Otherwise

$$x_1x_2 = y_1y_2 = x_1y_2 + x_2y_1 = 0$$

and $x_1y_2 - x_2y_1 = 0$ contrary to the fact that $\{u, v\}$ gives a basis of M/M^2.

Thus $uv, vu \in B$ satisfy $uv = vu \bmod M^3$ and $uv \neq 0$, $vu \neq 0 \bmod M^3$.

It follows that $uv = vu$. In fact more generally, if $u_1, u_2 \in B \backslash M^k$ and $u_1 = u_2 \bmod M^k$ then $u_1 = u_2$. Proof: $B \cap M^k$ is a basis of M^k, thus $u_1 - u_2 = \sum_{u \in B \cap M^k} \lambda_u u$. This is possible only if $u_1 - u_2 = 0$.

§ 3. The group of quaternion units

Let Q be defined by generators and relations:

$$Q = \langle a, b : a^4 = 1, b^2 = a^2, ab = b^3a \rangle .$$

Set $i = a, j = b, k = ab$ and $c = a^2$. Then

$$Q = \{1, c, i, ci, j, cj, k, ck\} .$$

Proposition 2. *Let K be a field of characteristic 2. The group algebra $K[Q]$ possesses a filtred multiplicative basis if and only if K contains a primitive cube root of unity.*

Proof. If K contains a primitive cube root of unity, say ω, let

$$B = \{1, u, v, uv, vu, u^2, v^2, u^3\} ,$$

where

$$u = \omega i + \omega^2 j + k$$
$$v = \omega^2 i + \omega j + k .$$

It is easily verified that B is a filtred multiplicative basis.

Conversely, suppose that $K[Q]$ possesses a filtred multiplicative basis B.

Observe that $\{1+i, 1+j\}$ is a K-basis of M/M^2, where again $M = \operatorname{rad} K[Q]$. Also $\{1+c, 1+i+j+k\}$ is a K-basis of M^2/M^3. Since $B \cap (M/M^2) = \{u, v\}$ must be a K-basis of $M \bmod M^2$, we have

$$u = x_1(1+i) + y_1(1+j) \bmod M^2 ,$$

$$v = x_2(1+i) + y_2(1+j) \bmod M^2 ,$$

with $x_1, y_1, x_2, y_2 \in K$ and $x_1y_2 + x_2y_1 \neq 0$.
Now

$$\begin{aligned} u \cdot v &= (x_1x_2+y_1y_2+x_2y_1)(1+c) \\ &\quad + (x_1y_2+x_2y_1)(1+i+j+k) \bmod M^3 , \\ v \cdot u &= (x_1x_2+y_1y_2+x_1y_2)(1+c) \\ &\quad + (x_1y_2+x_2y_1)(1+i+j+k) \bmod M^3 . \end{aligned}$$

Therefore,

$$u \cdot v + v \cdot u = (x_1y_2+x_2y_1)(1+c) \bmod M^3 \neq 0 \bmod M^3 ,$$

and so $u \cdot v \neq v \cdot u$.

We must have $uv \in B \cap (M^2 \backslash M^3)$ since the $(1+i+j+k)$-coordinate of $u \cdot v$ is non-zero. Similarly $v \cdot u \in B \cap (M^2 \backslash M^3)$. But $\dim (M^2/M^3) = 2$ and so

$$B \cap (M^2 \backslash M^3) = \{uv, vu\} .$$

Consider the element $u^2 \in M^2$. Either $u^2 = uv$ or $u^2 = vu$ or $u^2 \in M^3$. But $u^2 = (x_1^2+y_1^2+x_1y_1)(1+c) \bmod M^3$.

Since the $(1+i+j+k)$-coordinate of u^2 is 0, we have $u^2 \neq uv$, $u^2 \neq vu$. Hence $u^2 \in M^3$. This implies $u^2 = 0$, and it follows that the quadratic form $x_1^2 + y_1^2 + x_1y_1$ represents 0 non-trivially in K and $w = y_1/x_1$ is a primitive cube root of unity in K.