

|                     |   |
|---------------------|---|
| <b>Zeitschrift:</b> | L'Enseignement Mathématique   |
| <b>Herausgeber:</b> | Commission Internationale de l'Enseignement Mathématique                              |
| <b>Band:</b>        | 33 (1987)   |
| <b>Heft:</b>        | 1-2: L'ENSEIGNEMENT MATHÉMATIQUE  |
| <br>                |   |
| <b>Artikel:</b>     | SOME EXAMPLES OF GROUP ALGEBRAS WITHOUT FILTRÉD MULTIPLICATIVE BASIS                  |
| <b>Autor:</b>       | Paris, Luis   |
| <b>Kapitel:</b>     | §3. The group of quaternion units   |
| <b>DOI:</b>         | <a href="https://doi.org/10.5169/seals-87900">https://doi.org/10.5169/seals-87900</a> |

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

**Download PDF:** 21.05.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

since

$$(1-a)(1-b) = (1-b)(1-a) \bmod M^3.$$

We know that  $(1-a)^2, (1-b)^2, (1-a)(1-b)$  forms a  $K$ -basis of  $M^2/M^3$ . Hence  $u \cdot v \neq 0$  and  $v \cdot u \neq 0 \bmod M^3$ . Otherwise

$$x_1x_2 = y_1y_2 = x_1y_2 + x_2y_1 = 0$$

and  $x_1y_2 - x_2y_1 = 0$  contrary to the fact that  $\{u, v\}$  gives a basis of  $M/M^2$ .

Thus  $uv, vu \in B$  satisfy  $uv = vu \bmod M^3$  and  $uv \neq 0, vu \neq 0 \bmod M^3$ .

It follows that  $uv = vu$ . In fact more generally, if  $u_1, u_2 \in B \setminus M^k$  and  $u_1 = u_2 \bmod M^k$  then  $u_1 = u_2$ . Proof:  $B \cap M^k$  is a basis of  $M^k$ , thus  $u_1 - u_2 = \sum_{u \in B \cap M^k} \lambda_u u$ . This is possible only if  $u_1 - u_2 = 0$ .

### § 3. THE GROUP OF QUATERNION UNITS

Let  $Q$  be defined by generators and relations:

$$Q = \langle a, b : a^4 = 1, b^2 = a^2, ab = b^3a \rangle.$$

Set  $i = a, j = b, k = ab$  and  $c = a^2$ . Then

$$Q = \{1, c, i, ci, j, cj, k, ck\}.$$

**PROPOSITION 2.** *Let  $K$  be a field of characteristic 2. The group algebra  $K[Q]$  possesses a filtered multiplicative basis if and only if  $K$  contains a primitive cube root of unity.*

*Proof.* If  $K$  contains a primitive cube root of unity, say  $\omega$ , let

$$B = \{1, u, v, uv, vu, u^2, v^2, u^3\},$$

where

$$u = \omega i + \omega^2 j + k$$

$$v = \omega^2 i + \omega j + k.$$

It is easily verified that  $B$  is a filtered multiplicative basis.

Conversely, suppose that  $K[Q]$  possesses a filtered multiplicative basis  $B$ .

Observe that  $\{1+i, 1+j\}$  is a  $K$ -basis of  $M/M^2$ , where again  $M = \text{rad } K[Q]$ . Also  $\{1+c, 1+i+j+k\}$  is a  $K$ -basis of  $M^2/M^3$ . Since  $B \cap (M/M^2) = \{u, v\}$  must be a  $K$ -basis of  $M \bmod M^2$ , we have

$$u = x_1(1+i) + y_1(1+j) \bmod M^2,$$

$$v = x_2(1+i) + y_2(1+j) \bmod M^2,$$

with  $x_1, y_1, x_2, y_2 \in K$  and  $x_1y_2 + x_2y_1 \neq 0$ .

Now

$$\begin{aligned} u \cdot v &= (x_1x_2 + y_1y_2 + x_2y_1)(1+c) \\ &\quad + (x_1y_2 + x_2y_1)(1+i+j+k) \bmod M^3, \end{aligned}$$

$$\begin{aligned} v \cdot u &= (x_1x_2 + y_1y_2 + x_1y_2)(1+c) \\ &\quad + (x_1y_2 + x_2y_1)(1+i+j+k) \bmod M^3. \end{aligned}$$

Therefore,

$$u \cdot v + v \cdot u = (x_1y_2 + x_2y_1)(1+c) \bmod M^3 \neq 0 \bmod M^3,$$

and so  $u \cdot v \neq v \cdot u$ .

We must have  $uv \in B \cap (M^2 \setminus M^3)$  since the  $(1+i+j+k)$ -coordinate of  $u \cdot v$  is non-zero. Similarly  $v \cdot u \in B \cap (M^2 \setminus M^3)$ . But  $\dim(M^2/M^3) = 2$  and so

$$B \cap (M^2 \setminus M^3) = \{uv, vu\}.$$

Consider the element  $u^2 \in M^2$ . Either  $u^2 = uv$  or  $u^2 = vu$  or  $u^2 \in M^3$ .

But  $u^2 = (x_1^2 + y_1^2 + x_1y_1)(1+c) \bmod M^3$ .

Since the  $(1+i+j+k)$ -coordinate of  $u^2$  is 0, we have  $u^2 \neq uv$ ,  $u^2 \neq vu$ .

Hence  $u^2 \in M^3$ . This implies  $u^2 = 0$ , and it follows that the quadratic form  $x_1^2 + y_1^2 + x_1y_1$  represents 0 non-trivially in  $K$  and  $w = y_1/x_1$  is a primitive cube root of unity in  $K$ .