

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 33 (1987)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: RADICALS AND HILBERT NULLSTELLENSATZ FOR NOT NECESSARILY ALGEBRAICALLY CLOSED FIELDS
Autor: Laksov, Dan
Kapitel: §1. The main result and definitions
DOI: <https://doi.org/10.5169/seals-87902>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 12.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

RADICALS AND HILBERT NULLSTELLENSATZ
FOR NOT NECESSARILY ALGEBRAICALLY CLOSED FIELDS

by Dan LAKSOV

§ 1. THE MAIN RESULT AND DEFINITIONS

We shall in the following fix a (commutative) field k and denote by \bar{k} the algebraic closure of k . Moreover, we shall denote by K a subfield of \bar{k} containing k . The polynomial ring $k[x_1, x_2, \dots, x_r]$ in r variables over k we denote by R . Given an ideal I in R , we denote by $Z_K(I)$ the algebraic subset of the r -dimensional affine space A_K^r which consists of the common zeroes of the polynomials in I . That is

$$Z_K(I) = \{(a_1, a_2, \dots, a_r) \mid a_i \in K \quad \text{for} \quad i = 1, 2, \dots, r \\ \text{and} \quad f(a_1, a_2, \dots, a_r) = 0 \quad \text{for all} \quad f \in I\}.$$

The HILBERT NULLSTELLENSATZ is usually stated as follows:

Given an ideal I in R and a polynomial f of R , then f vanishes at all points of $Z_{\bar{k}}(I)$ if and only if $f^n \in I$ for some positive integer n .

In symbols the Hilbert Nullstellensatz can be written in the form:

$$\sqrt{I} = \{f \in R \mid Z_{\bar{k}}(f) \supseteq Z_{\bar{k}}(I)\}.$$

Here \sqrt{I} denotes the *radical* of I , defined as the intersection of all prime ideals containing I , or equivalently by

$$\sqrt{I} = \{f \in R \mid f^n \in I \quad \text{for some positive integer} \quad n\}.$$

As an immediate consequence of the Hilbert Nullstellensatz we obtain the following result which is often referred to as the **WEAK HILBERT NULLSTELLENSATZ**:

Given an ideal I in R , then I is not all of R if and only if $Z_{\bar{k}}(I)$ is non-empty.

The Hilbert Nullstellensatz is one of the fundamental algebraic tools in geometry because it leads to a dictionary between algebraic subsets

of \mathbf{A}_k^r on the one hand and radical ideals in R on the other. In algebraic geometry over a not-necessarily algebraically closed field K the main objects of study are the algebraic subsets of the affine space \mathbf{A}_K^r . However, if K is not algebraically closed, there always exists ideals in R with no zeroes in \mathbf{A}_K^r . Hence the above correspondence between radical ideals and algebraic sets fails, even in the sense of the weak Nullstellensatz.

The purpose of this article is to prove a generalization of the Hilbert Nullstellensatz which makes it possible to set up a dictionary between the algebraic subsets of \mathbf{A}_K^r on the one hand and certain ideals of R , that we shall call K -radical, on the other.

To state our main result it is convenient to introduce the following notation:

Let y_1, y_2, \dots be a countably infinite set of elements that are algebraically independent over k . We denote by $P_K(m)$ the set of homogeneous polynomials in $k[y_1, y_2, \dots, y_m]$ whose zeroes in \mathbf{A}_K^m , if any, are of the form $(a_1, a_2, \dots, a_{m-1}, 0)$. That is,

$$P_K(m) = \{p \in k[y_1, y_2, \dots, y_m] \mid p \text{ is homogeneous and } Z_K(p) \subseteq Z_K(y_m)\}$$

Let A be a k -algebra and I an ideal of A . We denote by $\sqrt{K}I$ the subset $\{a \in A \mid \text{for some positive integer } m \text{ there exists a polynomial } p \in P_K(m) \text{ and elements } a_1, a_2, \dots, a_{m-1} \text{ of } A \text{ such that } p(a_1, a_2, \dots, a_{m-1}, a) \in I\}$.

Below we shall prove that $\sqrt{K}I$ is an ideal in A which we call the K -radical of I . We can now state the main result of this article, which we shall refer to as the HILBERT K -NULLSTELLENSATZ as follows:

Given an ideal I of R , then

$$\sqrt{K}I = \{f \in R \mid Z_K(f) \supseteq Z_K(I)\}.$$

As an immediate consequence of the Hilbert K -Nullstellensatz we obtain the following result which we refer to as the WEAK HILBERT K -NULLSTELLENSATZ:

Given an ideal I of R , then $\sqrt{K}I$ is not all of T if and only if $Z_K(I)$ is non-empty.

We observe that the Hilbert Nullstellensatz and its weak form are the Hilbert \bar{k} -Nullstellensatz and the weak Hilbert \bar{k} -Nullstellensatz. Indeed, it is clear that we have

$$P_{\bar{k}}(m) = \{1, y_m, y_m^2, y_m^3, \dots\} \quad \text{for } m = 1, 2 \dots$$

Hence it follows from the definition of $\sqrt[K]{I}$ that, if $K = \bar{k}$ then $\sqrt[K]{I} = \sqrt{I}$ is the usual radical of I .

A result in the direction of the Hilbert Nullstellensatz was given by D. W. Dubois [2] and J. J. Risler [7] when k is ordered and K the real closure of k . A similar weaker result, which is however valid for any field k , when $k = K$, was given by W. A. Adkins, P. Gianni and A. Tognoli [1]. We shall return to these results in § 4 and see how they relate to the results of this article. In that section we also discuss some open problems related to the previous work.

In the process of generalizing the Hilbert Nullstellensatz we introduce, for each pair of fields k and K with $k \subseteq K$, the K -radical of an ideal in any k -algebra. The K -radical of ideals in R makes it possible to give a treatment of the Nullstellensatz over an arbitrary field which is analogous to the traditional presentation over algebraically closed fields. Most properties that hold for the usual radical of an ideal can be seen to hold for the K -radical and the K -radical merits some interest of its own. Below we shall however only give those properties needed in our presentation of the Nullstellensatz. These properties we have collected in § 2. For a more complete treatment see Laksov [4] and [5].

The results of Dubois and Risler strongly suggest that the K -radical of an ideal can be defined by much smaller sets of polynomials than the sets $P_K(m)$. Restricting the set of polynomials used to define the K -radical would be the first step towards generalizing Hilbert's 17'th problem and would give extremely interesting information about the fields involved. We shall however show that even modest advances in this direction may be very difficult. To be more precise we introduce, for each natural number m a set

$$P_K^0(m) = \{p \in k[y_1, y_2, \dots, y_m] \mid p \text{ is homogeneous and the only zero of } p \text{ in } \mathbf{A}_K^m \text{ is the origin}\}$$

and for any ideal I in R we define a subset I_T of R by

$$I_T = \{f_i \in R \mid \text{for some positive integer } m \geq i \text{ there exists a polynomial } p \in P_K^0(m) \text{ and elements } f_1, f_2, \dots, f_m \text{ in } R \text{ such that } p(f_1, f_2, \dots, f_m) \in I\}$$

Then $P_K^0(m) \subseteq P_K(m)$ and consequently $I_T \subseteq \sqrt[K]{I}$. Moreover we have that $I \subseteq I_T$. The definition of I_T is apparently more natural and symmetric than that of $\sqrt[K]{I}$.

An intriguing problem raised by Tognoli is:

For which pair of fields $k \subseteq K$ do we have that $I_T = \sqrt[k]{I}$ for all ideals I of R ?

It was long conjectured that equality holds for all pairs of fields (at least when the characteristics of k is zero). We shall however, in section 5, give examples showing that one may have strict inequality $I_T \subset \sqrt[k]{I}$ for the two pairs $k = K = \mathbf{Z}/2\mathbf{Z}$ and $k = K = \mathbf{Q}$.

Before we proceed (in § 3) to prove the Hilbert K -Nullstellensatz we shall in § 2 collect all the results that we need about the K -radicals and the polynomials $P_K(m)$ in the next section.

§ 2. SOME PROPERTIES OF THE K -RADICAL

We shall denote by $S(m)$ the polynomial ring $k[y_1, y_2, \dots, y_m]$.

LEMMA 1. Let $p \in P_K(m)$ and $q \in P_K(n)$. For each polynomial $s = s(y_1, y_2, \dots, y_{m+n}) \in S(m+n)$ of degree one less than q , we have that,

$$r = p(y_1 \cdot s, y_2 \cdot s, \dots, y_{m-1} \cdot s, q(y_{m+1}, y_{m+2}, \dots, y_{m+n})) \in P_K(m+n).$$

Proof. It is clear that r is a homogeneous polynomial in $S(m+n)$. Let $(a_1, a_2, \dots, a_{m+n}) \in \mathbf{A}_K^{m+n}$ be a zero of r . Since $p \in P_K(m)$, we have that $q(a_{m+1}, a_{m+2}, \dots, a_{m+n}) = 0$. However, we have that $q \in P_K(m)$ so that $a_{m+n} = 0$. Consequently $r \in P_K(m+n)$ as asserted.

PROPOSITION 2. Let A be a k -algebra and I an ideal of A . Then the K -radical $\sqrt[k]{I}$ of I is an ideal of A (possibly A itself) which contains the radical of I .

Proof. Since $P_K(1) = \{1, y_1, y_1^2, \dots\}$ it is clear that the set $\sqrt[k]{I}$ contains \sqrt{I} .

Let f and g be elements in $\sqrt[k]{I}$. Then by the definition of the K -radical there are positive integers m and n , polynomials $p \in P_K(m)$ and $q \in P_K(n)$ and elements f_1, f_2, \dots, f_{m-1} and g_1, g_2, \dots, g_{n-1} of A such that

$$p(f_1, f_2, \dots, f_{m-1}) \in I \quad \text{and}$$

$$q(g_1, g_2, \dots, g_{n-1}, g) \in I$$

Let h be an element of A and let d be the degree of p . Then we have that

$$p(hf_1, hf_2, \dots, hf_{m-1}, hf) = h^d p(f_1, f_2, \dots, f_{m-1}, f) \in I.$$