Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 33 (1987)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: NONLINEAR ANALYSIS IN GEOMETRY

Autor: Yau, Shing Tung

Kapitel: 3. Hermitian Manifolds and Stable Vector Bundles

DOI: https://doi.org/10.5169/seals-87888

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 25.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

144 S. T. YAU

3. HERMITIAN MANIFOLDS AND STABLE VECTOR BUNDLES

We will consider canonical metrics on compact complex manifolds which are not necessarily Kählerian. For Hermitian manifolds in general, it is difficult to find canonical metrics because the Hermitian connection has torsion and hence is not Riemannian. Therefore one would like to assume extra conditions on M. Let g be a Hermitian metric on M and ω its Kähler form. One natural condition is to assume that

$$\partial \overline{\partial}(\omega^{n-1}) = 0,$$

which is weaker than the condition of being Kähler. One would like to put more conditions on g, besides (1), to make the metric more canonical. Motivated by the theory of supersymmetry, Hull and Witten [HW] proposed the following condition on ω . Locally one should be able to write ω as $\partial\theta + \bar{\partial}\bar{\theta}$ where θ is a (0, 1) form. Notice that if ω is Kähler, it can always be written as $\partial\bar{\partial}f$.

Let us now demonstrate that the above condition is equivalent to the condition $\partial \overline{\partial} \omega = 0$. Clearly, we have only to prove the condition $\partial \overline{\partial} \omega = 0$ implies that ω can be written in the above form. As $\overline{\partial} \omega$ is a closed form, it is locally exact. By comparting the types, we can find a (0,2) form Ω and a (1,1) form ω , so that $\overline{\partial} \omega = \partial \Omega + \overline{\partial} \omega'$ with $\overline{\partial} \Omega = 0$ and $\partial \omega' = 0$. Noticing that $\omega = \overline{\omega}$, we can then prove that $\omega - \omega' - \overline{\omega}' - \Omega - \overline{\Omega}$ is a closed form. Therefore, locally it is exact and we can find a (0,1) form so that $\omega - \omega' - \overline{\omega}' = \partial \theta + \overline{\partial} \overline{\theta}$. Since $\partial \omega' = 0$, locally ω' is ∂ -exact and we have proved locally ω is the form that we seek.

Recently Todorov observed that any compact complex manifold admits a Hermitian form ω with $\partial \overline{\partial} \omega = 0$. Therefore it seems that for any compact complex manifold, it is of interest to study the group obtained by taking the quotient of (1, 1) form ω with $\partial \overline{\partial} \omega = 0$ by the subgroup cosets of $\partial \theta + \overline{\partial} \overline{\theta}$ where θ is globally defined (0, 1) form.

Now let V be a holomorphic vector bundle over a compact manifold M with the property $\partial \bar{\partial}(\omega^{n-1}) = 0$. We can define the degree of the bundle V with respect to ω by

$$\deg_{\omega} V = \int_{M} \Xi_{1}(V) \wedge \omega^{n-1}.$$

where $\Xi_1(V)$ denotes the Ricci form of the bundle V. Since $\partial \bar{\partial}(\omega^{n-1}) = 0$, this definition is independent of the choice of metric on V.

In [U-Y], Uhlenbeck and Yau proved the following:

(2) Suppose V is a holomorphic vector bundle over a compact Kähler manifold M. If V is stable, i.e., $\frac{\deg_{\omega}V'}{\operatorname{rank}V'} < \frac{\deg_{\omega}V}{\operatorname{rank}V}$ for every coherent subsheaf $V' \subseteq V$ such that $0 < \operatorname{rank}(V') < \operatorname{rank}(V)$, then there exists a Hermitian-Einstein metric on V which is unique up to a constant.

Conversely, the existence of a Hermitian-Einstein metric on V implies that V is direct sum of stable bundles. This was proved by Kobayashi and Lübke [Lu]. Moreover, it is likely that the condition M be Kähler can be replaced by (1). It should be noted that the above theorem was proved by Donaldson [D2] for algebraic surfaces.

We now state some corollaries of (2). First of all, the symmetric tensor product bundle of a stable holomorphic vector bundle is also stable. Secondly, if V is a stable bundle, then for r = rank (V),

(3)
$$\int_{M} (2r c_{2}(V) - (r-1)c_{1}^{2}(V)) \wedge \omega^{n-2} \geq 0,$$

and equality holds if and only if up to finite cover of M, V is a direct sum of line bundles (when n=2, this was due to Bogomolov [Bo]) without dealing with the case of equality. Therefore, if $c_1^2(V)=0$ then

$$\int_{M} c_{2}(V) \wedge \omega^{n-2} \ge 0$$
 and equality holds if and only if V is flat and unique

up to a scalar. These results are in fact generalizations of those in the Riemann surface case. In particular, let V be a holomorphic vector bundle over a Riemann surface Σ_g . Then V is stable and $c_1(V)=0$ if and only if there exists a Hermitian metric on V with zero curvature, i.e., if and if there is a unitary representation of $\pi_1(\Sigma_g)$ (see Narashimhan and Seshadri [N-S] for details.

We now consider the moduli space of stable vector bundles. Let M(r, d) be a complete family of stable vector bundles of fixed rank r and fixed degree d over a Riemann surface Σ_g . Can one prove that $c_1(M_g) > 0$, in particular, can one construct a Kähler metric on M_g with positive Ricci curvature? Cho [Co] proved that there exists a Kähler metric on $M_g(r, d)$ with nonnegative holomorphic sectional curvature. However, even the positivity of the holomorphic sectional curvature does not imply the positivity of the Ricci curvature. For example, let H be the hyperplane bundle over \mathbb{CP}^1 and (1) the trivial line bundle. Then the Hirzebruch surfaces M_d

146 S. T. YAU

= $P(H^d+(1))$ have Kähler metrics with positive holomorphic sectional curvature. On the other hand, for $d \ge 3$, M_d does not have positive first Chern class.

4. CHERN NUMBER INEQUALITIES

In 1976, the author proved the Calabi conjecture and demonstrated the following Chern number inequality for algebraic manifolds with either ample or trivial canonical line bundles:

$$(*) (-1)^n c_2 c_1^{n-2} \geqslant \frac{(-1)^n}{2(n+1)} c_1^n$$

where equality holds if and only if M is covered by the ball, i.e., $M = B/\Gamma$ for some $\Gamma \subseteq SU(n, 1)$. Around the same time, Miyaoka [M3], extending the method of Bogomolov, obtained the same inequality for n = 2 under the weaker assumption that the Kodaira dimension of the surface is nonnegative. However, he has not shown that equality holds if and only if M is covered by the ball.

By studying surfaces with singularities, Cheng and Yau [C-Y2] proved inequality (*) for surfaces of general type (equality holds if and only if M^2 is covered by the ball). The arguments in [C-Y2] can also be generalized to higher dimensions. One can also characterize surfaces M which are biholomorphic to B^n/Γ where $\Gamma \subseteq SU(2, 1)$ is allowed to have fixed points. Note that M is, in general, a variety since Γ may have fixed points.

It is also interesting to study manifolds which satisfy certain Chern number inequalities. Surfaces which satisfy inequality (*) have been studied by Hirzebruch, Deligne, Mostow, etc. A corollary of [Y2] is the following rigidity theorem for Kählerian structures on \mathbb{CP}^n : The only Kählerian structure on \mathbb{CP}^n is the standard one; moreover, the only complex structure on \mathbb{CP}^2 is the standard one. For n odd, this result was due to Hirzebruch and Kodaira [H-K].

We now sketch the proof of inequality (*) when the canonical line bundle K of M is ample. In this case, there exists a Kähler-Einstein metric on K. For Kähler-Einstein metrics one observes that the Chern integral associated to the left hand side of (*) can be expressed in terms of the length squared of the curvature tensor. Since the Ricci tensor is the only part of the curvature tensor, the right hand side, which can be written as the determinant of the Ricci tensor, can be dominated by the left hand side.