

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 33 (1987)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: INVOLUTIONS IN SURFACE MAPPING CLASS GROUPS
Autor: McCarthy, John / Papadopoulos, Athanase
Kapitel: 1. Introduction
DOI: <https://doi.org/10.5169/seals-87897>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

INVOLUTIONS IN SURFACE MAPPING CLASS GROUPS

by John McCARTHY and Athanase PAPADOPOULOS

1. INTRODUCTION

Let F be a compact orientable surface with negative Euler characteristic. A *mapping class* is an isotopy class of orientation-preserving homeomorphisms of F . Mapping classes form a group under composition, the *mapping class group*, $M(F)$. An element of order two of the mapping class group will be called an involution. In this article, we prove two theorems about products of involutions.

The first theorem is group-theoretical. We assume that the surface F is closed and we study the subgroup of $M(F)$ which is generated by involutions. In particular, for closed surfaces of genus greater than or equal to three, we prove that the mapping class group is generated by involutions.

The second theorem is geometric. There is a classification, into 3 types, of mapping classes, which is due to Thurston. This theorem is about the type of the product of two involutions. As is natural in this setting, the theorem and its proof are in terms of the action of the mapping class group on Teichmüller space and its Thurston boundary.

The second theorem is analogous in nature to the following elementary facts about the product of order-two elements of a discrete group of isometries of hyperbolic 2 or 3-space:

(i) the product is an elliptic isometry if and only if the two order-2 isometries have a common fixed-point in hyperbolic space.

(ii) if the two order-2 isometries have no common fixed point in hyperbolic space and have a common fixed point on the boundary at infinity, their product is a parabolic isometry.

(iii) if the two isometries do not have any common fixed point, neither in hyperbolic space nor on the boundary at infinity, their product is of hyperbolic type.

In section 2, we prove the group theoretical result. Then, in section 3, we give an outline of some of the background material on Thurston's

classification. For a complete exposition of Thurston's theory, we refer the reader to [4], and for more information on Teichmüller space, to [1]. Finally, in section 4, we prove the theorem on the type of the product of two involutions.

The problem of studying the types of products of involutions in the mapping class group was suggested to the second author by François Laudenbach. The theorem on the subgroup generated by involutions arose out of an attempt to obtain more precise information about the mapping classes which occur as products of two involutions.

The first author acknowledges partial support from Université René Descartes (Strasbourg), and the second author acknowledges support from CNRS (France) and Universidad Nacional Autónoma de México.

2. THE SUBGROUP GENERATED BY INVOLUTIONS

Let $M(F_g)$ denote the mapping class group of a closed orientable surface F_g of genus $g \geq 2$. Let $I(F_g)$ denote the subgroup of $M(F_g)$ which is generated by involutions. We wish to describe $I(F_g)$ as a subgroup of $M(F_g)$. Clearly $I(F_g)$ is a normal subgroup of $M(F_g)$. Hence, we shall give our description in terms of the quotient, $M(F_g)/I(F_g)$.

We begin by recalling some algebraic facts about $M(F_g)$. For a general introduction to the algebraic structure of $M(F_g)$, we refer the reader to [2].

It is a classical fact that $M(F_g)$ is generated by Dehn twists about nonseparating simple closed curves on M . In fact, $M(F_g)$ is generated by a finite number of such twists. The minimum number of twists which is required to generate $M(F_g)$ was given by Humphries [5].




FIGURE 1.