

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 34 (1988)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE SCHUR SUBGROUP OF THE BRAUER GROUP OF A LOCAL FIELD
Autor: Riehm, C.

Bibliographie
DOI: <https://doi.org/10.5169/seals-56585>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 21.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

satisfied. Let the the residue class field of $K(\varepsilon_n)$ have 2^k elements. Set $n' = (2^k)^{2^h} - 1$. Then $n \mid n'$, n' is odd, and $K(\varepsilon_{n'})/K(\varepsilon_n)$ is unramified of degree 2^h . Consider the conditions (i)-(v) with n' instead of n . Then (i) is unchanged, (ii) holds because $n \mid n'$, (iii) holds trivially and (v) holds vacuously because $2^h \mid (K(\varepsilon_{n'}) : K)$. Finally $K(\varepsilon_{n'}) \cap K(\varepsilon_4) = K$ since one is ramified and the other is not, so the non-trivial automorphism of $K(\varepsilon_{4n})/K(\varepsilon_n)$ is the restriction of that of $K(\varepsilon_{4n'})/K(\varepsilon_{n'})$, so (iv) holds also for n' .

We can deduce from this abbreviated form of Janusz' theorem that it is equivalent to Yamada's. Suppose that Janusz' conditions are satisfied, and consider the extension $\mathbf{Q}_2(\varepsilon_{2^{h+1}}, \varepsilon_n)/K$. The inertia subgroup of its Galois group is $\mathcal{g} = \mathcal{G}(\mathbf{Q}_2(\varepsilon_{2^{h+1}}, \varepsilon_n)/K(\varepsilon_n))$, a group of order 4. Suppose that ρ is an extension of the non-trivial automorphism of $\mathbf{Q}_2(\varepsilon_{2^h}, \varepsilon_n)/K(\varepsilon_n)$ to $\mathbf{Q}_2(\varepsilon_{2^{h+1}}, \varepsilon_n)$, so $\rho \in \mathcal{g}$. By condition (iv), there is an integer $a \equiv -1 \pmod{2^h}$ such that $\rho(\varepsilon_{2^{h+1}}) = \varepsilon_{2^{h+1}}^a$. It follows that ρ^2 is the identity. Thus \mathcal{g} is non-cyclic. Conversely suppose that there is an extension $\mathbf{Q}_2(\zeta)/K$ whose inertia subgroup \mathcal{g} is non-cyclic. As we saw in 1., this means that σ_{-1} is in the Galois group of \mathbf{Q}_2^c/K and so its restriction (which we also call σ_{-1}) is in $\mathcal{G}(\mathbf{Q}_2(\varepsilon_{2^h}, \varepsilon_c)/K)$ and is non-trivial. Its fixed field contains $K(\varepsilon_c)$; by Lemma 3.3 of [J], $K(\varepsilon_c, \varepsilon_4) = \mathbf{Q}_2(\varepsilon_{2^h}, \varepsilon_c)$ and so the fixed field is *exactly* $K(\varepsilon_c)$. Thus both (iv) and (ii) are also fulfilled. (i) holds by Lemma 1.

4. *F. Lorenz*, [L], p. 463. His condition for *non-triviality of $S(K)$* is that -1 is a norm in the extension K/\mathbf{Q}_2 . The norm residue symbol in the extension $\mathbf{Q}_2^c/\mathbf{Q}_2$ sends -1 to $\sigma_{-1} \in \mathcal{G}(\mathbf{Q}_2^c/\mathbf{Q}_2)$. Thus it follows from [S], pp. 204-205, that -1 is a norm in K/\mathbf{Q}_2 iff $\sigma_{-1} \in \mathcal{G}(\mathbf{Q}_2^c/K)$.

REFERENCES

- [C-F] CASSELS, J. W. S. and A. FROHLICH. *Algebraic Number Theory*. Thompson Book Co. Inc., Washington (1967).
- [F] FONTAINE, J.-M. Sur la décomposition des algèbres de groupes. *Ann. Sc. Ec. Norm. Sup.* 49 (1971), 121-180.
- [H] HASSE, H. *Zahlentheorie*, 2nd ed. Akademie-Verlag, Berlin, 1963.
- [I] ISAACS, I. M. *Character Theory of Finite Groups*. Academic Press, New York (1976).

- [J] JANUSZ, G. J. Generators for the Schur group of local and global number fields. *Pac. J. Math.* 56 (1975), 525-546.
- [L] LORENZ, F. Zur Schurgruppe eines lokalen Körpers. *Arch. Math.* 32 (1979), 458-468.
- [M] MOLLIN, R. The Schur group of a field of characteristic zero. *Pac. J. Math.* 76 (1978), 471-478.
- [N] NEUKIRCH, J. Über das Einbettungsproblem der algebraischen Zahlentheorie. *Inv. Math.* 21 (1973), 59-116.
- [S] SERRE, J.-P. *Corps Locaux*. Act. Sci. et Ind. 1296, Hermann, Paris (1962).
- [W] WEISS, E. *Cohomology of Finite Groups*. Pure and Appl. Math. 34, Academic Press, New York (1969).
- [Y] YAMADA, T. *The Schur Subgroup of the Brauer Group*. Lecture Notes in Math. 397, Springer-Verlag, Berlin (1974).
- [Z] ZASSENHAUS, H. J. *The Theory of Groups*. 2nd ed. Chelsea, New York, 1949.

(Reçu le 12 décembre 1986)

C. Riehm

Dept. of Mathematics and Statistics
McMaster University
Hamilton, Ontario
Canada L8S 4K1

vide-leer-empty