

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 34 (1988)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: GEODESICS IN THE UNIT TANGENT BUNDLE OF A ROUND SPHERE
Autor: Gluck, Herman
Kapitel: 3. Helices in S^3
DOI: <https://doi.org/10.5169/seals-56596>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

3. HELICES IN S^3

A spherical helix in S^3 is a curve $p(t)$ of constant geodesic curvature and torsion. As in R^3 , two spherical helices of the same curvature and torsion are congruent.

If the curvature is nonzero, then we can define a Frenet frame $T(t)$, $N(t)$, $B(t)$ along $p(t)$ in the usual way, and get the Frenet equations:

$$T' = \kappa N, \quad N' = -\kappa T - \tau B, \quad B' = \tau N.$$

Here we assume that t is an arc length parameter along $p(t)$, and use primes ' to denote covariant differentiation of vector fields along this path.

A model helix in S^3 is given by

$$p(t) = (\cos \alpha \cos at, \cos \alpha \sin at, \sin \alpha \cos bt, \sin \alpha \sin bt).$$

Here α ranges between 0 and $\pi/2$ and determines the shape of the flat torus

$$x_1^2 + x_2^2 = \cos^2 \alpha, \quad x_3^2 + x_4^2 = \sin^2 \alpha,$$

on which the helix $p(t)$ lies. We take the numbers a and b to be ≥ 0 , and require that

$$a^2 \cos^2 \alpha + b^2 \sin^2 \alpha = 1,$$

so that the helix will be traversed at unit speed. Every spherical helix in S^3 is congruent to one of these models.

Next, we give formulas for the curvature κ , torsion τ , and writhe $\rho = \sqrt{\kappa^2 + \tau^2}$ of the model helix $p(t)$ in terms of the descriptive parameters α , a and b . These formulas are given as general information only, and will not be used here.

We first record two simple inequalities which follow from the equality $a^2 \cos^2 \alpha + b^2 \sin^2 \alpha = 1$.

Note that $a = 1$ and $b = 1$ satisfies this equation. So if one of these quantities increases above 1, the other must decrease below 1. Arranging matters so that a is the larger of the two, we will then have

$$(a^2 - 1)(1 - b^2) \geq 0.$$

In addition,

$$a^2 + b^2 \geq a^2 \cos^2 \alpha + b^2 \sin^2 \alpha = 1,$$

so we have

$$a^2 + b^2 - 1 \geq 0.$$

The formulas for curvature, torsion and writhe are as follows.

$$\text{Curvature} = \kappa = \sqrt{(a^2 - 1)(1 - b^2)}$$

$$\text{Torsion} = \tau = ab$$

$$\text{Writhe} = \rho = \sqrt{a^2 + b^2 - 1}.$$

Consider the 3-dimensional linear space of vector fields

$$aT(t) + bN(t) + cB(t)$$

which can be written as constant coefficient combinations of the Frenet vectors along the helix $p(t)$. Covariant differentiation along the helix maps this linear space to itself according to the Frenet formulas.

We've already noted in the introduction that the instantaneous axis vector $U = \tau T - \kappa B$ satisfies $U' = 0$.

Consider the vectors N and $V = (\kappa/\rho)T + (\tau/\rho)B$, which form an orthonormal basis for the orthogonal complement of U . Note that

$$N' = -\kappa T - \tau B = -\rho V, \quad \text{and}$$

$$V' = (\kappa/\rho)T' + (\tau/\rho)B' = (\kappa/\rho)(\kappa N) + (\tau/\rho)(\tau N) = \rho N.$$

Thus, covariant differentiation along the helix kills the instantaneous axis vector and takes the orthogonal 2-plane to itself by a 90° rotation, followed by multiplication by the writhe.

4. SASAKI'S EQUATIONS

Let M be any Riemannian manifold, and UM its unit tangent bundle with the Riemannian metric described in section 1.

THEOREM (Sasaki [Sa], 1958). *The curve $(p(t), v(t))$ in UM is a constant speed geodesic there if and only if both of the following equations hold:*

$$1) \quad v'' = -\langle v', v' \rangle v$$

$$2) \quad p'' = R(v', v)p'.$$

Here, primes denote ordinary derivatives with respect to t when applied to functions, and covariant derivatives along the path $p(t)$ when applied to vector fields. For example, the first prime in p'' represents ordinary differentiation, the second, covariant differentiation. The symbol R denotes the Riemann curvature transformation

$$R: TM_p \times TM_p \rightarrow \text{Hom}(TM_p, TM_p).$$