

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 34 (1988)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** AN ELEMENTARY PROOF OF THE STRUCTURE THEOREM FOR  
CONNECTED SOLVABLE AFFINE ALGEBRAIC GROUPS  
**Autor:** Dokovi, Dragomir Ž.  
**DOI:** <https://doi.org/10.5169/seals-56599>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 08.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## AN ELEMENTARY PROOF OF THE STRUCTURE THEOREM FOR CONNECTED SOLVABLE AFFINE ALGEBRAIC GROUPS

by Dragomir Ž. ĐOKOVIĆ <sup>1)</sup>

### ABSTRACT

We give an elementary proof of the basic structure theorem for connected solvable affine algebraic groups  $G$  over an algebraically closed field  $k$ . The main feature of our proof is that we first establish the important fact that the centralizer in  $G$  of a semisimple element  $s$  is connected. Then the main structure theorem follows easily. We also prove that such  $s$  is contained in a maximal torus and that all maximal tori of  $G$  are conjugate. The structure theorem for connected nilpotent affine groups is not needed in the proof; it is obtained at the end as a simple consequence of the main results. In our proof we avoid the use of quotients and Lie algebras of affine groups. On the other hand we use the Lie-Kolchin theorem, Chevalley's theorem, the existence and uniqueness of the Jordan decomposition, and some other elementary facts.

Let  $k$  be an algebraically closed field. All algebraic groups will be defined over  $k$  and are assumed to be affine. By  $N \rtimes H$  we denote the semidirect product of affine algebraic groups where  $N$  is a normal and  $H$  a complementary subgroup. If  $G$  is any affine algebraic group we shall denote by  $G_u$  (resp.  $G_s$ ) the set of all unipotent (resp. semisimple) elements of  $G$ . By  $G^0$  we denote the identity component of  $G$  and by  $G'$  the derived subgroup of  $G$ . A torus  $S$  in  $G$  will be called maximal if  $S \subset T$  implies that  $T = S$  for any torus  $T$  of  $G$ . The center of  $G$  is denoted by  $Z(G)$ . The centralizer of  $s \in G$  resp.  $S \subset G$  in a subgroup  $H \subset G$  will be denoted by  $Z_H(s)$  resp.  $Z_H(S)$ . The existence and uniqueness of the Jordan decomposition for elements of  $G$  will be used without explicit reference. All group homomorphisms will be homomorphisms of affine algebraic groups. For other proofs of the structure theorem for connected solvable affine algebraic groups we refer the reader to the references [1-5].

<sup>1)</sup> This work was supported by the NSERC of Canada Grant A-5285.

1980 *Mathematics Subject Classification*: primary 20G15, secondary 22E25.

The group of all invertible  $n \times n$  upper triangular matrices will be denoted by  $B_n$ . Its subgroup consisting of all diagonal matrices is denoted by  $D_n$ . We have  $B_n = U_n \rtimes D_n$  where  $U_n$  is the closed connected subgroup of  $B_n$  consisting of all unipotent elements of  $B_n$ .

We start with some preliminary facts.

**THEOREM 1 (Lie-Kolchin).** *Every connected solvable affine algebraic group can be embedded in some  $B_n$  as a closed subgroup.*  $\square$

**COROLLARY.** *If  $G$  is a connected solvable affine group then  $G' \subset G_u$ .*  $\square$

**THEOREM 2 (Chevalley).** *If  $N$  is a closed normal subgroup of an affine group  $G$  then there exists a homomorphism  $f: G \rightarrow GL_n(k)$  such that  $\text{Ker } f = N$ .*  $\square$

For the proofs of Theorems 1 and 2 see, for instance, [5, Theorems 6.7 and 5.1.3].

**LEMMA 1.** *If  $f: G \rightarrow H$  is a surjective homomorphism of affine algebraic groups and  $N := \text{Ker } f$  then:*

- (i)  $f(G^0) = H^0$ ;
- (ii)  $f(G_u) = H_u$  and  $f(G_s) = H_s$ ;
- (iii)  $\dim G = \dim N + \dim H$ ;
- (iv) *If  $N$  and  $H$  are connected then  $G$  is connected.*

*Proof.* For the proofs of (i) and (iii) see for instance [4, Section 7.4]. (ii) follows from the fact that  $f$  preserves the Jordan decomposition [4, Theorem 2.4.8]. We shall sketch the proof of (iv). Since  $N$  is connected, we have  $N \subset G^0$ . By (i) we have  $f(G^0) = H^0 = H$ , and consequently  $G = NG^0 = G^0$ .  $\square$

We need a lemma to prove the centralizer theorem. For a more general version of this lemma see [2, Proposition (9.3)].

**LEMMA 2.** *Let  $N$  be a closed normal connected abelian unipotent subgroup of an affine group  $G$  and let  $s \in G_s$ . Then  $M := \{sus^{-1}u^{-1} : u \in N\}$  is a closed connected subgroup of  $N$ , the multiplication map  $\mu: M \times Z_N(s) \rightarrow N$  is bijective, and  $Z_N(s)$  is connected.*

*Proof.* Since  $N$  is abelian, the map  $f: N \rightarrow N$ , defined by  $f(u) = sus^{-1}u^{-1}$ , is a morphism of algebraic groups whose kernel is  $Z_N(s)$  and image  $M$ , so  $M$  is a closed connected subgroup of  $N$ . If  $x \in M \cap Z_N(s)$  then  $x = sus^{-1}u^{-1}$  for some  $u \in N$ . Since  $usu^{-1} = x^{-1}s = sx^{-1}$  is semi-simple and  $x$  is unipotent, the uniqueness of the Jordan decomposition implies that  $x = 1$ . Hence  $M \cap Z_N(s) = 1$  and so  $\mu$  is injective. By Lemma 1 (iii) we have  $\dim N = \dim M + \dim Z_N(s)$ , which implies that the homomorphism  $\mu$  is also surjective, i.e.,  $MZ_N(s) = N$ . The same argument shows that  $MZ_N(s)^0 = N$ , and so  $Z_N(s)$  must be connected.  $\square$

**THEOREM 3.** *If  $G$  is a connected solvable affine group and  $s \in G_s$  then  $Z_G(s)$  is connected and  $G = G_u Z_G(s)$ .*

*Proof.* We use induction on  $\dim G$ . If  $G$  is abelian the assertions are trivial. Otherwise let  $N$  be the last non-trivial term of the derived series of  $G$ . By the Corollary of Theorem 1,  $N$  is unipotent. We now apply Theorem 2 to this  $G$  and  $N$ . Let  $f$  be as in that theorem. We shall write  $\bar{x}$  for  $f(x)$  and  $\bar{G}$  for  $f(G)$ .

Let  $z \in G$  be such that  $\bar{z} \in Z_{\bar{G}}(\bar{s})$ . Then  $szs^{-1}z^{-1} \in N$ . By Lemma 2 there exists  $u \in N$  and  $v \in Z_N(s)$  such that  $szs^{-1}z^{-1} = sus^{-1}u^{-1} \cdot v$ . Since  $v$  commutes with  $u$  and  $s$ , and  $zsz^{-1} = v^{-1} \cdot usu^{-1}$ , it follows that  $v = 1$ . Thus  $u^{-1}z \in Z_G(s)$  and consequently we have a short exact sequence

$$1 \rightarrow Z_N(s) \hookrightarrow Z_G(s) \rightarrow Z_{\bar{G}}(\bar{s}) \rightarrow 1.$$

By Lemma 2,  $Z_N(s)$  is connected. By Lemma 1 (iii) we have  $\dim \bar{G} < \dim G$ . By induction hypothesis, we conclude that  $Z_{\bar{G}}(\bar{s})$  is connected and that  $\bar{G} = (\bar{G})_u \cdot Z_{\bar{G}}(\bar{s})$ . Now Lemma 1 (iv) implies that  $Z_G(s)$  is connected. By part (ii) of the same lemma we have  $f(G_u) = (\bar{G})_u$  and so  $f(G_u Z_G(s)) = (\bar{G})_u Z_{\bar{G}}(\bar{s}) = \bar{G}$ . Since  $N \subset G_u$ , it follows that  $G = G_u Z_G(s)$ .  $\square$

We now proceed to prove the main results about the structure of connected solvable affine groups. But first we need two lemmas.

**LEMMA 3.** *Let  $S \subset B_n$  be a commuting set of semisimple elements. Then there exists  $b \in B_n$  such that  $b^{-1}Sb \subset D_n$ .*

*Proof.* It is an elementary fact of linear algebra that there exists  $a \in GL_n(k)$  such that  $a^{-1}Sa \subset D_n$ . Hence if  $M_n(k)$  is the algebra of  $n$  by  $n$  matrices over  $k$  and  $A$  its subalgebra generated by  $S$ , we know that  $A$  is semisimple (and commutative). Let  $V := k^n$  be the space of column

vectors and let  $e_1, \dots, e_n$  be its standard basis. We shall view  $V$  as a left  $M_n(k)$ -module via matrix multiplication. The subspace  $V_i$  spanned by the vectors  $e_1, \dots, e_i$  is an  $A$ -submodule of  $V$  for each  $i$ . Since  $A$  is semisimple, there exist  $v_i \in V_i \setminus V_{i-1}$ ,  $1 \leq i \leq n$ , such that  $Av_i = kv_i$ . Thus if  $b$  is the matrix whose  $i$ -th column is  $v_i$ ,  $1 \leq i \leq n$ , then  $b \in B_n$  and  $b^{-1}Sb \subset D_n$ .  $\square$

LEMMA 4. *If  $G$  is a connected solvable affine group,  $T \subset G_s$  a closed subgroup, and  $G = G_u T$  then  $T$  is a torus and  $G = G_u \rtimes T$ .*

*Proof.* By the Lie-Kolchin theorem we may assume that  $G$  is a closed subgroup of some  $B_n$ . By using the projection map  $B_n \rightarrow D_n$  we obtain a short exact sequence  $1 \rightarrow G_u \hookrightarrow G \xrightarrow{p} D \rightarrow 1$ , where  $D \subset D_n$  is a torus. Since  $D = p(G) = p(G_u T) = p(T)$ , Lemma 1 (i) implies that  $p(T^0) = D$ . Thus  $G = G_u T^0$  and using  $T \cap G_u = 1$  we conclude that  $T = T^0$ . In particular  $T$  is abelian and by Lemma 3 we may assume that  $T \subset D_n$ , i.e.,  $T = D$ . Since  $B_n = U_n \rtimes D_n$ ,  $G_u \subset U_n$ ,  $T = D \subset D_n$ , and  $G = G_u T$ , it follows that  $G = G_u \rtimes T$ .  $\square$

THEOREM 4. *Let  $G$  be a connected solvable affine group. Then  $G = G_u \rtimes T$  where  $T$  is a maximal torus. In particular,  $G_u$  is connected.*

*Proof.* We use induction on  $\dim G$ . Assume first that  $G_s \subset Z(G)$ . Then  $G_s = Z(G)_s$  is a closed subgroup of  $G$  and  $G = G_u G_s$ . The assertion then follows from Lemma 4. Now assume that there exists  $s \in G_s \setminus Z(G)$ . Then  $Z_G(s)$  is a proper closed subgroup of  $G$ , see e.g. [4, Section 8.2]. By Theorem 3 it is connected and  $G = G_u Z_G(s)$ . By induction hypothesis there exists a torus  $T$  such that  $Z_G(s) = Z_G(s)_u T$ . Then  $G = G_u Z_G(s) = G_u T$  and  $G = G_u \rtimes T$  by Lemma 4.  $\square$

THEOREM 5. *Let  $G = G_u \rtimes T$  be a connected solvable affine group. Then every  $s \in G_s$  is conjugate to an element of  $T$ .*

*Proof.* We use induction on  $\dim G$ . We have  $s = ut$  where  $u \in G_u$  and  $t \in T$ . If  $G$  is abelian then  $u = 1$  and  $s = t$ . Otherwise let  $N$  be the last non-trivial term of the derived series of  $G$ . By the corollary of Theorem 1 we have  $N \subset G_u$ . Hence  $N$  is a closed connected normal abelian unipotent subgroup of  $G$ . By Theorem 2 and the induction hypothesis there exists  $x \in G$  such that  $xsx^{-1} = tv$  where  $v \in N$ . By Lemma 2,  $v = t^{-1}utu^{-1}z$  where  $u \in N$  and  $z \in Z_N(t)$ . Hence  $xsx^{-1} = utu^{-1}z$ . Since  $xsx^{-1}, utu^{-1} \in G_s$ ,  $z \in G_u$ , and  $z$  commutes with  $u$  and  $t$ , it follows that  $z = 1$  and consequently  $xsx^{-1} = utu^{-1}$ .  $\square$

THEOREM 6. If  $G = G_u \rtimes T$  is a connected solvable affine group and  $S \subset G_s$  is a commuting set then  $Z_G(S)$  is connected and  $aSa^{-1} \subset T$  for some  $a \in G$ . In particular, all maximal tori of  $G$  are conjugate.

*Proof.* We use induction on  $\dim G$ . The assertions are obvious if  $S \subset Z(G)$ . Otherwise choose  $s \in S \setminus Z(G)$ . By Theorem 5 we may assume that  $s \in T$ . Then  $Z_G(s)$  is a proper closed subgroup of  $G$  containing  $T$  and  $S$ . By Theorem 3,  $Z_G(s)$  is connected. Since  $\dim Z_G(s) < \dim G$ , we can apply the induction hypothesis to conclude the proof.  $\square$

It is now easy to describe connected nilpotent affine groups.

THEOREM 7. A connected solvable affine group  $G = G_u \rtimes T$  is nilpotent if and only if  $G_s = T \subset Z(G)$ . In that case  $G = G_u \times T$ .

*Proof.* Assume that  $G$  is nilpotent. We prove that  $G_s = T \subset Z(G)$  by induction on  $\dim G$ . We may assume that  $G$  is not abelian. Let  $N$  be the last non-trivial term of the lower central series of  $G$ . Let  $f$  be as in Theorem 2 and  $\bar{G} = f(G)$ . Then  $\bar{G} = f(G_u T) = (\bar{G})_u f(T)$ . By induction hypothesis we have  $f(T) = (\bar{G})_s \subset Z(\bar{G})$ . Consequently if  $t \in T$  and  $x \in G$  then  $u := txt^{-1}x^{-1} \in N$ . Since  $N \subset Z(G) \cap G_u$ , and  $xtx^{-1} = u^{-1}t = tu^{-1}$  we must have  $u = 1$ . Thus  $T \subset Z(G)$  and, by Theorem 5,  $G_s = T$ . The converse is obvious.  $\square$

## REFERENCES

- [1] ABE, E. *Hopf Algebras*. Cambridge University Press 1977.
- [2] BOREL, A. *Linear Algebraic Groups*. Notes by H. Bass, New York: W.A. Benjamin 1969.
- [3] HOCHSCHILD, G.P. *Basic Theory of Algebraic Groups and Lie Algebras*. New York-Heidelberg-Berlin: Springer-Verlag 1981.
- [4] HUMPHREYS, J.E. *Linear Algebraic Groups*. New York-Heidelberg-Berlin: Springer-Verlag 1975.
- [5] SPRINGER, T.A. *Linear Algebraic Groups*. Boston-Basel-Stuttgart: Birkhäuser 1981.

(Reçu le 23 janvier 1988)

Dragomir Ž. Đoković

Department of Pure Mathematics  
University of Waterloo  
Waterloo, Ontario, Canada N2L 3G1

**Vide-leer-empty**