

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 34 (1988)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ABOUT THE PROOFS OF CALABI'S CONJECTURES ON COMPACT KÄHLER MANIFOLDS  
**Autor:** Delanoë, Ph. / Hirschowitz, A.  
**Kapitel:** 1. The Monge-Ampère equation  
**DOI:** <https://doi.org/10.5169/seals-56591>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 18.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

note is to analyze how far these tools are necessary for the proof. It turns out that it is possible to reduce the contribution of elliptic theory mainly to a suitable local inverse function theorem for nonlinear elliptic operators acting on smooth functions [22] [11].

The proof presented below deals only with the reduction to the crucial estimates of order zero, two and three, already obtained by Aubin and Yau. Although it is not so clear in [21] [24] these estimates were performed essentially through coordinate free tensor calculus. We show how higher order estimates may be obtained in the same way.

The whole approach applies as well to the corresponding *real* elliptic Monge-Ampère equation on compact Riemannian manifolds [9] and to various generalizations of it. We shall freely use arguments of Calabi [6, 7, 8], Aubin [1, 2, 3], Yau [23, 24], Bourguignon *et al.* [5] [21].

ACKNOWLEDGEMENTS. The second author thanks Otto Forster for drawing him into the subject and François Rouvière for stimulating conversations. Both authors thank Jean-Pierre Bourguignon who originally suggested that they get in contact together.

## 1. THE MONGE-AMPÈRE EQUATION

Let  $X$  be a compact connected finite-dimensional Kähler manifold.  $\omega$  denotes the original  $C^\infty$  Kähler form,  $g$  the corresponding Kähler metric,  $\varphi \in C^\infty(X)$  denotes a  $C^\infty$  real-valued function on  $X$ , and we set

$$\omega' = \omega + \sqrt{-1} \partial \bar{\partial} \varphi$$

where  $\partial$  and  $\bar{\partial}$  are the usual first order differential operators. Let  $g'$  denotes the Kähler metric corresponding to  $\omega'$ . In the sequel, "smooth" means  $C^\infty$ .

If  $g$  and  $g'$  are viewed as morphisms from the antiholomorphic tangent bundle into the holomorphic cotangent bundle  $T^*$ , then  $(g'g^{-1})$  is an endomorphism of  $T^*$  the determinant of which,  $\det(g'g^{-1})$  is a smooth function on  $X$ . The function  $\varphi$  is said to be *admissible* if and only if  $\det(g'g^{-1})$  is strictly positive on  $X$ . One proves easily that if  $\varphi$  is admissible, then  $g'$  is again a (positive definite Kähler) metric e.g. [2], p. 119.

Let  $\lambda \in [0, +\infty)$ . It is convenient to denote by  $A_\lambda$  the subset of  $C^\infty(X)$  consisting in all admissible real-valued smooth functions  $\varphi$  on  $X$ , satisfying, in case  $\lambda = 0$  the further zero average condition

$$\int_X \varphi dX_g = 0,$$

where  $dX_g$  denotes the volume form associated with  $g$ . When  $\lambda > 0$ ,  $A_\lambda$  is an open subset of  $C^\infty(X)$ : indeed, the natural injection  $C^\infty(X) \hookrightarrow C^2(X)$  is continuous, with respect to the Fréchet and Banach topologies;  $A_\lambda$  is the pull back, by this injection of the open set

$$\{\varphi \in C^2(X), \det(g'g^{-1}) > 0\}.$$

*Definition 1.1.* Let  $X$  be a smooth compact manifold,  $V$  a smooth vector bundle on  $X$ ,  $C^\infty(X, V)$  the Fréchet space of smooth sections of  $V$ . A LCFC submanifold of  $C^\infty(X, V)$ , is a locally closed finite codimensional Fréchet submanifold of  $C^\infty(X, V)$ .

The set  $A_0$  is an open subset of the LCFC submanifold

$$\{\varphi \in C^\infty(X), \int_X \varphi dX_g = 0\}.$$

We define the map  $P_\lambda$ , from  $A_\lambda$  to  $C^\infty(X)$ , by

$$P_\lambda(\varphi) = \lambda\varphi - \text{Log det}(g'g^{-1}).$$

The proofs of theorems 0.1 and 0.2 have been reduced to the solution, when  $\lambda \geq 0$ , of the following complex Monge-Ampère equation (e.g. [21], (lecture n° V), [4] p. 143):

$$(1) \quad P_\lambda(\varphi) = f,$$

where  $f \in C^\infty(X)$  is given, and in case  $\lambda$  vanishes, has to satisfy the natural constraint (e.g. [1] p. 403, [24] p. 361, [21] p. 85),

$$\int_X e^{-f} dX_g = \int_X dX_g.$$

In any case,  $f$  ranges in a connected LCFC submanifold  $B_\lambda$  of  $C^\infty(X)$ . To see that  $B_0$  is connected, notice that  $0 \in B_0$  and that given any  $f \in B_0$ , the following path connects  $f$  to 0 in  $B_0$ :

$$t \in [0, 1] \rightarrow f_t =: tf + \text{Log} \left( \int_X e^{-tf} dX_g \right) - \text{Log} \left( \int_X dX_g \right).$$

The derivative of the map  $P_\lambda$  at  $\varphi \in A_\lambda$ , is given by

$$(2) \quad dP_\lambda(\varphi, \delta\varphi) = (\Delta' + \lambda) \delta\varphi$$

where  $\Delta'$  stands for the Laplace operator on functions in the metric  $g'$  [21] p. 96. Classically, it follows from the Maximum Principle [20], the Fredholm Alternative theorem and from the elliptic regularity theory, that  $dP_\lambda(\varphi, \cdot)$  is invertible  $\forall \varphi \in A_\lambda$ , either from  $C^\infty(X)$  to itself when  $\lambda > 0$ , or

from  $\{u \in C^\infty(X), \int u dX_g = 0\}$  to  $\{v \in C^\infty(X), \int v dX_{g'} = 0\}$  ( $dX_{g'}$  denotes the volume form in the metric  $g'$ ) when  $\lambda = 0$ .

For completeness, let us indicate how, for instance theorem 0.2, can be reduced to equation (1) with  $\lambda = 0$ . It is quite straightforward. First of all we are given a cohomology class  $c \in H^2(X, \mathbf{R})$  such that there exists a Kähler form  $\omega$  in  $c$ ; let  $\rho$  be the Ricci form of  $\omega$ :  $\rho \in C_1(X)$ , the first Chern class of  $X$ .

Then we are given  $\gamma \in C_1(X)$  and hence  $f \in C^\infty(X)$  a real function (defined up to an additive constant), which measures the deviation for  $\omega$  from satisfying 0.2:

$$\gamma - \rho = \sqrt{-1} \partial \bar{\partial} f.$$

Now we look for another Kähler form  $\omega' \in c$ , i.e. we look for a smooth real function  $\varphi$  (also defined up to an additive constant), where

$$\omega' - \omega = \sqrt{-1} \partial \bar{\partial} \varphi$$

such that the Ricci form  $\rho'$  of  $\omega'$  coincides with  $\gamma$ .

In other words, we want  $\varphi$  to satisfy

$$\rho' - \rho \equiv \sqrt{-1} \partial \bar{\partial} \varphi,$$

or equivalently, if  $g$  and  $g'$  are the Kähler metrics respectively associated with  $\omega$  and  $\omega'$ ,

$$\partial \bar{\partial} \{-\text{Log det}(g'g^{-1})\} \equiv \partial \bar{\partial} \varphi$$

which immediately yields equation (1) with  $\lambda = 0$ :

$$-\text{Log det}(g'g^{-1}) = f,$$

since anyway  $f$  is only defined up to an additive constant.

As  $\omega$  and  $\omega'$  are cohomologous and closed, so are the corresponding volume forms, therefore  $X$  has same volume measured with the metrics  $g$  and  $g'$ ; this defines completely  $f$ , subject to the natural constraint mentioned above.

## 2. A TOPOLOGICAL LEMMA

In our setting, the continuity method becomes a "surjectivity method" since it is based on the following