

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 34 (1988)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ABOUT THE PROOFS OF CALABI'S CONJECTURES ON COMPACT
KÄHLER MANIFOLDS
Autor: Delanoë, Ph. / Hirschowitz, A.
Kapitel: 3. Local inversion
DOI: <https://doi.org/10.5169/seals-56591>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 16.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

LEMMA 2.1. Let A, B be metric spaces, with $A \neq \emptyset$ and B connected. Let $P: A \rightarrow B$ be a continuous map. Assume:

- (i) P is open,
- (ii) P is proper, that is, for any compact subset K in B , $P^{-1}(K)$ is compact. Then P is surjective.

Proof. We only need to prove that $P(A)$ is closed. Let b_0 be a point in $\overline{P(A)}$. Since B is a metric space, there exists a sequence $(b_i)_{i>0}$ in $P(A)$ converging to b_0 . The subset $K = \{b_0, b_1, b_2, \dots\}$ is compact, hence so is $PP^{-1}(K)$. The latter contains b_1, \dots, b_i, \dots , hence b_0 , and it is obviously contained in $P(A)$. Q.E.D.

In order to make use of this lemma, we shall need some inverse function theorem for (i), and some *a priori* estimates for (ii).

3. LOCAL INVERSION

THEOREM 3.1. Let X be a smooth compact manifold, V and W smooth vector bundles on X , U an open set in $C^\infty(X, V)$, and $P: U \rightarrow C^\infty(X, W)$, a smooth nonlinear elliptic partial differential operator. Let A and B be LCFC submanifolds of U and of $C^\infty(X, W)$ respectively, such that the restriction P_A of P to A , sends A into B . Then the Jacobian criterion holds for P_A , namely, if the derivative of $P_A: A \rightarrow B$ is invertible at $\varphi_0 \in A$, then P_A is a local diffeomorphism near φ_0 .

This is a convenient variant of the Nash-Moser theorem (e.g. [14]) regarding suitable restrictions of elliptic operators. It is established in a separate paper [11] (see also [22]). It relies only on the classical (Banach) inverse function theorem combined with *elliptic regularity*.

Remark 3.2. The Nash-Moser theorem has been studied by many authors, see the bibliography below and further references in [14] [15] [25].

4. PROPERNESS

In view of (2), theorem 3.1 implies that P_λ is open. We want to apply lemma 2.1 in order to prove that P_λ is surjective from A_λ to B_λ . Since $P_\lambda(A_\lambda) \neq \emptyset$ (it contains 0), and since B_λ is connected, this amounts to proving that P_λ is *proper*. Let us explain why *a priori* estimates imply properness.

Concerning subsets in A_λ we have