

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 34 (1988)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ABOUT THE PROOFS OF CALABI'S CONJECTURES ON COMPACT KÄHLER MANIFOLDS
Autor: Delanoë, Ph. / Hirschowitz, A.
Kapitel: 4. Properness
DOI: <https://doi.org/10.5169/seals-56591>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 08.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

LEMMA 2.1. Let A, B be metric spaces, with $A \neq \emptyset$ and B connected. Let $P: A \rightarrow B$ be a continuous map. Assume:

- (i) P is open,
- (ii) P is proper, that is, for any compact subset K in B , $P^{-1}(K)$ is compact. Then P is surjective.

Proof. We only need to prove that $P(A)$ is closed. Let b_0 be a point in $\overline{P(A)}$. Since B is a metric space, there exists a sequence $(b_i)_{i>0}$ in $P(A)$ converging to b_0 . The subset $K = \{b_0, b_1, b_2, \dots\}$ is compact, hence so is $PP^{-1}(K)$. The latter contains b_1, \dots, b_i, \dots , hence b_0 , and it is obviously contained in $P(A)$. Q.E.D.

In order to make use of this lemma, we shall need some inverse function theorem for (i), and some *a priori* estimates for (ii).

3. LOCAL INVERSION

THEOREM 3.1. Let X be a smooth compact manifold, V and W smooth vector bundles on X , U an open set in $C^\infty(X, V)$, and $P: U \rightarrow C^\infty(X, W)$, a smooth nonlinear elliptic partial differential operator. Let A and B be LCFC submanifolds of U and of $C^\infty(X, W)$ respectively, such that the restriction P_A of P to A , sends A into B . Then the Jacobian criterion holds for P_A , namely, if the derivative of $P_A: A \rightarrow B$ is invertible at $\varphi_0 \in A$, then P_A is a local diffeomorphism near φ_0 .

This is a convenient variant of the Nash-Moser theorem (e.g. [14]) regarding suitable restrictions of elliptic operators. It is established in a separate paper [11] (see also [22]). It relies only on the classical (Banach) inverse function theorem combined with *elliptic regularity*.

Remark 3.2. The Nash-Moser theorem has been studied by many authors, see the bibliography below and further references in [14] [15] [25].

4. PROPERNESS

In view of (2), theorem 3.1 implies that P_λ is open. We want to apply lemma 2.1 in order to prove that P_λ is surjective from A_λ to B_λ . Since $P_\lambda(A_\lambda) \neq \emptyset$ (it contains 0), and since B_λ is connected, this amounts to proving that P_λ is *proper*. Let us explain why *a priori* estimates imply properness.

Concerning subsets in A_λ we have

PROPOSITION 4.1. *A subset S in A_λ is relatively compact in A_λ iff its closure \bar{S} in $C^\infty(X)$ lies inside A_λ and S is bounded in $C^\infty(X)$.*

This readily follows from Ascoli theorem which implies the well-known fact [12] (p. 231) that in $C^\infty(X)$ (and in any closed LCFC submanifold of $C^\infty(X)$, such as B_λ , as well) bounded subsets are relatively compact and vice-versa; hence, compact subset of A_λ are nothing but bounded closed strictly interior subsets of A_λ . Explicitely, let us state the

COROLLARY 4.2. *A closed subset S in A_λ is compact if and only if there exists a sequence $(C_i), i \in \mathbb{N}$, of positive numbers, such that for any φ in S the following estimates hold:*

$$\begin{aligned} \|(g')^{-1}\| &=: \sup_X |(g')^{-1}| \leq C_0, \\ \forall i \in \mathbb{N}, \quad \|D^i \varphi\| &=: \sup_X |D^i \varphi| \leq C_i, \end{aligned}$$

where $|\cdot|$ denotes some natural norms of tensors in the original metric g , and $D =: (\nabla, \bar{\nabla})$ is the total covariant differentiation with respect to the metric g .

Proof. Indeed S is closed and bounded. Moreover, since for $\varphi \in S$,

$$\|(g')^{-1}\| \leq C_0$$

all the eigenvalues of $(g')^{-1}$ (which are positive) are uniformly bounded from above, hence those of g' are uniformly bounded from below, in other words:

$$\exists \varepsilon > 0, \quad \forall \varphi \in S, \quad g' \geq \varepsilon g,$$

or equivalently \bar{S} lies strictly inside A_λ . Q.E.D.

In the next sections we will show that if f belongs to some compact (i.e. bounded and closed) subset K of B_λ , defined by a sequence $(K_i), i \in \mathbb{N}$, such that $\|D^i f\| \leq K_i$, then for $\varphi \in A_\lambda$ satisfying $P_\lambda(\varphi) = f$, the following *a priori* estimates hold:

$$\|\varphi\| \leq C_0, \quad \forall i \in \mathbb{N}, \quad \|D^i \nabla \bar{\nabla} \varphi\| \leq C_{i+2}.$$

These estimates imply that P_λ is proper, i.e. that $S = P_\lambda^{-1}(K)$ is compact, according to the following

PROPOSITION 4.3. *Let S be a closed subset in A_λ . Suppose that there exists a sequence $(C_i), i \in \mathbb{N}$, such that for any φ in S , the following estimates hold:*

$$\| \varphi \| \leq C_0, \quad \| P_\lambda(\varphi) \| \leq C_0, \quad \forall i \in \mathbf{N}, \quad \| D^i \nabla \bar{\nabla} \varphi \| \leq C_{i+2}.$$

Then S is compact.

Proof. The first two estimates imply a uniform estimate

$$| \operatorname{Log} \det (g' g^{-1}) | \leq E.$$

The estimate on $\| \nabla \bar{\nabla} \varphi \|$ yields another one:

$$\| g' \| \leq F.$$

These two estimates yield

$$\| (g')^{-1} \| \leq G.$$

Now from $\| D^i \nabla \bar{\nabla} \varphi \| \leq C_{i+2}$ we infer

$$\| D^i \Delta \varphi \| \leq \tilde{C}_{i+2}$$

since D and g^{-1} commute (Δ denotes the Laplacian in the metric g). As Δ performs a continuous linear automorphism of the Fréchet space of smooth functions *with zero average* (by Fredholm theory), the Closed Graph Theorem implies the missing estimates. Q.E.D.

Remark 4.4. Actually we have been considering two *gradings* of $C^\infty(X)$ [14]. The usual one, namely the one defined, $\forall u \in C^\infty(X)$, by

$$\| u \|_0 = \sup_X | u |,$$

$$\| u \|_i = \| u_i \|_{i-1} + \| D^i u \|, \quad i \geq 1,$$

and another one, well-adapted here since the true unknown is a Kähler metric, defined by

$$\| u \|_0^* = \| u \|_0, \quad \| u \|_1^* = \| u \|_1,$$

$$\| u \|_i^* = \| u \|_{i-1}^* + \| D^{i-2}(\nabla \bar{\nabla} u) \|, \quad i \geq 2.$$

Although it is unnecessary for the purpose of proposition 4.3, it can be shown globally (without Schauder theory) that these two gradings are *tamely* equivalent [14] of degree 2 and base 0 [10] (section 5). Hence, they define the same topology.