

<b>Zeitschrift:</b>	L'Enseignement Mathématique
<b>Herausgeber:</b>	Commission Internationale de l'Enseignement Mathématique
<b>Band:</b>	34 (1988)
<b>Heft:</b>	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
 <b>Artikel:</b>	ABOUT THE PROOFS OF CALABI'S CONJECTURES ON COMPACT KÄHLER MANIFOLDS
<b>Autor:</b>	Delanoë, Ph. / Hirschowitz, A.
<b>Kapitel:</b>	6. Coordinate free tensor calculus
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-56591">https://doi.org/10.5169/seals-56591</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

### 5. A PRIORI ESTIMATES: THE ORIGINAL WAY

According to proposition 4.3 we must prove now that, given any sequence of positive real numbers  $(K_i)$ ,  $i \in \mathbf{N}$ , there exists a sequence  $(C_i)$  such that

$$\forall i \in \mathbf{N}, \quad \| D^i P_\lambda(\varphi) \| \leq K_i$$

implies

$$\| \varphi \| \leq C_0, \quad \forall i \in \mathbf{N}, \quad \| D^i \nabla \bar{\nabla} \varphi \| \leq C_{i+2}.$$

These are *a priori* estimates of order zero, two, three, and so on .... In case  $\lambda > 0$ , the  $C^0$  estimate is straightforward [2]. In case  $\lambda = 0$ , it becomes very tricky; proofs simpler than Yau's original one [24] (p. 352-359), based on the idea of uniformly estimating the  $L^p(dX_g)$  norms of  $\varphi$ , may be found in [16] (dimension 2), [3] [21] and [4] (p. 148-149).

Estimates of order two and three are carried out by means of tensor calculus and of the Maximum Principle (for elliptic equations) [20] applied to *suitable* test functions. Though it is not everywhere clear in [21] [24], it is worth noting that the computations can be written intrinsically, i.e. without any reference to a *particular* system of coordinates (e.g. [2]), or even *coordinate free* (see section 6 below).

Further regularity is then recovered by Schauder theory e.g. [5] (lemma 1). In the sequel, we show how further estimates can be carried out instead, *just going ahead with coordinate free tensor calculus*. This occurs actually for any fully nonlinear second order elliptic equation on a compact Riemannian manifold, via a straightforward imitation of the device below.

*Remark 5.1.* It follows from the  $C^2$  *a priori* estimates that the metrics  $g'$  are *a priori* uniformly equivalent to the original metric  $g$  (see e.g. [3], p. 75).

### 6. COORDINATE FREE TENSOR CALCULUS

Even coordinate free tensor calculus needs indices. Usually these indices refer to a *local* frame. Another way is to view these indices *globally* as labelling copies of the holomorphic and antiholomorphic tangent and cotangent bundles. From this point of view, a tensor written with indices is a section of the tensor product of a family of bundles indexed by an *unordered* set of indices (disregarding those indices subject to the summation convention).

We extend the summation convention as follows: we will be concerned only with lower indices. If a letter occurs twice, it refers to a contraction, which is taken with respect to  $g$  or to  $g'$  according to whether the letter occurs with a bar or with a prime. So,

$$\begin{aligned} T_{\dots a \dots \bar{a} \dots} &\text{ stands for } g^{a\bar{b}} T_{\dots a \dots \bar{b} \dots}, \quad \text{while} \\ T_{\dots a \dots a' \dots} &\text{ stands for } g'^{a\bar{b}} T_{\dots a \dots \bar{b} \dots}. \end{aligned}$$

As usual if  $T_{a\dots l}$  is a tensor, further lower indices refer to covariant differentiation (with respect to  $g$ ); so,

$$\begin{aligned} T_{a\dots lm} &\text{ stands for } \nabla_m T_{a\dots l}, \quad \text{while} \\ T_{a\dots l\bar{m}} &\text{ stands for } \bar{\nabla}_{\bar{m}} T_{a\dots l}. \end{aligned}$$

Our indices will be latin letters; greek letters will denote multi-indices. If  $\alpha$  is a multi-index,  $\bar{\alpha}$  will denote the *conjugate* multi-index (for instance if  $\alpha = abc$ , then  $\bar{\alpha} = \bar{a}\bar{b}\bar{c}$ ), while  $|\alpha|$  denotes its length. We shall say that  $\alpha$  is *mixed* if its length is at least two and, among the first two letters, *exactly* one has a bar.

The notations  $D, \nabla, \bar{\nabla}, \| \cdot \|$ , were introduced in section 4.

*Remark 6.1.* Since covariant differentiation (with respect to  $g$ ) and contraction with respect to  $g'$  do not commute, we observe that, for instance, the difference (recall  $g' = g + \nabla\bar{\nabla}\varphi$ )

$$(3) \quad \Phi_{aa'\alpha b} - (\Phi_{aa'\alpha})_b \equiv \Phi_{ac\alpha} \Phi_{a'c'b}$$

does not vanish.

## 7. HIGHER ORDER A PRIORI ESTIMATES: GENERALITIES

We want to prove by induction,

**PROPOSITION 7.1.** *Given  $n \geq 4$ , a sequence  $(K_i)$ ,  $i \in \mathbb{N}$ , and a finite sequence  $C_0, \dots, C_{n-1}$ , there exists  $C_n$  such that:*

$$\begin{aligned} \|\varphi\| &\leq C_0, \quad \forall i = 0, \dots, n-3, \quad \|D^i \nabla \bar{\nabla} \varphi\| \leq C_{i+2} \\ \text{and } \forall i \in \mathbb{N}, \quad \|D^i P_\lambda(\varphi)\| &\leq K_i, \end{aligned}$$

implies

$$\|D^{n-2} \nabla \bar{\nabla} \varphi\| \leq C_n.$$