

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 35 (1989)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: CAUCHY RESIDUES AND DE RHAM HOMOLOGY
Autor: Iversen, Birger
Kapitel: 1. Compact chains
DOI: <https://doi.org/10.5169/seals-57358>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

CAUCHY RESIDUES AND DE RHAM HOMOLOGY

by Birger IVERSEN

This paper represents my third attempt to write up a suitable generalization of the classical Cauchy Residue theorem. As I pushed the earlier versions for naturality and generality I was ultimately lead to a new foundation of de Rham homology free of the theory of distributions but relying on basic sheaf theory much like the Borel Moore homology theory.

Singular homology and de Rham homology agree on a smooth manifold. The whole point in introducing de Rham homology is the possibility of alternative representations of homology classes. This is amply illustrated by the general Cauchy residue formula given at the end of the paper.

I would like to thank the Mittag Leffler Institute at Stockholm for hospitality while this paper was worked out.

1. COMPACT CHAINS

Let X denote a smooth n -dimensional manifold. For an integer p we let $\Gamma(X, \Omega^p)$ denote the space of \mathbf{C} -valued differential p -forms on X . By a *compact p -chain* on X we understand a \mathbf{C} -valued linear form T on $\Gamma(X, \Omega^p)$ for which there exists a compact subset K of X such that

$$(1.1) \quad \langle T, \omega \rangle = 0, \quad \omega \in \Gamma(X, \Omega^p), \quad \text{Supp}(\omega) \cap K = \emptyset,$$

where the bracket denotes simple evaluation of a linear form. The compact p -chain T on X gives rise to a $(p-1)$ -chain bT on X given by

$$(1.2) \quad \langle bT, \omega \rangle = \langle T, d\omega \rangle, \quad \omega \in \Gamma(X, \Omega^{p-1}).$$

The operator b makes the compact p -chains on X into a complex, which we denote $D^c(X, \mathbf{C})$. A compact p -chain T is *closed* if $bT = 0$. By a *compact p -cycle* we understand a closed p -chain, while a *compact p -boundary* is a p -cycle of the form bW , where W is a compact $(p+1)$ -chain. We say that p -cycles S and T are *homologous* if $T - S$ is a p -boundary. An explicit relation of the form

$$(1.3) \quad T - S = bW, \quad W \in D_{p+1}^c(X, \mathbf{C}),$$

is called a *de Rham homology* from S to T . Explicit de Rham homologies are often constructed on the basis of Stokes theorem, compare the formulas to the right of the drawings in section 7.

Let us make the important observation that homologous p -chains have the same evolution on any closed p -form. The group of de Rham homology classes is denoted by

$$(1.4) \quad H_p^c(X, \mathbf{C}) = H_p D_p^c(X, \mathbf{C}).$$

The letter c in the homology symbol is borrowed from Haefliger's exposé in [1].

A smooth map $f: X \rightarrow Y$ will induce a chain map from the complex of compact chains on X to the complex of compact chains on Y

$$f_*: D_p^c(X, \mathbf{C}) \rightarrow D_p^c(Y, \mathbf{C}).$$

To see this notice that a given compact subset K on X and a p -form ω on Y supported outside $f(K)$ pulls back to a form $f^*\omega$ supported outside K . We can now define f_*T by the formula

$$(1.5) \quad \langle f_*T, \omega \rangle = \langle T, f^*\omega \rangle, \quad T \in D_p^c(X, \mathbf{C}), \omega \in \Gamma(Y, \Omega^p).$$

These remarks make it clear how to turn de Rham homology into a covariant functor on the smooth category.

Zero cycles. Evaluation of a compact zero cycle Z against the constant function 1 defines the *degree of the zero cycle*

$$(1.6) \quad \text{dg}(Z) = \langle Z, 1 \rangle, \quad Z \in D_0^c(X, \mathbf{C}).$$

A point $x \in X$ defines a zero cycle of degree 1, the *Dirac 0-cycle* δ_x given by

$$(1.7) \quad \langle \delta_x, \varphi \rangle = \varphi(x), \quad \varphi \in \Gamma(X, \Omega^0).$$

A continuously differentiable curve $\gamma: [a, b] \rightarrow X$ with endpoints $x = \gamma(a)$ and $y = \gamma(b)$ defines a de Rham homology from δ_x to δ_y :

$$(1.8) \quad \int_{\gamma} d\varphi = \varphi(y) - \varphi(x), \quad \varphi \in \Gamma(X, \Omega^0).$$

In case X is connected, then all zero cycles of degree zero are homologous to zero as it follows from the result of the next section.

A smooth map $f: X \rightarrow Y$ will preserve the degree of a zero cycle in the sense that

$$(1.9) \quad \text{dg}(f_* T) = \text{dg } T, \quad T \in D_0^c(X, \mathbf{C}),$$

as it follows from (1.5).

The reader is invited to replace \mathbf{C} by \mathbf{R} and change the meaning of the symbol Ω from complex to real differential forms.

2. BIDUALITY

In this section we shall show that de Rham cohomology can be calculated as the linear dual of de Rham homology in the same way singular cohomology can be obtained from singular homology.

(2.1) THEOREM. *Let X denote a smooth manifold. Evaluation of a compact p -chain against a p -form induces an isomorphism*

$$H^p(X, \mathbf{C}) \xrightarrow{\sim} \text{Hom}(H_p^c(X, \mathbf{C}), \mathbf{C})$$

for all integers p .

Proof. The heart of the matter is of sheaf theoretic nature, so we start with a brief review during which the reader is invited to change the meaning of the letter X to denote a general locally compact space and the letter \mathbf{C} to denote an arbitrary field. For notation and details the reader may consult [5] V.1, and the references given there.

To a *soft* \mathbf{C} -sheaf F on X we can associate the sheaf F^\vee whose sections over the open subset U of X are given by

$$(2.2) \quad \Gamma(U, F^\vee) = \text{Hom}(\Gamma_c(U, F), \mathbf{C})$$

Restriction in the sheaf F^\vee from U to a smaller open subset V is the \mathbf{C} -linear dual of "extension by zero"

$$\Gamma_c(V, F) \rightarrow \Gamma_c(U, F), \quad V \subseteq U.$$

The presheaf F^\vee we have described is actually a sheaf and indeed a *soft* sheaf. This allows us to iterate the construction and form $F^{\vee\vee}$. We shall construct a natural *biduality morphism* of \mathbf{C} -sheaves on X

$$(2.3) \quad F \rightarrow F^{\vee\vee}.$$