

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 35 (1989)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: AN ALGEBRAIC PROOF OF VAN DER WAERDEN'S THEOREM
Autor: Bergelson, Vitaly / Furstenberg, Hillel / Hindman, Neil / Katznelson, Yitzhak
Kapitel: 2. Minimal left ideals in right topological semigroups
DOI: <https://doi.org/10.5169/seals-57373>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 28.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

2. MINIMAL LEFT IDEALS IN RIGHT TOPOLOGICAL SEMIGROUPS

We present in this section several well known facts which are not usually seen in early graduate courses.

2.1 LEMMA (Ellis [2]). *Let S be a compact Hausdorff right topological semigroup. Then S has an idempotent, that is there exists $x \in S$ with $x + x = x$.*

Proof. Let $\mathcal{A} = \{A \subseteq S : A \neq \emptyset, A \text{ is compact, and } A + A \subseteq A\}$. Now $\mathcal{A} \neq \emptyset$ since $S \in \mathcal{A}$. Let \mathcal{C} be a chain in \mathcal{A} . Then \mathcal{C} is a collection of closed subsets of S with the finite intersection property so $\bigcap \mathcal{C} \neq \emptyset$ and $\bigcap \mathcal{C}$ is compact. Trivially $(\bigcap \mathcal{C}) + (\bigcap \mathcal{C}) \subseteq \bigcap \mathcal{C}$. Pick by Zorn's Lemma a minimal member A of \mathcal{A} .

Pick $x \in A$ and let $B = A + x$. Now $B = \rho_x[A] (= \{\rho_x(y) : y \in A\})$ so, as the continuous image of a compact set, B is compact (and trivially non-empty). Also $B + B = A + x + A + x \subseteq A + A + A + x \subseteq A + x = B$. Thus $B \in \mathcal{A}$. Since $B = A + x \subseteq A + A \subseteq A$ and A is minimal, $B = A$ so $x \in B = A + x$. That is, there exists $y \in A$ with $x = y + x$.

Let $C = \{y \in A : x = y + x\}$. ρ_x is continuous so $\rho_x^{-1}[\{x\}]$ is closed. Thus C is closed, hence compact. To see that $C + C \subseteq C$, let $y, z \in C$. Then $y + z \in A$ and $(y+z) + x = y + (z+x) = y + x = x$ so $y + z \in C$. Thus $C \in \mathcal{A}$. Since $C \subseteq A$ and A is minimal, $C = A$. Then $x \in C$ and hence $x + x = x$. \square

A non-empty subset I of a semigroup S is a left ideal if $S + I \subseteq I$, a right ideal if $I + S \subseteq I$, and a two-sided ideal if it is both a left ideal and a right ideal. It is a fact (which we will not need) that any right ideal in a compact right topological semigroup contains a minimal right ideal, which need not be closed. (For this and other interesting facts see [1].) We do need a similar fact about left ideals.

2.2 LEMMA. *Let S be a compact Hausdorff right topological semigroup. Any left ideal contains a minimal left ideal and minimal left ideals are closed.*

Proof. Let L be a left ideal of S . Let $\mathcal{A} = \{A \subseteq L : A \text{ is a left ideal and } A \text{ is compact}\}$. Choose $x \in L$. Then $S + x = \rho_x[S]$, the continuous image of a compact space. Also $S + (S+x) = (S+S) + x \subseteq S + x$ so $S + x$ is a left ideal. Since $S + x \subseteq S + L \subseteq L$, we have $\mathcal{A} \neq \emptyset$. One easily sees that the intersection of a chain in \mathcal{A} is again in \mathcal{A} . Choose by Zorn's Lemma a minimal member A of \mathcal{A} .

To see that A is in fact a minimal left ideal, assume we have a left ideal $B \subseteq A$ and pick $x \in B$. Then as above $S + x \in \mathcal{A}$ while $S + x \subseteq S + B \subseteq B \subseteq A$ so $S + x = A$ so $B = A$ \square

2.3 *Definition.* Let S be a semigroup. Then $M(S) = \cup \{L : L \text{ is a minimal left ideal of } S\}$.

It is a fact (which we will not need) that if S is a compact Hausdorff right topological semigroup, then $M(S)$ is a two-sided ideal of S .

2.4 *LEMMA.* Let S be a compact Hausdorff right topological semigroup and let I be a two-sided ideal of S . Then $M(S) \neq \emptyset$ and $M(S) \subseteq I$.

Proof. Since S is a left ideal of S it contains by Lemma 2.2 a minimal left ideal so $M(S) \neq \emptyset$. So see that $M(S) \subseteq I$, let $x \in M(S)$. There is a minimal left ideal L of S with $x \in L$. Also choose some $y \in I$. Then $y + x \in L \cap I$ (since I is a right ideal) so $L \cap I \neq \emptyset$. Thus $L \cap I$ is a left ideal contained in L so that $L \cap I = L$. \square

The proof of the following lemma is an easy exercise and we omit it.

2.5 *LEMMA.* Let S_1 and S_2 be compact right topological semigroups and let $S_1 \times S_2$ have the product topology and coordinatewise operations. Then $S_1 \times S_2$ is a compact right topological semigroup. Given $x \in S_1$ and $y \in S_2$, λ_x and λ_y may or may not be continuous (where $\lambda_x(t) = x + t$). If $\lambda_x : S_1 \rightarrow S_1$ and $\lambda_y : S_2 \rightarrow S_2$ are continuous, then $\lambda_{(x,y)} : S_1 \times S_2 \rightarrow S_1 \times S_2$ is continuous.

3. VAN DER WAERDEN'S THEOREM

We let $l \in \mathbf{N}$ be fixed throughout and show that given any finite partition of \mathbf{N} some one cell contains a length l arithmetic progression.

3.1 *Definition.* (a) Let $Y = (\beta\mathbf{N})^l$ with the product topology and coordinatewise operations.

$$(b) \quad E^* = \{(a, a + d, a + 2d, \dots, a + (l-1)d) : a \in \mathbf{N} \text{ and } d \in \mathbf{N} \cup \{0\}\}.$$

$$(c) \quad I^* = \{(a, a + d, a + 2d, \dots, a + (l-1)d) : a, d \in \mathbf{N}\}.$$

$$(d) \quad E = cl_Y E^*$$

$$(e) \quad I = cl_Y I^*.$$

Note that by Lemmas 1.1 and 2.5, Y is a compact Hausdorff right topological semigroup and whenever $\mathbf{x} = (x_1, x_2, \dots, x_l) \in \mathbf{N}^l$, $\lambda_{\mathbf{x}}$ is continuous.