

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	35 (1989)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	AN ALGEBRAIC PROOF OF VAN DER WAERDEN'S THEOREM
Autor:	Bergelson, Vitaly / Furstenberg, Hillel / Hindman, Neil / Katznelsion, Yitzhak
Kapitel:	3. Van der Waerden's Theorem
DOI:	https://doi.org/10.5169/seals-57373

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

To see that A is in fact a minimal left ideal, assume we have a left ideal $B \subseteq A$ and pick $x \in B$. Then as above $S + x \in \mathcal{A}$ while $S + x \subseteq S + B \subseteq B \subseteq A$ so $S + x = A$ so $B = A$ \square

2.3 *Definition.* Let S be a semigroup. Then $M(S) = \cup \{L : L \text{ is a minimal left ideal of } S\}$.

It is a fact (which we will not need) that if S is a compact Hausdorff right topological semigroup, then $M(S)$ is a two-sided ideal of S .

2.4 *LEMMA.* Let S be a compact Hausdorff right topological semigroup and let I be a two-sided ideal of S . Then $M(S) \neq \emptyset$ and $M(S) \subseteq I$.

Proof. Since S is a left ideal of S it contains by Lemma 2.2 a minimal left ideal so $M(S) \neq \emptyset$. So see that $M(S) \subseteq I$, let $x \in M(S)$. There is a minimal left ideal L of S with $x \in L$. Also choose some $y \in I$. Then $y + x \in L \cap I$ (since I is a right ideal) so $L \cap I \neq \emptyset$. Thus $L \cap I$ is a left ideal contained in L so that $L \cap I = L$. \square

The proof of the following lemma is an easy exercise and we omit it.

2.5 *LEMMA.* Let S_1 and S_2 be compact right topological semigroups and let $S_1 \times S_2$ have the product topology and coordinatewise operations. Then $S_1 \times S_2$ is a compact right topological semigroup. Given $x \in S_1$ and $y \in S_2$, λ_x and λ_y may or may not be continuous (where $\lambda_x(t) = x + t$). If $\lambda_x : S_1 \rightarrow S_1$ and $\lambda_y : S_2 \rightarrow S_2$ are continuous, then $\lambda_{(x, y)} : S_1 \times S_2 \rightarrow S_1 \times S_2$ is continuous.

3. VAN DER WAERDEN'S THEOREM

We let $l \in \mathbf{N}$ be fixed throughout and show that given any finite partition of \mathbf{N} some one cell contains a length l arithmetic progression.

3.1 *Definition.* (a) Let $Y = (\beta\mathbf{N})^l$ with the product topology and coordinatewise operations.

- (b) $E^* = \{(a, a + d, a + 2d, \dots, a + (l-1)d) : a \in \mathbf{N} \text{ and } d \in \mathbf{N} \cup \{0\}\}$.
- (c) $I^* = \{(a, a + d, a + 2d, \dots, a + (l-1)d) : a, d \in \mathbf{N}\}$.
- (d) $E = cl_Y E^*$
- (e) $I = cl_Y I^*$.

Note that by Lemmas 1.1 and 2.5, Y is a compact Hausdorff right topological semigroup and whenever $\mathbf{x} = (x_1, x_2, \dots, x_l) \in \mathbf{N}^l$, $\lambda_{\mathbf{x}}$ is continuous.

3.2 LEMMA. E is a compact Hausdorff right topological semigroup and I is a two sided ideal of E .

Proof. Compactness is immediate and the Hausdorff property and right continuity are inherited from Y . We let $\mathbf{p} = (p_1, p_2, \dots, p_l)$ and $\mathbf{q} = (q_1, q_2, \dots, q_l)$ be members of E and show that $\mathbf{p} + \mathbf{q} \in E$. We show further that if either \mathbf{p} or \mathbf{q} is in I , then $\mathbf{p} + \mathbf{q} \in I$.

To see that $\mathbf{p} + \mathbf{q} \in E$, let U be a neighborhood of $\mathbf{p} + \mathbf{q}$. By the continuity of $\rho_{\mathbf{q}}$, pick a neighborhood V of \mathbf{p} with $V + \mathbf{q} = \rho_{\mathbf{q}}[V] \subseteq U$. Since $\mathbf{p} \in cl E^*$ we may pick $a \in \mathbf{N}$ and $d \in \mathbf{N} \cup \{0\}$ with

$$(a, a+d, a+2d, \dots, a+(l-1)d) \in V.$$

If $\mathbf{p} \in I$ we may presume $d \neq 0$. Let $\mathbf{x} = (a, a+d, a+2d, \dots, a+(l-1)d)$. Then $\mathbf{x} \in V$ so $\mathbf{x} + \mathbf{q} \in U$. By the continuity of $\lambda_{\mathbf{x}}$, pick a neighborhood W of \mathbf{q} with $\mathbf{x} + W = \lambda_{\mathbf{x}}[W] \subseteq U$. Since $\mathbf{q} \in cl E^*$, pick $b \in \mathbf{N}$ and $c \in \mathbf{N} \cup \{0\}$ (with $c \neq 0$ if $\mathbf{q} \in I$) such that $(b, b+c, b+2c, \dots, b+(l-1)c) \in W$. Let $\mathbf{y} = (b, b+c, b+2c, \dots, b+(l-1)c)$. Then $\mathbf{x} + \mathbf{y} \in U \cap E^*$. If either $d \neq 0$ or $c \neq 0$, then $c + d \neq 0$ so $\mathbf{x} + \mathbf{y} \in U \cap I^*$. \square

3.3 THEOREM. Let $p \in M(\beta\mathbf{N})$ and let $\mathbf{p} = (p, p, \dots, p)$. Then $\mathbf{p} \in I$.

Proof. We first show that $\mathbf{p} \in E$. Let $U_1 \times U_2 \times \dots \times U_l$ be a basic neighborhood of \mathbf{p} . Then $U_1 \cap U_2 \cap \dots \cap U_l$ is a neighborhood of p in $\beta\mathbf{N}$. Since \mathbf{N} is dense, pick $a \in \mathbf{N} \cap (U_1 \cap U_2 \cap \dots \cap U_l)$. Then $(a, a, \dots, a) \in E^* \cap (U_1 \times U_2 \times \dots \times U_l)$. Thus $\mathbf{p} \in cl E^* = E$.

Since $p \in M(\beta\mathbf{N})$, there is a minimal left ideal L of $\beta\mathbf{N}$ with $p \in L$. Since $E + \mathbf{p}$ is a left ideal of E , pick by Lemma 2.2 a minimal left ideal L^* of E with $L^* \subset E + \mathbf{p}$. Since L^* is closed, hence compact, pick by Lemma 2.1 an idempotent $\mathbf{q} = (q_1, q_2, \dots, q_l)$ in L^* . Now $\mathbf{q} \in L^* \subseteq E + \mathbf{p}$ so pick some $\mathbf{s} = (s_1, s_2, \dots, s_l)$ in E with $\mathbf{q} = \mathbf{s} + \mathbf{p}$.

We show that $\mathbf{p} + \mathbf{q} = \mathbf{p}$. To this end let $i \in \{1, 2, \dots, l\}$. Now $q_i = s_i + p \in L$ so $\beta\mathbf{N} + q_i \subseteq \beta\mathbf{N} + L \subseteq L$. Thus $\beta\mathbf{N} + q_i$ is a left ideal contained in the minimal left ideal L so that $\beta\mathbf{N} + q_i = L$. Thus since $p \in L$ there exists $t_i \in \beta\mathbf{N}$ with $t_i + q_i = p$. But then $p + q_i = t_i + q_i + q_i = t_i + q_i = p$ as required.

Since $\mathbf{p} \in E$ and $\mathbf{q} \in L^*$, a left ideal of E , we have $\mathbf{p} = \mathbf{p} + \mathbf{q} \in L^*$ so that $\mathbf{p} \in M(E)$. Thus by Lemma 2.4, $\mathbf{p} \in I$.

3.4 COROLLARY (van der Waerden). Let $m \in \mathbf{N}$ and let $\{A_1, A_2, \dots, A_m\}$ be a partition of \mathbf{N} . There exist $i \in \{1, 2, \dots, m\}$ and $a, d \in \mathbf{N}$ with $\{a, a+d, a+2d, \dots, a+(l-1)d\} \subseteq A_i$.

Proof. By Lemma 2.4 $M(\beta\mathbf{N}) \neq \emptyset$ so pick $p \in M(\beta\mathbf{N})$ and let $\mathbf{p} = (p, p, \dots, p)$. By Lemma 1.2 pick $i \in \{1, 2, \dots, m\}$ such that $cl A_i$ is a neighborhood of p and let $U = cl A_i$. Then $U \times U \times \dots \times U$ is a neighborhood of \mathbf{p} while, by Theorem 2.3, $\mathbf{p} \in I = cl I^*$. Pick $a, d \in \mathbf{N}$ with $(a, a+d, a+2d, \dots, a+(l-1)d) \in U \times U \times \dots \times U$. Then

$$\{a, a+d, a+2d, \dots, a+(l-1)d\} \subseteq U \cap \mathbf{N} = (cl A_i) \cap N = A_i. \quad \square$$

We remark that if one starts with the free semigroup on l letters in place of \mathbf{N} , essentially the same proof yields the Hales-Jewett Theorem. See [3] for the details.

REFERENCES

- [1] BERGLUND, J., H. JUNGHENN and P. MILNES. *Compact right topological semigroups and generalizations of almost periodicity*. Lecture Notes in Math. 663, Springer-Verlag, Berlin (1978).
- [2] ELLIS, R. *Lectures on Topological Dynamics*. Benjamin, New York, 1969.
- [3] FURSTENBERG, H. and Y. KATZNELSON. Idempotents and coloring theorems. To appear.
- [4] FURSTENBERG, H. and B. WEISS. Topological dynamics and combinatorial number theory. *J. d'Analyse Math.* 34 (1978), 61-85.
- [5] GRAHAM, R. and B. ROTHSCHILD. A short proof of van der Waerden's Theorem. *Proc. Amer. Math. Soc.* 42 (1974), 385-386.
- [6] HINDMAN, N. Ultrafilters and Ramsey Theory — an update. *Proceedings of "Set Theory and its Applications – Conference at York"*. To appear.
- [7] SHELAH, S. Primitive recursive bounds for van der Waerden numbers. *J. Amer. Math. Soc.* 1 (1989), 683-697.
- [8] TAYLOR, A. A note on van der Waerden's Theorem. *J. Comb. Theory (Series A)* 33 (1982), 215-219.
- [9] VAN DER WAERDEN, B. Beweis einer Baudetschen Vermutung. *Nieuw Arch. Wisk* (3) 19 (1927), 212-216.

(Reçu le 14 juin 1989)

Vitaly Bergelson

Department of Mathematics
Ohio State University
Columbus, OH 43210
(USA)

Hillel Furstenberg

Department of Mathematics
Hebrew University
Jerusalem (Israel)

Neil Hindman

Department of Mathematics
Howard University
Washington, DC 20059
(USA)

Yitzhak Katznelson

Department of Mathematics
Stanford University
Stanford, CA 94305
(USA)