

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 35 (1989)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: CAUCHY RESIDUES AND DE RHAM HOMOLOGY
Autor: Iversen, Birger
Kapitel: 2. Biduality
DOI: <https://doi.org/10.5169/seals-57358>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A smooth map $f: X \rightarrow Y$ will preserve the degree of a zero cycle in the sense that

$$(1.9) \quad \text{dg}(f_* T) = \text{dg } T, \quad T \in D_0^c(X, \mathbf{C}),$$

as it follows from (1.5).

The reader is invited to replace \mathbf{C} by \mathbf{R} and change the meaning of the symbol Ω from complex to real differential forms.

2. BIDUALITY

In this section we shall show that de Rham cohomology can be calculated as the linear dual of de Rham homology in the same way singular cohomology can be obtained from singular homology.

(2.1) THEOREM. *Let X denote a smooth manifold. Evaluation of a compact p -chain against a p -form induces an isomorphism*

$$H^p(X, \mathbf{C}) \xrightarrow{\sim} \text{Hom}(H_p^c(X, \mathbf{C}), \mathbf{C})$$

for all integers p .

Proof. The heart of the matter is of sheaf theoretic nature, so we start with a brief review during which the reader is invited to change the meaning of the letter X to denote a general locally compact space and the letter \mathbf{C} to denote an arbitrary field. For notation and details the reader may consult [5] V.1, and the references given there.

To a soft \mathbf{C} -sheaf F on X we can associate the sheaf F^\vee whose sections over the open subset U of X are given by

$$(2.2) \quad \Gamma(U, F^\vee) = \text{Hom}(\Gamma_c(U, F), \mathbf{C})$$

Restriction in the sheaf F^\vee from U to a smaller open subset V is the \mathbf{C} -linear dual of "extension by zero"

$$\Gamma_c(V, F) \rightarrow \Gamma_c(U, F), \quad V \subseteq U.$$

The presheaf F^\vee we have described is actually a sheaf and indeed a soft sheaf. This allows us to iterate the construction and form $F^{\vee\vee}$. We shall construct a natural *biduality morphism* of \mathbf{C} -sheaves on X

$$(2.3) \quad F \rightarrow F^{\vee\vee}.$$

To this end consider the tautological evaluation

$$\Gamma(U, F^\vee) \times \Gamma_c(U, F) \rightarrow \mathbb{C}.$$

This can be modified to yield a pairing

$$\text{ev}: \Gamma_c(U, F^\vee) \times \Gamma(U, F) \rightarrow \mathbb{C}$$

namely, for $T \in \Gamma_c(U, F^\vee)$ and $\omega \in \Gamma(U, F)$ choose $v \in \Gamma_c(U, F)$, such that ω and v has the same restriction to $\text{Supp}(T)$, and put $\text{ev}(T, \omega) = T(v)$. The evaluation map may be interpreted as a transformation

$$(2.4) \quad a_U: \Gamma_c(U, F^\vee) \rightarrow \text{Hom}(\Gamma(U, F), \mathbb{C}).$$

An open subset V of U gives rise to a commutative diagram

$$(2.5) \quad \begin{array}{ccc} \Gamma_c(V, F^\vee) & \xrightarrow{a_V} & \text{Hom}(\Gamma(V, F), \mathbb{C}) \\ \downarrow & & \downarrow \\ \Gamma_c(U, F^\vee) & \xrightarrow{a_U} & \text{Hom}(\Gamma(U, F), \mathbb{C}) \end{array}$$

where the first vertical arrow is “extension by zero” in the soft sheaf F^\vee and the second vertical arrow is the linear dual of restriction in the sheaf F . Let us return to the open subset U and consider the composite

$$\Gamma(U, F) \rightarrow \text{Hom}(\text{Hom}(\Gamma(U, F), \mathbb{C}), \mathbb{C}) \xrightarrow{a_U^*} \Gamma(U, F^{\vee\vee})$$

where the first arrow is the biduality map from linear algebra. By variation of U we obtain the biduality morphism $b: F \rightarrow F^{\vee\vee}$ announced in (2.3).

Let us now return to the situation at hand and consider the biduality morphism for the de Rham complex.

$$(2.6) \quad b: \Omega^\bullet \rightarrow \Omega^{\bullet\vee\vee}$$

which we shall prove to be a quasi-isomorphism. The question is local, so it suffices to check the case $X = \mathbf{R}^n$, which can be done by the Poincaré Lemma with and without compact support. Both complexes are made of soft sheaves, so we lean on the fact, implicit in the definition of a manifold, that X is countable at infinity to conclude that b induces isomorphisms, compare [5] IV.2.2,

$$H^p(X, \mathbb{C}) \xrightarrow{\sim} H^p\Gamma(X, \Omega^{\bullet\vee\vee}), \quad p \in \mathbb{N}.$$

In order to identify the right hand side notice first that

$$\Gamma(X, \Omega^{\bullet\vee\vee}) = \text{Hom}(\Gamma_c(X, \Omega^\bullet), \mathbb{C})$$

and second, that the map a_X introduced in (2.4) induces an isomorphism

$$(2.7) \quad a: \Gamma_c(X, \Omega^{\bullet \vee}) \xrightarrow{\sim} D^c_c(X, \mathbb{C}).$$

Collect this together to conclude the proof. Q.E.D.

The de Rham homology as defined here agrees with the original theory based on currents [6]: the inclusion of the complex of currents in $\Omega^{\bullet \vee}$ is a quasi-isomorphism as can be seen by the method used in the last third of the proof above.

As a consequence of the isomorphism (2.7) we can of course redefine de Rham homology as

$$(2.8) \quad H^c_p(X, \mathbb{C}) = H_p \Gamma_c(X, \Omega^{\bullet \vee}).$$

If the letter c is dropped we obtain Borel Moore homology, compare [5] IX and the references given there.

The biduality theorem 2.1 is certainly related to that of Verdier [7], [1]. In fact most of the material presented here may be extended to a context of similar generality. I hope to return to this point in the near future.

3. SMOOTH SINGULAR HOMOLOGY

Let us consider an n -dimensional smooth manifold X . Integration over smooth singular simplexes defines a map

$$(3.1) \quad S^{\infty}_*(X, \mathbb{C}) \rightarrow D^c_c(X, \mathbb{C})$$

from the complex of smooth singular simplexes to the complex of compact chains on X .

(3.2) THEOREM. *Integration induces an isomorphism*

$$H^{\infty}_*(X, \mathbb{C}) \xrightarrow{\sim} H^c_c(X, \mathbb{C})$$

from smooth singular homology to de Rham homology.

Proof. Let us first discuss *Mayer-Vietoris* sequences in de Rham homology. For open subsets U and V of X a Mayer-Vietoris sequence arises from the following exact sequence of complexes

$$(3.3) \quad 0 \rightarrow \Gamma_c(U \cap V, \Omega^{\bullet \vee}) \xrightarrow{+} \Gamma_c(U, \Omega^{\bullet \vee}) \oplus \Gamma_c(V, \Omega^{\bullet \vee}) \xrightarrow{+} \Gamma_c(U \cup V, \Omega^{\bullet \vee}) \rightarrow 0.$$