

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	35 (1989)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
 Artikel:	ZETA FUNCTIONS AND GENUS OF QUADRATIC FORMS
Autor:	Bayer, Pilar / Nart, Enric
Kapitel:	§2. Local representation masses and Z_p -equivalence OF FORMS
DOI:	https://doi.org/10.5169/seals-57377

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

possible anymore, as the following example shows: By Proposition 1.1 the diagonal forms $f = \langle 1, 2, 2, 2, 4 \rangle$, $g = \langle 1, 1, 2, 4, 4 \rangle$ have the same Gauss sums $\theta(\ , f, 2^t) = \theta(\ , g, 2^t)$ for all $t \geq 1$, however they are obviously not \mathbf{Z}_2 -equivalent.

The theory of Minkowski reproduced in this section was extended by O'Meara to integral quadratic forms over local fields.

§ 2. LOCAL REPRESENTATION MASSES AND \mathbf{Z}_p -EQUIVALENCE OF FORMS

We identify \mathbf{Q}_p with its topological dual by defining $\langle n, m \rangle = \chi_p(nm)$, where χ_p is Tate's character:

$$\chi_p(a) = \exp(2\pi i \sum_{s < 0} a_s p^s),$$

if $a = \sum_{s \geq s_0} a_s p^s$. Let dn be the Haar measure of \mathbf{Q}_p normalized by $dn(\mathbf{Z}_p) = 1$.

As is well-known, dn is selfdual. Let dx be the Haar measure of \mathbf{Q}_p naturally induced by dn .

Let f be a non-singular integral p -adic quadratic form in $k \geq 1$ variables. We shall deal in this section with the representation mass function given by (0.1) for $\phi = 1_{(\mathbf{Z}_p)^k}$. That is, we define for all $n_o \in \mathbf{Q}_p$:

$$r(n_o, f, \mathbf{Z}_p) = \lim_{U \rightarrow \{n_o\}} (dx(f^{-1}(U) \cap \mathbf{Z}_p^k) / dnU),$$

whenever this limit exists. Clearly r has support contained in \mathbf{Z}_p . We can also consider the Gauss-Weil transform of $1_{(\mathbf{Z}_p)^k}$ by f given by

$$\theta(m, f, \mathbf{Q}_p) = \int_{\mathbf{Z}_p^k} \langle f(x), m \rangle dx.$$

The relationship between these representation masses and the ones introduced in the preceding section is given in the following

LEMMA 2.1. i) *Let $n \in \mathbf{Z}_p$, $n \neq 0$, and $t > v_p(4n)$. Then*

$$r(n, f, \mathbf{Z}_p) = \lim_{s \rightarrow \infty} p^{(1-k)s} r(n, f, p^s) = p^{(1-k)t} r(n, f, p^t).$$

ii) *Let $m \in \mathbf{Z}_p$ and $u \in \mathbf{Z}_p$, $t \geq 1$ be chosen arbitrarily satisfying $m = up^{-t}$. Then*

$$\theta(m, f, \mathbf{Q}_p) = p^{-kt} \theta(u, f, p^t).$$

Proof. i) Let $U_t = n + p^t \mathbf{Z}_p$. We have $dn(U_t) = p^{-t}$ and

$$dx(f^{-1}(U_t) \cap \mathbf{Z}_p^k) = \sum_{a \in (\mathbf{Z}/p^t \mathbf{Z})^k} dx(f^{-1}(U_t) \cap (a + p^t \mathbf{Z}_p^k)) = p^{-kt} r(n, f, p^t),$$

since $f^{-1}(U_t) \cap (a + p^t \mathbf{Z}_p^k)$ is equal to $a + p^t \mathbf{Z}_p^k$ or vacuous, according to $f(a) \equiv n \pmod{p^t}$ or not. This proves the first equality in i).

We want now to show that $p^{(1-k)s} r_{p^s}(n) = p^{(1-k)(s-1)} r_{p^{s-1}}(n)$, for all $s > t$. We know that

$$r(n, f, p^s) = p^{-s} \sum_{u=1}^{p^s} \theta(u, f, p^s) \exp(-2\pi i u n p^{-s}).$$

Let us denote by A and B the sum of the terms satisfying $p \mid u$ and $p \nmid u$, respectively. Clearly $A = p^{k-1} r(n, f, p^{s-1})$; hence, we are reduced to proving $B = 0$. Taking into account the explicit computations of Gauss sums (Proposition 1.1), we can express the sum B as

$$B = \begin{cases} C \sum_{u \in (\mathbf{Z}/p^s \mathbf{Z})^*} \left(\frac{u}{p}\right)^a \exp(-2\pi i u n p^{-s}) & \text{if } p > 2 \\ D \sum_{u \in (\mathbf{Z}/2^s \mathbf{Z})^*} \left(\frac{2}{u}\right)^b \exp\left(\frac{2\pi i u}{8}\right)^c \exp(-2\pi i u n 2^{-s}) & \text{if } p = 2, \end{cases}$$

where C, D, a, b, c depend on f and s , but are independent of u . Now, $\exp(-2\pi i u n p^{-s})$ is a primitive p^l -th root of 1 with $l > 1$ if $p > 2$, and $l > 3$ if $p = 2$. One can check that, for any function φ defined on $(\mathbf{Z}/p^m \mathbf{Z})^*$, $m \geq 1$ and ξ any primitive p^l -th root of 1, $l > m$, one has

$$\sum_{u \in (\mathbf{Z}/p^l \mathbf{Z})^*} \varphi(u) \xi^u = 0.$$

In particular, B must be zero.

In order to prove ii) we need only to observe that

$$\begin{aligned} \theta(m, f, \mathbf{Q}_p) &= \int_{\mathbf{Z}_p^k} \exp(2\pi i f(x) u p^{-t}) dx \\ &= \sum_{a \in (\mathbf{Z}/p^t \mathbf{Z})^k} \exp(2\pi i f(a) u p^{-t}) \int_{a + p^t \mathbf{Z}_p^k} dx = p^{-kt} \theta(u, f, p^t). \quad \square \end{aligned}$$

Remark. After Siegel [13], it was very well known that for $n \neq 0$ the values $p^{(1-k)t} r(n, f, p^t)$ become constant for $t > 2v_p(4n)$. Lemma 2.1 shows that the minimum value of t with this property can be taken equal to half of the one found by Siegel.

By Lemma 2.1, $r(\ , f, \mathbf{Z}_p)$ is locally constant, hence continuous on \mathbf{Q}_p^* , and $r(n, f, \mathbf{Z}_p) = 0$ if and only if n is not represented by f in \mathbf{Z}_p . The fundamental fact is that r is integrable on \mathbf{Z}_p and θ is its Fourier transform. This is well-known [4]. For the sake of completeness we give a short proof of this result using only the background introduced up to now.

PROPOSITION 2.2. $r \in L^1(\mathbf{Z}_p)$ and

$$\theta(m, f, \mathbf{Q}_p) = \int_{\mathbf{Z}_p} r(n, f, \mathbf{Z}_p) \langle n, m \rangle dn.$$

Proof. We assume $p > 2$. For $p = 2$ the proof works in the same way with minor modifications left to the reader. Let $m = up^{-s}$, $u \in \mathbf{Z}_p$, $s \geq 0$. For all $t > s$, $\mathbf{Z}_p \setminus p^t \mathbf{Z}_p$ is compact, hence $r(n)$, being continuous, is integrable and we have by Lemma 2.1:

$$\begin{aligned} \int_{\mathbf{Z}_p \setminus p^t \mathbf{Z}_p} r(n, f, \mathbf{Z}_p) \langle n, m \rangle dn &= \sum_{\substack{a \in \mathbf{Z}/p^t \mathbf{Z} \\ a \neq 0}} \int_{a + p^t \mathbf{Z}_p} r(n, f, \mathbf{Z}_p) \langle n, m \rangle dn \\ &= \sum_{\substack{a \in \mathbf{Z}/p^t \mathbf{Z} \\ a \neq 0}} p^{-kt} r(a, f, p^t) \exp(2\pi i a u p^{-s}) = p^{-kt} (\theta(p^{t-s} u, f, p^t) - r(0, f, p^t)) \\ &= \theta(m, f, \mathbf{Q}_p) - p^{-kt} r(0, f, p^t). \end{aligned}$$

Both assertions of the proposition are consequences of Lebesgue's dominated convergence theorem if $p^{-kt} r(0, f, p^t)$ tends to zero as t tends to infinity. This is checked immediately for $k = 1$. For $k > 1$ it can be easily deduced from (1.1) and the explicit computation of Gauss sums in the preceding section. \square

We are ready to prove a crucial fact for the rest of the paper:

THEOREM 2.3. *Let f, g be two non-singular integral p -adic quadratic forms in k variables. If $p = 2$, assume that they are of the same type. The following conditions are equivalent:*

- i) $f \sim g$ over \mathbf{Z}_p ,
- ii) $r(\ , f, \mathbf{Z}_p) = r(\ , g, \mathbf{Z}_p)$,
- iii) $\theta(\ , f, \mathbf{Q}_p) = \theta(\ , g, \mathbf{Q}_p)$.

Proof. If $f \sim g$ over \mathbf{Z}_p , then $f \sim g$ over $\mathbf{Z}/p^t \mathbf{Z}$ and $r(\ , f, p^t) = r(\ , g, p^t)$ for all $t \geq 1$. By Lemma 2.1 this implies ii). By Proposition 2.2, ii) implies iii). Again by Lemma 2.1, iii) implies that $\theta(\ , f, p^t) = \theta(\ , g, p^t)$ for all $t \geq 1$, therefore condition i) follows now from Theorem 1.2. \square

Let K be a local field and f a non-singular quadratic form in k variables defined over K . If ϕ is a Schwartz-Bruhat function on K^k , the representation mass function $r_\phi(\ , f, K)$ defined as in (0.1) coincides with another classical representation mass function introduced by Weil. This is Weil's procedure (see [4] for the details): for $n \neq 0$, the $(k-1)$ -differential forms

$$\omega_i(x) = (-1)^{i-1} (D_i f)^{-1} dx_1 \wedge \dots \wedge d\hat{x}_i \wedge \dots \wedge dx_k,$$

induce a gauge form ω_n on the affine variety $f^{-1}(n)$. Since we are in a local field, ω_n induces a positive measure $|\omega_n|$ on $f^{-1}(n)$ such that for every continuous function φ on K^k with compact support not containing zero we have

$$(2.1) \quad \int_{K^k} \varphi(x) dx = \int_K \left(\int_{f^{-1}(n)} \varphi |\omega_n| \right) dn.$$

The representation mass of $n \in K^*$ by f with respect to ϕ is then defined as

$$F_\phi(n) = \int_{f^{-1}(n)} \phi |\omega_n|.$$

This function is continuous and after (2.1) it is easy to prove that F_ϕ is integrable and its Fourier transform coincides with the Gauss-Weil transform:

$$\int_{K^k} \phi(x) \langle f(x), m \rangle dx = \int_K F_\phi(n) \langle n, m \rangle dn.$$

Let now $n_o \in K^*$ and let U be any open neighbourhood of n_o . From (2.1) it is also easy to justify that:

$$\int_{f^{-1}(U)} \phi(x) dx = \int_U F_\phi(n) dn.$$

Since F_ϕ is continuous and K is locally compact, we have also:

$$F_\phi(n_o) = \lim_{U \rightarrow \{n_o\}} \left(\int_U F_\phi(n) dn / \int_U dn \right) = r_\phi(n_o),$$

thus $F_\phi = r_\phi$ on K^* .