

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 35 (1989)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE TOWER OF HANOI
Autor: Hinz, Andreas M.
Kapitel: 3. Open problems
DOI: <https://doi.org/10.5169/seals-57378>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 29.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

In all the other cases, the moves of disc $n + 1$ are determined by Lemma 7, the moves of the other n discs are governed by Lemmas 5 and 6, and their uniqueness follows by induction hypothesis, keeping in mind that the paths of Lemma 5 are actually paths from $\bar{\sigma}^d$ to $\hat{j}^{n-h_\sigma(d)}$. \square

Using the methods of this chapter, one finds the shortest path from σ to $\hat{0}$ in Figure 1 with length 102.

3. OPEN PROBLEMS

Much of the discussion of the **TH** in computer science literature has been a controversy between recursion and iteration. It has turned out here that problems involving just regular states, can be solved by iteration very elegantly (Chapter 1). On the other hand, as soon as irregular states are considered, only recursive solutions are available (Chapter 2). While for $\mathfrak{P}3$ the solution is essentially unique and the recursion will work efficiently, the situation for $\mathfrak{P}4$ is less straightforward. Although the number of cases to be considered can be further limited by methods as in Section 2.3 (e.g. the shortest path (of length 108) from σ to r in Figure 1 is unique), and one can show that no three moves of the largest disc $n + 1$ occur if $r(n+1) = t(n+1)$ and $h(n+1) > 1$, it is not clear whether there are shortest path problems with even three different solutions. Also it seems that the minimal length in $\mathfrak{P}3$ and 4 can only be determined recursively.

The only existing solution to the **TH** with more than three pegs is also recursive, and the preceding chapters should have demonstrated that things are not as easy as many authors might hope (see the remarks in the Introduction). To move the largest disc $n + 1$ in the solution of $\mathfrak{P}0$ with four pegs, the n other discs have to be transferred to two different pegs; after the last move of disc $n + 1$, discs 1 to n have to be sent from some two pegs to the top of disc $n + 1$. Again it follows by symmetry that disc $n + 1$ will only be moved once in a shortest path. But this time, this does not reduce the problem for $n + 1$ discs to a similar one with only n discs, but to the different setting of how to transfer n discs from a perfect state to two different pegs in the shortest possible way. Here is where the hitherto unjustified assumption made in literature enters, namely that this will be achieved by dividing the perfect state in a suitable way into two parts, then first solving $\mathfrak{P}0$ for the smaller discs using four pegs, leaving them untouched thereafter and solving the old problem for the larger discs using three pegs only.

The validity of this hypothesis is the most interesting open problem about the TH. It might be found by checking a suitable guess about the minimal length for $\mathfrak{B}2$ with four pegs against the recursive solution which can easily be constructed using the fact, proved as Lemma 1, that the largest disc will not move more than three times.

In contrast to this recursive solution, the use of the hypothesis leads to a very elegant iterative solution to $\mathfrak{B}0$ with four (or more) pegs (see Hinz [26]), resembling algorithm i in 1.2.1, with the astonishing result that the transfer of 64 discs can be carried out in less than 6 hours (compare the time needed with three pegs, indicated in the Introduction!).

To conclude, it can be said that the invention of Edouard Lucas, besides its appeal as a puzzle for human beings as well as for computer performance, has been endowed with enough structure to be treated mathematically (the problem $\mathfrak{B}5 : = \text{irregular} \rightarrow \text{irregular without the "divine rule"}$ (0) seems to have almost no mathematical structure), but not with so much to be trivial and incapable of meaningful generalizations. As long as there are still open problems, a mathematical subject is not dead. The brahmins are alive and as long as they are still moving golden discs, the world will, according to legend, not fall to dust. Let us hope so!

REFERENCES

- [1] AFRIAT, S. N. *The Ring of Linked Rings*. Duckworth (London), 1982.
- [2] ALLARDICE, R. E. and A. Y. FRASER. La Tour d'Hanoi. *Proc. Edinburgh Math. Soc.* 2 (1883-84), 50-53.
- [3] BALL, W. W. R. *Mathematical recreations and problems of past and present times*. Macmillan (London), 1892 (2nd ed.).
- [4] BENDISCH, J. Generalized Sequencing Problem "Towers of Hanoi". *Z. Oper. Res.* 29 (1985), 31-45.
- [5] BERMAN, G. and K. D. FRYER. *Introduction to combinatorics*. Academic Press (New York), 1972.
- [6] BROUSSEAU, A. Tower of Hanoi with more pegs. *J. Recreational Math.* 8 (1975-76), 169-178.
- [7] CHARNIAK, E. and D. MCDERMOTT. *Introduction to Artificial Intelligence*. Addison-Wesley (Reading (Mass.)), 1986.
- [8] CLAUS, N. (= E. LUCAS). *La Tour d'Hanoi, Véritable casse-tête annamite*. P. Bousrez (Tours), 1883.
- [9] ———. La Tour d'Hanoi, Jeu de calcul. *Science et Nature, Vol. I, N° 8* (1884), 127-128.
- [10] CROWE, D. W. The n -dimensional cube and the Tower of Hanoi. *Amer. Math. Monthly* 63 (1956), 29-30.
- [11] CULL, P. and E. F. ECKLUND Jr. On the Towers of Hanoi and Generalized Towers of Hanoi Problems. *Congress. Numer.* 35 (1982), 229-238.