

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 35 (1989)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: CAUCHY RESIDUES AND DE RHAM HOMOLOGY
Autor: Iversen, Birger
Kapitel: 5. Stokes formula
DOI: <https://doi.org/10.5169/seals-57358>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

Download PDF: 21.05.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

and similarly with X replaced by U and X_α replaced by $U \cap X_\alpha$. Using this and the exact sequence 4.2 we get that

$$\lim_{\rightarrow} D^c(X_\alpha, U \cap X_\alpha; \mathbf{C}) = D^c(X, U; \mathbf{C})$$

from which the result follows by passing to homology.

Q.E.D.

Let us also notice that in case X is the disjoint union of a family (X_α) of open subsets we have that

$$(4.9) \quad \bigoplus_{\alpha} H^c(X_\alpha, U \cap X_\alpha; \mathbf{C}) \xrightarrow{\sim} H^c(X, U; \mathbf{C}).$$

5. STOKES FORMULA

Let us consider the open subset U of the n -dimensional smooth manifold X and the resulting exact sequences

$$(5.1) \quad \begin{aligned} &\rightarrow H_p^c(X, \mathbf{C}) \rightarrow H_p^c(X, U; \mathbf{C}) \xrightarrow{b} H_{p-1}^c(U, \mathbf{C}) \xrightarrow{j_*} H_{p-1}^c(X, \mathbf{C}) \rightarrow \\ &\leftarrow H^p(X, \mathbf{C}) \leftarrow H^p(X, U; \mathbf{C}) \xleftarrow{\partial} H^{p-1}(U, \mathbf{C}) \xleftarrow{j^*} H^{p-1}(X, \mathbf{C}) \leftarrow \end{aligned}$$

where the first is discussed in the previous section and the second is the sheaf cohomology sequence. The relative term in the second sequence is often written

$$(5.2) \quad H_Z^p(X, \mathbf{C}) = H^p(X, U; \mathbf{C}), \quad Z = X - U.$$

We can now extend the biduality theorem (2.1).

(5.3) THEOREM. *The cohomology sequence above is dual to the homology sequence. In particular we have a Stoke's formula*

$$\langle b\alpha, \omega \rangle = \langle \alpha, \partial\omega \rangle$$

for $\alpha \in H_p^c(X, U; \mathbf{C})$ and $\omega \in H^{p-1}(U, \mathbf{C})$.

Proof. The first sequence arises from the following short exact sequence of complexes, compare (4.2) and (4.7),

$$0 \rightarrow \Gamma_c(U, \Omega^{\cdot \vee \vee}) \xrightarrow{j_*} \Gamma_c(X, \Omega^{\cdot \vee \vee}) \rightarrow \Gamma_c(Z, \Omega^{\cdot \vee \vee}) \rightarrow 0.$$

In order to calculate the second sequence we depart from the flabby resolution $\Omega^{\cdot \vee \vee}$ of \mathbf{R} established in the proof of the biduality theorem (2.1).

The basic philosophy being that flabby sheaves are acyclic for local cohomology, [5] II. 9.3. Thus we can calculate the cohomology sequence (5.1) from the short exact sequence

$$0 \leftarrow \Gamma(U, \Omega^{\bullet \vee \vee}) \xleftarrow{j^*} \Gamma(X, \Omega^{\bullet \vee \vee}) \leftarrow \Gamma_Z(X, \Omega^{\bullet \vee \vee}) \leftarrow 0.$$

According to formula (2.4) we may identify the arrow marked j^* with the linear dual of the arrow marked j_* . Simple evaluation according to (2.4) will be written

$$\langle T, l \rangle, \quad T \in \Gamma_c(X,)^{\bullet \vee}, \quad l \in \Gamma(X, \Omega^{\bullet \vee \vee}).$$

This notation is compatible with the symbol introduced in section 1 taking the biduality morphism (2.6) into account. We leave the remaining details with the reader. Q.E.D.

6. POINCARÉ DUALITY

Let X be a n -dimensional *oriented* smooth manifold. A compactly supported $(n-p)$ -form α on X defines a compact p-chain $P\alpha$ given by

$$(6.1) \quad \langle P\alpha, \beta \rangle = \int_X \alpha \wedge \beta, \quad \beta \in \Gamma(X, \Omega^p).$$

(6.2) THEOREM. *For a smooth oriented n-dimensional manifold X , the transformation P induces an isomorphism*

$$P: H_c^{n-p}(X, \mathbf{C}) \rightarrow H_p^c(X, \mathbf{C}), \quad p \in \mathbf{N},$$

from de Rham cohomology with compact support to de Rham homology.

Proof. The following diagram is commutative

$$(6.3) \quad \begin{array}{ccc} \Gamma_c(X, \Omega^{n-p}) & \xrightarrow{P} & D_p^c(X, \mathbf{C}) \\ \downarrow (-1)^n d & & \downarrow (-1)^{p-1} b \\ \Gamma_c(X, \Omega^{n-p+1}) & \xrightarrow{P} & D_{p-1}^c(X, \mathbf{C}) \end{array}$$

as it follows from the relation

$$d(\alpha \wedge \beta) = (d\alpha) \wedge \beta + (-1)^{n-p} \alpha \wedge d\beta, \quad \alpha \in \Gamma_c(X, \Omega^{n-p}), \beta \in \Gamma(X, \Omega^p),$$