

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 35 (1989)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON CHERN CLASSES OF FINITE GROUP REPRESENTATIONS
Autor: Kroll, Ove
Kapitel: Section 3. Proof of Theorem 5
DOI: <https://doi.org/10.5169/seals-57380>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 30.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

$$\begin{aligned} \varphi: T_n(q) &\rightarrow \mu_\infty, \\ c^{(1)}(d_{p_1}(\varphi)) &= c^{(2)}(d_{p_2}(\varphi)). \end{aligned}$$

But

$$d_{p_i}(\varphi) = e_{p_i}^{-1} \circ \varphi$$

and

$$c^{(i)}(d_{p_i}(\varphi)) = \varphi^* \circ (e_{p_i}^{-1})^* (e_{p_i}^*(u)) = \varphi^*(u).$$

Remark. It is necessary to reduce to \mathbf{Z}_{l^a} coefficients as the restriction map

$$H^*(Gl_n(q), \hat{\mathbf{Z}}_l) \rightarrow H^*(T_n(q), \hat{\mathbf{Z}}_l)$$

is not injective in general.

SECTION 3. PROOF OF THEOREM 5

CH1 and CH2 clearly follow from resp. CH1 and CH2 in Theorem 2 together with the functoriality of the decomposition map d_p i.e. the diagram

$$\begin{array}{ccc} R_c(G) & \xrightarrow{f^*} & R_c(H) \\ \downarrow d_p & & \downarrow d_p \\ R_p(G) & \xrightarrow{f^*} & R_p(H) \end{array}$$

is commutative for a group homomorphism $f: H \rightarrow G$. To obtain CH3 note that $d_p(\varphi) = e_p^{-1} \circ \varphi$ so by definition

$$c_1(\varphi) = c_1(d_p(\varphi)) = (e_p^{-1} \circ \varphi)^*(e_p^*(u)) = \varphi^* \circ (e_p^{-1})^* \circ e_p^*(u) = \varphi^*(u).$$

Furthermore let δ be the connecting homomorphism obtained from the exact sequence

$$\mathbf{Z} \rightarrow \mathbf{Q} \rightarrow \mathbf{Q}/\mathbf{Z}$$

As the diagram

$$\begin{array}{ccc} H^1(\mu_\infty, \mathbf{Q}/\mathbf{Z}) & \xrightarrow{\sim \delta} & H^2(\mu_\infty, \mathbf{Z}) \\ \downarrow \varphi^* & & \downarrow \varphi^* \\ H^1(G, \mathbf{Q}/\mathbf{Z}) & \xrightarrow{\sim \delta} & H^2(G, \mathbf{Z}) \end{array}$$

is commutative and as both δ 's are isomorphisms it suffices to show that

$$\delta^{-1} \circ c_1: \text{Hom}(G, \mathbf{C}^*) \rightarrow H^1(G, \mathbf{Q}/\mathbf{Z}) \simeq \text{Hom}(G, \mathbf{Q}/\mathbf{Z})$$

is an isomorphism.

But by inspection $\delta^{-1} \circ c_1(\varphi) = \delta^{-1}(u) \circ \varphi$, and as u is a $\hat{\mathbf{Z}} = \text{End}_{\mathbf{Z}}(\mu_{\infty})$ generator for $H^2(\mu_{\infty}, \mathbf{Z}) \simeq \hat{\mathbf{Z}}$, $\delta^{-1}(u)$ is an isomorphism

$$\delta^{-1}(u): \mu_{\infty} \xrightarrow{\sim} \mathbf{Q}/\mathbf{Z}.$$

SECTION 4. CHERN CLASSES FOR LOCALLY FINITE GROUPS

The definition will be based on the following two observations. In the following, let $G = \varinjlim G_k$ be a locally finite group where $\{G_k\}$ is a family of finite subgroups.

LEMMA. *Let*

$$\varphi: G \rightarrow \text{Gl}_n(\mathbf{C})$$

be a representation of G . Then φ is uniquely determined by its restrictions

$$\varphi_k: G_k \rightarrow \text{Gl}_n(\mathbf{C}).$$

Conversely given a family of compatible representations $\varphi_k: G_k \rightarrow \text{Gl}_n(\mathbf{C})$, there exists a unique $\varphi: G \rightarrow \text{Gl}_n(\mathbf{C})$ which restricts to φ_k for all k .

Proof. From the universal property of the direct limit, we have

$$\text{Hom}(G, \text{Gl}_n(\mathbf{C})) \cong \varprojlim \text{Hom}(G_k, \text{Gl}_n(\mathbf{C})).$$

PROPOSITION. *For all $i \geq 0$, the natural map*

$$H^i(G, \mathbf{Z}) \cong \varprojlim H^i(G_k, \mathbf{Z}).$$

is an isomorphism.

Proof. Obvious for $i = 0, 1$. For $i \geq 1$, the homology groups

$$H_i(G, \mathbf{Z}) = \varinjlim H_i(G_k, \mathbf{Z})$$

are all abelian torsion groups.

Now by the universal coefficient theorem ($i \geq 1$)