

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 35 (1989)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: KNOTTING CODIMENSION 2 SUBMANIFOLDS LOCALLY
Autor: Masuda, Mikiya / Sakuma, Makoto
Kapitel: Introduction
DOI: <https://doi.org/10.5169/seals-57360>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

KNOTTING CODIMENSION 2 SUBMANIFOLDS LOCALLY

by Mikiya MASUDA and Makoto SAKUMA

INTRODUCTION

Let L be a connected oriented n -dimensional closed manifold smoothly embedded in a connected oriented $(n+2)$ -dimensional closed manifold M , and let K be an oriented n -dimensional smooth knot in the oriented S^{n+2} . Then we consider the connected sum $(M, L) \# (S^{n+2}, K)$. In other words, we knot L locally using K . It yields another embedding of L in M ; however, it does not always give a new embedding. In fact, the lightbulb theorem says that the connected sum of $(S^2 \times S^1, \{*\} \times S^1)$ with any knot in S^3 is always equivalent to the original embedding. Moreover, by the prime decomposition theorem for knots in 3-manifolds [My], $(S^2 \times S^1, \{*\} \times S^1)$ is essentially the only embedding of a circle with the above property. Litherland [Li] has generalized the lightbulb theorem to the higher dimensional cases. In the appendix of [V], Viro exhibits an example of a 2-knot whose connected sum with the standard projective plane in S^4 does not change the isotopy type of the projective plane. (See also [La].)

The purpose of this paper is to study under what conditions this phenomenon occurs (or does not occur). The first named author [Ms] studied this problem when the codimension is greater than 2.

Put it in another way. Let \mathcal{K}_n be the set of isotopy classes of oriented n -knots diffeomorphic to S^n in the oriented S^{n+2} . The set forms an abelian monoid under connected sum for pairs. Analogously to the inertia group of a manifold, we define

$$I(M, L) = \{(S^{n+2}, K) \in \mathcal{K}_n \mid (M, L) \# (S^{n+2}, K) = (M, L)\}$$

where $=$ in the parenthesis indicates that there is an orientation preserving diffeomorphism of pairs. The set forms a submonoid of \mathcal{K}_n and describes the effect of knotting L locally. We are also concerned with the following intermediate submonoid

$$I_0(M, L) = \{(S^{n+2}, K) \in I(M, L) \mid (M, L) \# (S^{n+2}, K) \equiv (M, L)\}$$

where \equiv indicates that there is an orientation preserving diffeomorphism of pairs which is concordant to the identity map as a diffeomorphism of the ambient space M .

Our results suggest that $I(M, L)$ and $I_0(M, L)$ depend only on the order of a meridian of L in $\pi_1(M - L)$ or $H_1(M - L; \mathbf{Z})$. Roughly speaking, according as the order is infinite, 1, or p ($1 < p < \infty$), they can be distinguished by (at least) these three types:

Type 1 $I(M, L) = \{0N\}$,

Type 2 $I(M, L) = \mathcal{K}_n$, $I_0(M, L) = \ker \sigma$,

Type 3 $\{0\} \subsetneq I(M, L) \subsetneq \mathcal{K}_n$, $\{0\} \subsetneq I_0(M, L) \subsetneq \ker \sigma$,

(see section 4 for $\sigma(S^{n+2}, K)$).

We refer the reader to 1.1, 2.6, 3.4, 5.1, 5.2, and 5.8 for the precise statement.

This paper consists of five sections. In Section 1, we deduce a necessary condition for $I_0(M, L)$, which is valid for any (M, L) . We treat type 1 in Section 2. Type 2 is discussed in Sections 3, 4 and type 3 is discussed in Section 5. We will find that type 3 is closely related to the generalized Smith conjecture.

The authors would like to express their hearty thanks to Professors A. Kawauchi and T. Maeda for helpful conversations and suggestions.

§ 1. GENERAL REMARKS ON $I_0(M, L)$

It is known (and it is easily verified) that the signature of a Seifert surface of an oriented n -knot K in S^{n+2} is independent of the choice of a Seifert surface; so it is an invariant of the oriented knot K . The invariant is called the signature of the knot K and denoted by $\text{Sign}(S^{n+2}, K)$. We note that $\text{Sign}(S^{n+2}, K)$ is trivially zero unless $n + 1 \equiv 0 \pmod{4}$.

As is seen in Section 3, there is a pair (M^{n+2}, L^n) such that $I(M, L) = \mathcal{K}_n$ for any $n \geq 3$. In contrast, we can deduce a necessary condition for $I_0(M, L)$ which holds for any pair (M, L) .

THEOREM 1.1. *If $(S^{n+2}, K) \in I_0(M, L)$, then $\text{Sign}(S^{n+2}, K) = 0$.*

Proof. Let V be a Seifert surface of K . Since $S^{n+2} = \partial D^{n+3}$, we can push the interior of V into the interior of D^{n+3} so that V is transverse to S^{n+2} . This yields an oriented pair (D^{n+3}, V) having (S^{n+2}, K) as the boundary.