

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 35 (1989)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: KNOTTING CODIMENSION 2 SUBMANIFOLDS LOCALLY
Autor: Masuda, Mikiya / Sakuma, Makoto
Kapitel: §2. Type 1 case
DOI: <https://doi.org/10.5169/seals-57360>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

§ 2. TYPE 1 CASE

In this section we consider the case where a meridian of L^n in M^{n+2} has infinite order in $H_1(M-L; \mathbf{Z})$. We shall denote by $[m]$ the homology class in $H_1(M-L; \mathbf{Z})$ represented by a meridian m of L in M . For a manifold pair (X, Y) of codimension 2 and an epimorphism γ from $\pi_1(X-Y)$ to a finite group, let $(X, Y)_\gamma$ be the branched covering of (X, Y) corresponding to γ . Each knot group $\pi_1(S^{n+2}-K)$ has a natural epimorphism to \mathbf{Z}_p for any positive integer p , and the corresponding p -fold branched cyclic covering of (S^{n+2}, K) is denoted by $(S^{n+2}, K)_p$.

LEMMA 2.1. *Suppose $[m]$ is of infinite order. Then if $(S^{n+2}, K) \in I(M, L)$ then $(S^{n+2}, K)_p$ is a homotopy $(n+2)$ -sphere for any positive integer p .*

Proof. Since $[m]$ represents a nontrivial element in the finitely generated free abelian group $B_1(M-L) \equiv H_1(M-L; \mathbf{Z})/\text{Tor } H_1(M-L; \mathbf{Z})$, there is a positive integer r and a primitive element x in $B_1(M-L)$ such that $[m] = rx$ in $B_1(M-L)$. For each positive integer p , let γ_p be the canonical epimorphism $\pi_1(M-L) \rightarrow B_1(M-L) \otimes \mathbf{Z}_{pr}$. Noting the naturality of the homomorphism γ_p , we can see the following:

$$\begin{aligned} (M, L)_{\gamma_p} &= ((M, L) \# (S^{n+2}, K))_{\gamma_p \circ f_*} \\ &= (M, L)_{\gamma_p} \# d_p(S^{n+2}, K)_p \end{aligned}$$

Here f is a diffeomorphism $(M, L) \# (S^{n+2}, K) \rightarrow (M, L)$ and d_p is the order of $B_1(M-L) \otimes \mathbf{Z}_{pr}$ divided by p . Hence $H_*(S^{n+2}, K)_p; \mathbf{Z} \simeq H_*(S^{n+2}; \mathbf{Z})$ and $\pi_1((S^{n+2}, K)_p) \simeq 1$ by the existence of prime decompositions of finitely generated groups into free products [Wg]. Q.E.D.

It is conjectured that those knots which satisfy the conclusion of the above lemma are trivial. In fact, for $n = 1$, it follows from the Smith conjecture [MB]. As a supporting evidence for higher dimensional cases, we have

LEMMA. *Suppose that $(S^{n+2}, K)_p$ is a homotopy $(n+2)$ -sphere for every positive integer p . Then the Alexander modules of K are trivial.*

Proof. Let $\tilde{E}(K)$ be the infinite cyclic cover of the exterior $E(K)$ of K in S^{n+2} , and let t denote the automorphism of the homology group of $\tilde{E}(K)$ induced by the action of a meridian. Then, by the arguments of [Sm1],

we can see that $t^p - 1 : H_q(\tilde{E}(K); \mathbf{Z}_r) \rightarrow H_q(\tilde{E}(K); \mathbf{Z}_r)$ is an isomorphism for any positive integers p , q , and r . Assume r is prime. Then $H_q(\tilde{E}(K); \mathbf{Z}_r)$ is a finite abelian group, since it is a finitely generated torsion module over the principal ideal domain $\mathbf{Z}_r\langle t \rangle$ (see [Le3, p. 8]). So the automorphism t on $H_q(\tilde{E}(K); \mathbf{Z}_r)$ has a finite order, say d , and we have $t^d - 1 = 0$. Hence $H_q(\tilde{E}(K); \mathbf{Z}_r) = 0$, and by the universal coefficient theorem, the following holds for any prime r and any positive integer q :

$$(2.3) \quad H_q(\tilde{E}(K); \mathbf{Z}) \otimes \mathbf{Z}_r = 0$$

$$(2.4) \quad \text{Tor}(H_q(\tilde{E}(K); \mathbf{Z}), \mathbf{Z}_r) = 0$$

By (2.4), $H_q(\tilde{E}(K); \mathbf{Z})$ has no nontrivial elements of finite order; so it has a square presentation matrix $M(t)$ as a $\mathbf{Z}\langle t \rangle$ -module by [Le3, Proposition 3.5]. By (2.3) the q -th Alexander polynomial $\det M_q(t)$ ($\in \mathbf{Z}\langle t \rangle$) is a unit mod. r for any prime r . Hence it is a unit in $\mathbf{Z}\langle t \rangle$, and we have $H_q(\tilde{E}(K); \mathbf{Z}) = 0$ for any positive integer q . Q.E.D.

Thus, as a consequence of Lemmas 2.1 and 2.2 and the results of [Le2] and [T], we have the following:

PROPOSITION 2.5. *Suppose $[m]$ is of infinite order. Then any knot in $I(M, L)$ has trivial Alexander modules and is null cobordant.*

Hence the only obstruction for a knot (S^{n+2}, K) in $I(M, L)$ to be trivial lies in the knot group $\pi_1(S^{n+2} - K)$. For the special case where $[m]$ generates $H_1(M - L)$, we can apply the result of Maeda [Ma] (cf. [DF]), and obtain the following:

THEOREM 2.6. *Suppose $n \geq 3$ and $H_1(M - L)$ is the infinite cyclic group generated by $[m]$. Then $I(M, L)$ is trivial.*

Proof. Let (S^{n+2}, K) be a knot in $I(M, L)$. Note that $\pi_1(M - L)$ is isomorphic to the amalgamated free product $\pi_1(M - L) *_{\langle m \rangle} \pi_1(S^{n+2} - K)$.

Then we can conclude $\pi_1(S^{n+2} - K) \simeq \mathbf{Z}$ by the result of [Ma] (cf. [DF]) which asserts the existence of a prime decomposition of a finitely presented group G with $G/[G, G] \simeq \mathbf{Z}$ with respect to such amalgamated free products. Combined with Proposition 2.5, we see $S^{n+2} - K$ is homotopy equivalent to a circle. Hence (S^{n+2}, K) is trivial by [Le1].