Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	35 (1989)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	HURWITZ-RADON MATRICES AND PERIODICITY MODULO 8
Autor:	Eckmann, Beno
Kapitel:	0. Introduction
DOI:	https://doi.org/10.5169/seals-57365

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

HURWITZ-RADON MATRICES AND PERIODICITY MODULO 8

by Beno Eckmann

0. INTRODUCTION

0.1. We consider complex $n \times n$ — matrices $A_1, A_2, ..., A_s$, either all unitary (case U) or all orthogonal (case O); they are called Hurwitz-Radon matrices, in short HR-matrices, if

(1)
$$A_{i}^{2} = -E, A_{i}A_{k} + A_{k}A_{j} = 0, \quad j, k = 1, 2, ..., s, j \neq k;$$

E or E_n denotes the unit matrix. Such matrices are well-known to exist, even with entries $0, \pm 1, \pm i$ (case *U*) or $0, \pm 1$ (case *O*). The possible values of *n* are multiples $mn_0, m = 1, 2, 3, ...$ where in case $U, n_0 = 2^{s/2}$ if *s* is even, $n_0 = 2^{(s-1)/2}$ if *s* is odd. In case $0, n_0 = 2^{(s-1)/2}$ if $s \equiv 7 \mod 8$; $n_0 = 2^{s/2}$ if $s \equiv 0, 6$; $n_0 = 2^{(s+1)/2}$ if $s \equiv 1, 3, 5$; and $n_0 = 2^{(s+2)/2}$ if $s \equiv 2, 4 \mod 8$.

If we put $A_0 = E$ the relations (1) are equivalent to

$$f_s(x_0, x_1, ..., x_s) = \sum_{0}^{s} x_j A_j$$

being a unitary, or orthogonal respectively, matrix for all real x_j with $\sum_{0}^{s} x_j^2 = 1$. Thus f_s can be considered as a map $S^s \to U$ via U(n), or $S^s \to O$ via O(n) where $U = \lim_{s \to 0} U(k)$ and $O = \lim_{s \to 0} O(k)$ are the infinite unitary and orthogonal groups. We also write f_s for the homotopy class of that map, $f_s \in \pi_s(U)$ or $\pi_s(O)$. We recall that by the Bott periodicity theorems these groups are cyclic or 0.

THEOREM A. If $A_1, A_2, ..., A_s$ are HR-matrices of minimal size $n = n_0(s)$ then f_s is a generator of $\pi_s(U)$, or $\pi_s(O)$ respectively, s = 0, 1, 2,

Remark 1. For s = 0 (empty set of HR-matrices) we have $f_0(x_0) = x_0(1)$, $x_0^2 = 1$; i.e., $f_0(1) = (1)$, $f_0(-1) = (-1)$, $f_0: S^0 \to O(1) \to O$. For s > 0, $f_0: S^s \to O$ clearly factors through $SO(n) \to SO$ (U being connected, the analogue is irrelevant in the unitary case).

B. ECKMANN

78

Remark 2. The problem originally solved by Hurwitz [H] and Radon [R] refers to the case O: One asks for complex bilinear forms $z = f(x, y) = (\sum_{0}^{s} x_{j}A_{j})y$, where $z = (z_{1}, ..., z_{n}), y = (y_{1}, ..., y_{n}), x = (x_{0}, ..., x_{s})$, such that $z_{1}^{2} + ... + z_{n}^{2} = (x_{0}^{2} + ... + x_{s}^{2})(y_{1}^{2} + ... + y_{n}^{2}).$

This means that $\sum_{0}^{s} x_{j}A_{j}$ is orthogonal, i.e. leaves invariant $\sum_{0}^{n} y_{j}^{2}$ except for the factor $\sum_{0}^{s} x_{j}^{2}$; and thus, since we may assume $A_{0} = E$, that $A_{1}, ..., A_{s}$ is a set of orthogonal HR-matrices in the sense of (1).

The case U refers to the analogous problem for the identity

$$|z_1|^2 + ... + |z_n|^2 = (x_0^2 + ... + x_s^2)(|y_1|^2 + ... + |y_n|^2)$$

where y and z are complex, and x real.

0.2. The symplectic case: It is also of interest to consider HR-matrices, i.e. matrices fulfilling (1), which are symplectic. A linear combination $\sum_{0}^{s} x_{j}A_{j}$ of $2n \times 2n$ -matrices with $A_{0} = E$ is symplectic up to the factor $\sum_{0}^{s} x_{j}^{2}$ if and only if $A_{1}, ..., A_{s}$ is a set of symplectic HR-matrices (Proposition 4.1).

We restrict attention to unitary symplectic matrices, i.e., to the group $Sp(n) \subset U(2n)$, and write Sp for the infinite symplectic group $\lim_{x \to 0} Sp(k)$. With a set $A_1, ..., A_s$ of unitary symplectic HR-matrices, and $A_0 = E$, we associate the map $f_s(x_0, x_1, ..., x_s) = \sum_{s=0}^{s} x_j A_j$, $\sum_{s=0}^{s} x_j^2 = 1$, of S^s into Sp via Sp(n); we also write f_s for the corresponding element of $\pi_s(Sp)$, known to be 0 or cyclic.

THEOREM A'. If $A_1, ..., A_s$ are unitary symplectic HR-matrices of minimal size $2n_0$ then f_s is a generator of $\pi_s(Sp)$.

0.3. The paper is organized as follows. We first recall (Section 1) that the HR-matrix problem can be formulated in terms of representations of certain finite group G_s , s = 0, 1, 2, ... introduced by the author [E], and discuss these representations using the elegant description of [LS]. In Section 2 the "reduced" Grothendieck groups of representations E_s^U and E_s^o are

computed; they turn out to be isomorphic to $\pi_s(U)$ and $\pi_s(O)$ respectively. Moreover a product is defined in the direct sum of the $E_s^U(E_s^O)$ turning it into a graded ring $E_*^U(E_*^O)$. The claim of Theorem A is proved in Section 3; we show that the maps $\phi: E_s^U \to \pi_s(U), \psi: E_s^O \to \pi_s(O)$ given by the f_s of 0.1 are isomorphisms. Using the product structure in $\pi_*(U)$ and $\pi_*(O)$ known from K-theory the proof reduces to simple verifications in low dimensions. The symplectic case is dealt with in Section 4. In Section 5 we make a remark concerning the "linearization phenomenon" for the homotopy groups of U, O and Sp.

1. The groups G_s and their representations

We will denote throughout by G_s the group given by the presentation 1.1. $G_s = \langle \varepsilon, a_1, ..., a_s | \varepsilon^2 = 1, a_j^2 = \varepsilon, a_j a_k = \varepsilon a_k a_j, j, k = 1, 2, ..., s, j \neq k > .$ Clearly any set $A_1, ..., A_s$ of HR-matrices yields a (unitary or orthogonal) representation of G_s of degree *n* by $\varepsilon \mapsto -E$, $a_j \mapsto A_j$, j = 1, 2, ..., s. Conversely a representation of G_s with $\varepsilon \mapsto -E$, in short an ε -representation, yields a set of s HR-matrices. For the elementary properties of G_s and its representations we refer to [E]. We just recall that the order of G_s is 2^{s+1} , that ε is central, and that the irreducible unitary ε -representations of G_s are of degree $2^{s/2}$ if s is even (one equivalence class), of degree $2^{(s-1)/2}$ if s is odd (two equivalence classes). These degrees are the minimal values n_0 in case U. As for the case O, one has to recall that a representation is equivalent to an orthogonal one if and only if it is equivalent to a real (and orthogonal) one. Thus, unless an irreducible unitary ε -representation is already real, one has to add its conjugate-complex representation, and the discussion of the various cases depending on s yields the minimal values n_0 (case O) mentioned in the introduction; in other words, the degrees of the irreducible orthogonal ε -representations of G_{s} .

1.2. A very simple and useful scheme for studying the groups G_s and their ε -representations (and many other things) has been deviced by T. Y. Lam and T. Smith [LS]. They have expressed the G_s as products of very small and well-known groups. Namely $C = G_1$, the cyclic group of order 4; $Q = G_2$, the quaternionic group of order 8; K, the Klein 4-group; and D, the dihedral group of order 8. Although K and D do not belong to the family G_s , they are of a similar nature and contain a distinguished central element ε of order 2 (distinguished arbitrarily in K). "Product" here means the central product obtained from the direct product by identifying the