

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	35 (1989)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
 Artikel:	HURWITZ-RADON MATRICES AND PERIODICITY MODULO 8
Autor:	Eckmann, Beno
Kapitel:	1. The groups G_s and their representations
DOI:	https://doi.org/10.5169/seals-57365

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 24.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

computed; they turn out to be isomorphic to $\pi_s(U)$ and $\pi_s(O)$ respectively. Moreover a product is defined in the direct sum of the $E_s^U(E_s^O)$ turning it into a graded ring $E_*^U(E_*^O)$. The claim of Theorem A is proved in Section 3; we show that the maps $\phi: E_s^U \rightarrow \pi_s(U)$, $\psi: E_s^O \rightarrow \pi_s(O)$ given by the f_s of 0.1 are isomorphisms. Using the product structure in $\pi_*(U)$ and $\pi_*(O)$ known from K -theory the proof reduces to simple verifications in low dimensions. The symplectic case is dealt with in Section 4. In Section 5 we make a remark concerning the “linearization phenomenon” for the homotopy groups of U , O and Sp .

1. THE GROUPS G_s AND THEIR REPRESENTATIONS

1.1. We will denote throughout by G_s the group given by the presentation

$$G_s = \langle \varepsilon, a_1, \dots, a_s \mid \varepsilon^2 = 1, a_j^2 = \varepsilon, a_j a_k = \varepsilon a_k a_j, j, k = 1, 2, \dots, s, j \neq k \rangle.$$

Clearly any set A_1, \dots, A_s of HR-matrices yields a (unitary or orthogonal) representation of G_s of degree n by $\varepsilon \mapsto -E$, $a_j \mapsto A_j$, $j = 1, 2, \dots, s$. Conversely a representation of G_s with $\varepsilon \mapsto -E$, in short an ε -representation, yields a set of s HR-matrices. For the elementary properties of G_s and its representations we refer to [E]. We just recall that the order of G_s is 2^{s+1} , that ε is central, and that the irreducible unitary ε -representations of G_s are of degree $2^{s/2}$ if s is even (one equivalence class), of degree $2^{(s-1)/2}$ if s is odd (two equivalence classes). These degrees are the minimal values n_0 in case U . As for the case O , one has to recall that a representation is equivalent to an orthogonal one if and only if it is equivalent to a real (and orthogonal) one. Thus, unless an irreducible unitary ε -representation is already real, one has to add its conjugate-complex representation, and the discussion of the various cases depending on s yields the minimal values n_0 (case O) mentioned in the introduction; in other words, the degrees of the irreducible orthogonal ε -representations of G_s .

1.2. A very simple and useful scheme for studying the groups G_s and their ε -representations (and many other things) has been devised by T. Y. Lam and T. Smith [LS]. They have expressed the G_s as products of very small and well-known groups. Namely $C = G_1$, the cyclic group of order 4; $Q = G_2$, the quaternionic group of order 8; K , the Klein 4-group; and D , the dihedral group of order 8. Although K and D do not belong to the family G_s , they are of a similar nature and contain a distinguished central element ε of order 2 (distinguished arbitrarily in K). “Product” here means the central product obtained from the direct product by identifying the

two ε 's. The expression for the G_s then is as follows, displaying a fundamental periodicity modulo 8:

(2)	s	0	1	2	3	4	5	6	7	8	9	...
	G_s	$\mathbf{Z}/2$	C	Q	QK	QD	D^2C	D^3	D^3K	D^4	D^4C	...

and $G_{s+8} = D^4G_s$.

The tensor product of ε -representations of two of the groups G_s, K, D is an ε -representation of their product above, and all ε -representations of the G_s can be obtained in that explicit way from those of C, Q, K, D , which are well-known. This yields, in particular, the characters χ and the Schur indices I of the irreducible unitary ε -representation (the Schur index $I = 1$ if the representation is equivalent to a real one; if it is not, $I = -1$ if it is equivalent to the conjugate-complex one, $I = 0$ otherwise). Both χ and I behave multiplicatively with respect to the central product.

1.3. The Schur indices of the irreducible ε -representations are: 0 for $C = G_1$, -1 for $Q = G_2$, and 1 for K and D (two equivalence classes for K , one for D). This yields the Schur indices I_s of the irreducible ε -representations of the G_s , as listed in (2) below; we further list the numbers v_s^U of inequivalent unitary, and v_s^O of inequivalent orthogonal irreducible ε -representations, and the respective degrees d_s^U, d_s^O . Note that I_s is periodic with period 8, and d_s^O is periodic with period 8 in the sense that $d_{s+8}^O = 16d_s^O$. Finally we include in the same table the Grothendieck groups D_s^U and D_s^O of (equivalence classes of) irreducible ε -representations of G_s , with respect to the direct sum of representations.

(3)	s	0	1	2	3	4	5	6	7	8	9	...
	I_s	1	0	-1	-1	-1	0	1	1	1	0	...
	v_s^U	1	2	1	2	1	2	1	2	1	2	
	v_s^O	1	1	1	2	1	1	1	2	1	1	
	d_s^U	1	1	2	2	4	4	8	8	16	16	
	d_s^O	1	2	4	4	8	8	8	8	16	32	
	D_s^U	\mathbf{Z}	$\mathbf{Z} \oplus \mathbf{Z}$									
	D_s^O	\mathbf{Z}	\mathbf{Z}	\mathbf{Z}	$\mathbf{Z} \oplus \mathbf{Z}$	\mathbf{Z}	\mathbf{Z}	\mathbf{Z}	$\mathbf{Z} \oplus \mathbf{Z}$	\mathbf{Z}	\mathbf{Z}	

The values of d_s^O follow immediately from the I_s and the d_s^U . The values n_0 for the case O , as given in the Introduction, are the d_s^O .

2. THE REDUCED ε -REPRESENTATION RING

2.1. For all $s \geq 0$ the group G_s is the subgroup of G_{s+1} obtained by omitting the generator a_{s+1} ; let $h_s: G_s \rightarrow G_{s+1}$ be the embedding homomorphism. Via h_s we can restrict an ε -representation of G_{s+1} to G_s , which in terms of HR-matrices means omitting A_{s+1} .

Let $h_s^*: D_{s+1}^U \rightarrow D_s^U$ be the corresponding homomorphism of Grothendieck groups, and $E_s^U = D_s^U/h_s^* D_{s+1}^U$ the “reduced” groups; similarly $E_s^O = D_s^O/h_s^* D_{s+1}^O$. They can easily be computed by means of the characters of ε -representations, as follows.

For Q and D the character of an irreducible unitary ε -representation is 0 except on 1 and ε . For C and K it is $\neq 0$ on all 4 elements; on the essential generator ($\neq \varepsilon$) of C it is $+i$ or $-i$ for the two inequivalent representations, and $+1$ or -1 in the case of K . For G_s , s even, we infer from the table (2) that the character is 0 except on 1, ε . For G_s , s odd, the character is 0 except on 1, ε and two further elements $z, \varepsilon z$; on these the two inequivalent ε -representations differ just by the sign of the character.

If s is even, $d_{s+1}^U = d_s^U = 2^{s/2}$; thus the restriction of an irreducible ε -representation must be irreducible, whence $h_s^* D_{s+1}^U = D_s^U$, $E_s^U = 0$. If s is odd, $d_{s+1}^U = 2d_s^U = 2^{(s+1)/2}$; thus the restriction is the sum of two irreducible ε -representations, and since the character is 0 (except on 1, ε) these two must be inequivalent. Therefore $h_s^* D_{s+1}^U$ is the “diagonal” of $D_s^U = \mathbf{Z} \oplus \mathbf{Z}$, and $E_s^U = \mathbf{Z}$; its generator ρ_s is represented by either of the two inequivalent irreducible ε -representations of G_s , $-\rho_s$ by the other one.

In the orthogonal case the E_s^O are computed similarly from (3). Since $d_1^O = 2$ and $d_0^O = 1$, the restriction from D_1^O to D_0^O yields twice the generator, and $E_0^O = \mathbf{Z}/2$; the same argument holds for $s \equiv 0 \pmod{8}$, $d_{s+1}^O = 2d_s^O$. Since $d_2^O = 4$ and $d_1^O = 2$, we get $E_1^O = \mathbf{Z}/2$. From $d_3^O = d_2^O = 4$ we get $E_2^O = 0$. As for $s = 3$, the character argument shows that $h_3^* D_4^O$ = diagonal of $D_3^O (= \mathbf{Z} \oplus \mathbf{Z})$, and $E_3^O = \mathbf{Z}$. For $s = 4, 5, 6$ the dimensions $d_{s+1}^O = d_s^O$ show that $E_4^O = E_5^O = E_6^O = 0$. For $s = 7$, the character argument yields $h_7^* D_8^O$ = diagonal of $D_7^O (= \mathbf{Z} \oplus \mathbf{Z})$, and $E_7^O = \mathbf{Z}$. Finally one has, for all s , $E_{s+8}^O \cong E_s^O$.