

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 35 (1989)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE FIXED POINT SET OF A FINITE GROUP ACTION ON A
HOMOLOGY FOUR SPHERE
Autor: Demichelis, Stefano
Kapitel: 5. LOCALLY LINEAR REPRESENTATION
DOI: <https://doi.org/10.5169/seals-57368>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

$\text{Fix}(A_4)$ is a sphere. It cannot be S^2 since the representation of A_4 in $SO(3)$ is irreducible, so it is S^1 . The only closed 1-dimensional submanifold of S^1 is S^1 itself, so $\text{Fix}(G) = S^1$.

b. As in subcase a., a linear change in coordinates allows us to assume that h is actually \tilde{i} , and as before if $G_2 \in G$ the proposition is proved applying 4.1.

If it is not the case, let α correspond to the cycle $(12345) \in A_5$, β to (123) and γ to (345) . We observe that β and γ generate A_5 and so:

1. $\text{Fix}(A_5) = \text{Fix}(\beta) \cap \text{Fix}(\gamma)$,
2. $\text{Fix}(A_5) \subset \text{Fix}(\alpha)$.

We claim that $\text{Fix}(\alpha)$ is S^0 . According to Smith's theorem it is enough to prove that the representation of α around x_0 has an isolated fixed point, i.e. is the sum of two irreducible complex ones.

If not by Lemma 3.3 $(\bar{i}(\alpha); i(\alpha))$ would be conjugate in $SO(3) \times SO(3)$ to an element on the diagonal. From the explicit description of i and \bar{i} (see the end of section 7.1 of [22]), it follows that they send all the five cycles to non conjugate elements in $SO(3)$, so this is impossible, and $\text{Fix}(\alpha) = S^0$.

As for β and γ , their images under (\bar{i}, i) are conjugate to elements on the diagonal, by 3.3 and 3.4 their fixed point sets have two-dimensional components, and so by Smith's theorem they are copies of S^2 .

So $\text{Fix}(G)$ is the intersection of a couple of S^2 s and is contained in $\text{Fix}(\alpha)$ which is S^0 . If this set is empty or equal to S^0 , the proposition follows. If it were a single point, it would be a transverse intersection, by local linearity, but it is not possible since a homology S^4 does not contain any two cycles with intersection number odd. This ends the proof.

5. LOCALLY LINEAR REPRESENTATION

Let's now consider the case of G acting on a homology S^4 with two fixed points, P_0 and P_1 .

THEOREM 5.1. *The unoriented representations of G around P_0 and P_1 are linearly equivalent.* ¹⁾

Proof. It will suffice to show that the characters associated to the representations around the P_i s agree on every cyclic subgroup C_k of G .

¹⁾ See the note in the introduction.

Observe that by Lemma 3.4 and Smith's theorem the fixed point set of an element of G different from the identity is either S^0 or S^2 .

Let g generate C_k , we distinguish three cases:

1. $\text{Fix}(g^r) = \{P_1; P_2\}$ for every $r \equiv 0(\text{mod } k)$,
2. $\text{Fix}(g) = S^2$,
3. $\text{Fix}(g) = \{P_1; P_2\}$ but $\text{Fix}(g^n) = S^2$ for some $g^n \neq \text{id}$.

Case 1. The hypothesis means that the action is semifree and the claim follows from the work of Atiyah and Bott, see [1] and [14].

Case 2. The action of C_k on the normal bundle of the fixed S^2 defines an element N of $K_{C_k}(S^2)$. Since C_k acts trivially on S^2 the two inclusions $P_i \rightarrow S^2$ are obviously C_k homotopic so that the diagram:

$$\begin{array}{ccccc}
 & & K_{C_k}(P_2) & & \\
 & \nearrow & & \searrow & \\
 [N] \in K_{C_k}(S^2) & & & & R(C_k) \\
 & \searrow & & \nearrow & \\
 & & K_{C_k}(P_1) & &
 \end{array}$$

commutes. This means that the representation of C_k in the normal component to S^2 are conjugate, the tangential representations are of course both the identity, so the statement is proved.

Case 3. We can assume, by [8], that the action on $S^2 = \text{Fix}(g^n)$ is linear. S^2 has zero intersection number in Σ so its normal bundle N can be identified to $S^2 \times R^2$, and we fix a trivialization. Denote a point of $S^2 - \{P_1; P_2\}$ by (x, t) with $x \in S^1$ and $t \in (0, 1)$. Let C_0 be the space $\{\phi: S^1 \rightarrow SO(2) \mid \text{deg } \phi = 0\}$, it is an abelian group by pointwise multiplication and a C_k module with structure given by:

$$(h\phi)(x) = \phi(hx), \quad h \in C_k \quad \text{and} \quad x \in S^1 \subset S^2$$

acted on by the obvious induced action.

By [5], chapter VI, prop. 11.1, the action is given by a θ_t such that

1. $\theta_t \in Z^1(C_k; C_0)$ and depends continuously on $t \in [0, 1]$.
2. $\theta_i(h)(x)$ is constant on $x \in S^1$ and equal to the representation of h at P_i for $i = 0; 1$.

A change in the trivialization adds to each θ_t a coboundary so there is a well defined continuous family $\theta_t: [0, 1] \rightarrow H^1(C_k; C_0)$.

A straightforward calculation shows that $H^1(C_k; C_0) = H^2(C_k; Z) = C_k$. Since θ_t is continuous it has to be constant, so $\theta_0 = \theta_1$ and by 2. the

two normal representations are equal. In the topological case, by the results of Cappel and Shaneson topological equivalence of matrices in dimension 4 implies linear equivalence, so the statement of Theorem 5.1 makes sense also for a group of homeomorphism.

The proof given can be adapted to this more general case provided that the followings are true:

1. the topological Atiyah-Singer signature formula holds,
2. a locally flat S^2 in Σ has a normal bundle,
3. the argument in case 3 works with $\text{Homeo}(S^1)$ instead of $SO(2)$.

Assertion 1 is proved, in the case of the semi-free action, in [21], page 188; assertion 2 follows from the work of Freedman, see [10]; assertion 3 is proved using the retraction $\text{Homeo}(S^1)$ into $SO(2)$ given by the Poincaré number, see [7].

APPENDIX

LEMMA. *The extensions:*

$$\begin{array}{ccccccccc}
 0 & \rightarrow & C_2 & \rightarrow & \tilde{A}_5 & \rightarrow & A_5 & \rightarrow & 0 \\
 & & & & \downarrow & & \downarrow & & \\
 0 & \rightarrow & C_2 & \rightarrow & A_5 \times A_5 & \rightarrow & A_5 \times A_5 & \rightarrow & 0 \\
 & & & & \downarrow & & \downarrow^{(h, h')} & & \\
 0 & \rightarrow & C_2 & \rightarrow & SO(4) & \rightarrow & SO(3) \times SO(3) & \rightarrow & 0
 \end{array}$$

are not split, h and h' can be any nontrivial representations of A_5 and f is either $(Id \times \{I\})$ or $(\{I\} \times Id)$.

Proof. Standard theory of group extensions and cohomology (see [4]) allows us to reduce to the:

PROPOSITION. *Any non trivial homomorphism $A_5 \xrightarrow{i} SO(3)$ induces an isomorphism $Z/2 = H^2(BSO(3); Z/2) \xrightarrow{i} H^2(BA_5; Z/2) = Z/2$.*

Proof of the Proposition. If the corresponding extension is split, then $Z/2 \times A_5 \subset S^3$, but $A_5 = 60$ so there exists a $Z/2 \subset A_5$ so $Z/2 \times Z/2$ would act freely on S^3 , which cannot happen.