

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 35 (1989)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: AN ELEMENTARY PROOF OF A THEOREM ON QUADRATIC FORMS OVER THE RATIONAL NUMBERS
Autor: Leep, David B.
Kapitel: §3. Consequences of the Main Theorem
DOI: <https://doi.org/10.5169/seals-57371>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

At this point we simply count lattice points directly and construct the table below. The table shows $P(\sqrt{p}) > \frac{p-1}{4}$ for all primes $p < 100$ except $p = 5$. This completes the proof of Theorem 1.7 and hence the Main Theorem has been proved completely.

p	$P(\sqrt{p})$	Range of primes where $P(\sqrt{p}) > \frac{p-1}{4}$
59	27	$59 \leq p \leq 97$
31	15	$31 \leq p \leq 53$
19	9	$19 \leq p \leq 29$
11	5	$11 \leq p \leq 17$
7	3	$p = 7$
3	1	$p = 3$
5	1	$1 \not> \frac{5-1}{4}$

§ 3. CONSEQUENCES OF THE MAIN THEOREM

In this section we derive some consequences of the Main Theorem that have applications to the algebraic theory of quadratic forms. The results in this section are well known ([La], Chapter 6). The point is that we have new and more elementary proofs.

Let $\langle\langle a, b, c \rangle\rangle$ denote the 3-fold Pfister form

$$\langle 1, a \rangle \otimes \langle 1, b \rangle \otimes \langle 1, c \rangle = \langle 1, a, b, c, ab, ac, bc, abc \rangle.$$

3.1. PROPOSITION. *Let $a, b, c \in \mathbf{Q}^\times$. Then $\langle\langle a, b, c \rangle\rangle$ is hyperbolic over \mathbf{Q} if and only if a, b, c are not all positive.*

Proof. If $\langle\langle a, b, c \rangle\rangle$ is hyperbolic, then consideration of $\langle\langle a, b, c \rangle\rangle$ over the field of real numbers shows at least one of a, b, c is negative.

Now suppose $a < 0$. Then the Main Theorem implies $-c \in D_{\mathbf{Q}}(\langle\langle a, b \rangle\rangle)$ and $\langle\langle a, b \rangle\rangle \perp \langle c \rangle$ is isotropic over \mathbf{Q} by Lemma 1.2(b). A theorem of Pfister ([La], p. 279] implies $\langle\langle a, b, c \rangle\rangle$ is hyperbolic over \mathbf{Q} .

3.2. PROPOSITION. Let $a, b, c \in \mathbf{Q}^\times$, $a, b, c > 0$. Then

$$\langle\langle a, b, c \rangle\rangle \cong \langle\langle 1, 1, 1 \rangle\rangle = 8\langle 1 \rangle.$$

Proof. Calculating in the Witt ring WF we have

$$\begin{aligned} \langle\langle a, b, 1 \rangle\rangle \perp (-1)\langle\langle a, b, c \rangle\rangle &= \langle\langle a, b \rangle\rangle (\langle 1, 1 \rangle \perp (-1)\langle 1, c \rangle) \\ &= \langle\langle a, b \rangle\rangle \langle 1, -c \rangle = \langle\langle a, b, -c \rangle\rangle = 0 \text{ by Proposition 3.1.} \end{aligned}$$

Therefore $\langle\langle a, b, 1 \rangle\rangle \cong \langle\langle a, b, c \rangle\rangle$. Repeating the same calculation with a, b in place of c yields the result.

3.3. COROLLARY. Let $a, b, c \in \mathbf{Q}^\times$ and let $\mathbf{H} = \langle 1, -1 \rangle$. Then

$$\langle\langle a, b, c \rangle\rangle \cong \begin{cases} \langle\langle 1, 1, 1 \rangle\rangle & \text{if } a, b, c > 0 \\ 4\mathbf{H} & \text{otherwise.} \end{cases}$$

3.4. THEOREM. $I^3\mathbf{Q}$ is torsion-free.

Proof. Corollary 3.3 shows that the only nonzero 3-fold Pfister form in $I^3\mathbf{Q}$ is $\langle\langle 1, 1, 1 \rangle\rangle$. Therefore $I^3\mathbf{Q} \cong \mathbf{Z}$ and $I^3\mathbf{Q}$ is torsion-free.

REFERENCES

- [An] ANDREWS, G. *Number Theory*. W.S. Saunders, Philadelphia, 1971.
- [BS] BOREVICH, Z. I. and I. R. SHAFAREVICH. *Number Theory*. Academic Press, New York, 1966.
- [Ca] CASSELS, J. W. S. *Rational Quadratic Forms*. Academic Press, New York, 1978.
- [Ha] HASSE, H. *Vorlesungen Über Zahlentheorie*. Springer-Verlag, Berlin, 1964.
- [HW] HARDY, G. H. and E. M. WRIGHT. *An Introduction to the Theory of Numbers*. Oxford University Press, 4th ed., 1960.
- [La] LAM, T. Y. *The Algebraic Theory of Quadratic Forms*. Benjamin, 1973.
- [Mo] MORDELL, L. J. *Diophantine Equations*. Academic Press, New York, 1969.
- [Om] O'MEARA, O. T. *Introduction to Quadratic Forms*. Springer-Verlag, Berlin, 1963.
- [Se] SERRE, J.-P. *A Course in Arithmetic*. Springer-Verlag, 1973.
- [Sk] SKOLEM, Th. On the Diophantine Equation $ax^2 + by^2 + cz^2 + dv^2 = 0$. *Norske Videnskabers Selsk. Forh.* 21 (1948), 76-79.

(Reçu le 10 avril 1989)

David B. Leep

Department of Mathematics
University of Kentucky
Lexington, KY 40506
(USA)