

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 36 (1990)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** THE DISTANCE BETWEEN IDEALS IN THE ORDERS OF A REAL QUADRATIC FIELD  
**Autor:** Kaplan, Pierre / Williams, Kenneth S.  
**Kapitel:** 1. Introduction  
**DOI:** <https://doi.org/10.5169/seals-57912>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 05.03.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## THE DISTANCE BETWEEN IDEALS IN THE ORDERS OF A REAL QUADRATIC FIELD

par Pierre KAPLAN and Kenneth S. WILLIAMS <sup>1)</sup>

### 1. INTRODUCTION

The notion of the distance between two equivalent, reduced, primitive ideals of an order in the ring of integers of a real quadratic field was first introduced by Shanks [7] in 1972 in order to develop a more efficient algorithm for computing the fundamental unit of the field, although this notion was already implicit in the work of earlier authors including Lagrange [2]. Shanks used the language of binary quadratic forms to describe the concept of distance. This concept, still described in terms of binary quadratic forms, was made more precise and exploited by Lenstra [4] (1982) and Schoof [6] (1983) in their work on quadratic fields and factorization. In 1986 Williams and Wunderlich [12] gave a treatment of distance in terms of ideals, and used it to develop a simple algorithm for use in the continued fraction factoring algorithm. Parts of their theory have also been used in numerical studies of Eisenstein's problem [9] [11].

The aim of this paper is two-fold. We first give a complete treatment of the basic theory of the distance between equivalent, reduced, primitive ideals in the hope of making this attractive and useful theory better known and more readily available for further research. Our treatment is based mainly on the presentation of Williams and Wunderlich [12], but, in our view, is simpler in some aspects. Our second objective is to define a homomorphism between the ideal class groups of different orders and to apply this theory to compare distances between corresponding ideals in the two orders. The presentation is self-contained in that factorization of ideals in an order of a quadratic field is not needed, nor do we use the theory of the units of a real quadratic field. Indeed the theory of units is a consequence of our presentation, see Corollary 5. We give known results as Propositions and new results as Theorems.

---

<sup>1)</sup> Research supported by Natural Sciences and Engineering Research Council of Canada Grant A-7233.

Throughout this paper, if  $A$  is a unitary commutative ring, and  $\alpha_1, \alpha_2, \dots, \alpha_m$  are elements of  $A$ , the  $\mathbb{Z}$ -module generated by  $\alpha_1, \alpha_2, \dots, \alpha_m$  is denoted by  $[\alpha_1, \alpha_2, \dots, \alpha_m]$  and the  $A$ -module (ideal) generated by  $\alpha_1, \alpha_2, \dots, \alpha_m$  by  $(\alpha_1, \alpha_2, \dots, \alpha_m)$ . The product of the ideals  $(\alpha_1, \dots, \alpha_m)$  and  $(\alpha'_1, \dots, \alpha'_n)$  is the ideal  $(\alpha_1 \alpha'_1, \dots, \alpha_i \alpha'_j, \dots, \alpha_m \alpha'_r)$ . If  $I$  is an ideal, we often write the product ideal  $(\alpha)I$  as  $\alpha I$ .

## 2. BASIC DEFINITIONS

Let  $K$  be a quadratic field of discriminant  $D_0$ . As  $D_0$  is a discriminant we have  $D_0 \equiv 0 \pmod{4}$  or  $D_0 \equiv 1 \pmod{4}$ . In §2 and §3  $K$  may be real ( $D_0 > 0$ ) or imaginary ( $D_0 < 0$ ) but in the remaining sections  $K$  will be assumed to be real. An element  $\alpha$  of  $K$  can be written  $\alpha = x + y\sqrt{D_0}$ , where  $x$  and  $y$  are rational numbers. The conjugate of  $\alpha$  is the element  $\bar{\alpha} = x - y\sqrt{D_0}$  of  $K$ . The norm of  $\alpha$  is the rational number  $N(\alpha) = \alpha\bar{\alpha} = x^2 - D_0y^2$ . We define the integer  $\omega_0$  of  $K$  by

$$(2.1) \quad \omega_0 = \begin{cases} \frac{\sqrt{D_0}}{2}, & \text{if } D_0 \equiv 0 \pmod{4}, \\ \frac{1}{2}(1 + \sqrt{D_0}), & \text{if } D_0 \equiv 1 \pmod{4}. \end{cases}$$

The ring of integers of  $K$  is  $O_{D_0} = [1, \omega_0]$ . For a positive integer  $f$ , we set

$$(2.2) \quad D = D_0f^2, \omega = \begin{cases} \frac{\sqrt{D}}{2}, & \text{if } D \equiv 0 \pmod{4} \\ \frac{1}{2}(1 + \sqrt{D}), & \text{if } D \equiv 1 \pmod{4}. \end{cases}$$

and

$$(2.3) \quad O_D = [1, \omega] = [1, f\omega_0].$$

It is easy to check that  $O_D$  is the subring of index  $f$  in  $O_{D_0}$ , called the order of discriminant  $D$ . We note that

$$(2.4) \quad \omega^2 = \begin{cases} \frac{D}{4}, & \text{if } D \equiv 0 \pmod{4}, \\ \omega + \frac{(D-1)}{4}, & \text{if } D \equiv 1 \pmod{4}. \end{cases}$$