Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 36 (1990)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: STATE MODELS FOR LINK POLYNOMIALS
Autor: Kauffman, Louis H.
Kapitel: IV. A Skein Model for the Kauffman polynomial
DOI: https://doi.org/10.5169/seals-57900

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 24.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

is admissible and contributes $(-z)^3\delta$ to the summation for the negative trefoil



IV. A Skein Model for the Kauffman polynomial

The work of section 3 goes over essentially verbatim for the Dubrovnik version of the Kauffman polynomial. Recall that in this context a template is obtained by first orienting the edges of the universe U underlying the unoriented K, and then labelling the edges of U.

The chart in Figure 7 shows the cases of admissible splices at crossings (with respect to the skein template algorithm). Each splice has been labelled with its corresponding vertex weight. Note that a splice is admissible if it indicates the form of passage that is obtained from an approach to the crossing that meets it as an under-crossing. Such approaches give active crossings in the skein template algorithm.

I have retained only the arrow for the first passage after each split, because the orientation on the other edge may change under the direction of the template. *The crossings are oriented because each end-node (unlink) produced by the skein template algorithm acquires an orientation from the directions of travel given by the template.*



Figure 7

In order to understand the pattern of these admissible splices, consider an unoriented crossing



I have labelled two of its regions A. These are swept out by turning the overcrossing line counterclockwise. The other two regions are labelled B. The *A-splice* of the crossing is that splice that joins the two regions labelled A. The *B-splice* joins the two regions labelled B.

We then see that a passage is admissible in the A-splice if it occurs on the *right* for an observer who stands in between the strands, facing in the direction of the passage from basepoint. Similarly, the admissible B-splices are on the left for such an observer.

Call an admissible splice *negative* if it is of B-type. (This receives a $(-z)$ in Figure 7.)

With these definitions we have

$$t(L) = \text{number of splices to obtain } L \text{ from } K\,.$$
$$t_-(L) = \text{number of negative splices}\,.$$

$\mathrm{Au}(K, T)$ = set of admissible unlinks relative to K and T (here K is unoriented).

Then the Dubrovnik polynomial is given by the formula

$$D_K = \sum_{L \in \mathrm{Au}(K,\,T)} (-1)^{t_-(L)} z^{t(L)} a^{w(L)} \mu^{|L|-1}$$
$$\left(\mu = 1 + (a - a^{-1})/z\right).$$

As a state-expansion we can write



Once again,

$$D_K = \sum_{L \in \mathrm{Au}(K,\,T)} \langle K \mid L \rangle\, \mu^{|L|-1}$$

where $<K \mid L>$ denotes the product of vertex weights (all relative to the given template). Independence of the template follows from the well-definedness of the polynomial itself.

Remark. It would be very interesting to know the relationship between this state model for the Kauffman polynomial and the extraordinary model of Jaeger [34]. Jaeger gives a state expansion where the states are a collection of oriented knots and links. Each state is itself evaluated via the regular isotopy version of the Homfly polynomial.

V. Graph polynomials

The two skein polynomials (Homfly and Kauffman) each have three variable extensions to rigid vertex isotopy invariants of 4-valent graphs imbedded in three-space. This construction has been announced in [45]. (See also [56] and [74].) Our skein models involve 4-valent graphs implicitly, and so give rise to a natural definition for these extended polynomials as state models.

Let the new variables A and B be given, with $z = A - B$ the usual z for the skein polynomials. The extended polynomials are then defined by the axioms:

Homfly Extension Axioms

1. R $= AR$ $+ R$,

   R $= BR$ $+ R$,

2. R_K = usual regular isotopy Homfly polynomial if K is free of graphical vertices ( ).

Kauffman Extension Axioms

1. D $= AD$ $+ BD$ $+ D$,

2. D_K = usual regular isotopy Dubrovnik polynomial if K is free of graphical vertices ( ).