

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 36 (1990)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: MANIN'S PROOF OF THE MORDELL CONJECTURE OVER
FUNCTION FIELDS
Autor: Coleman, Robert F.
Kapitel: 2. The Gauss-Manin connection
DOI: <https://doi.org/10.5169/seals-57915>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

since $\pi(s(v)) = v$ and $\pi \nabla(e) = \nabla_H(\pi(e))$. The lemma now follows from

$$(t \circ \nabla_G^\vee)(w)(e) = \nabla_G^\vee(w)(e - s(\pi(e))) = 0. \quad \square$$

Suppose W is an \mathcal{O}_S submodule of H . We let $[W]$ denote the smallest subconnection of H containing W .

2. THE GAUSS-MANIN CONNECTION

Here we will recall the definition and some basic properties of the Gauss-Manin connection which we will need in this paper. For more details see [K-O]. If \mathcal{S}^\bullet is a complex, $\mathcal{S}^\bullet(k)$ will denote the complex obtained from \mathcal{S}^\bullet by setting $\mathcal{S}^i(k) = \mathcal{S}^{i+k}$. For any scheme Y over K will let $K[Y]$ denote $\Gamma(\mathcal{O}_Y)$.

Suppose S is a smooth connected affine scheme over K . Suppose $f: X \rightarrow S$ is a smooth morphism, Z is a closed subscheme of X , smooth over S . Suppose T is either $\text{Spec}(K)$ or S . Then we define the subcomplex $\Omega_{X/T,Z}^\bullet$ of $\Omega_{X/T}^\bullet$ by the exactness of the sequence.

$$0 \rightarrow \Omega_{X/T,Z}^\bullet \rightarrow \Omega_{X/T}^\bullet \rightarrow \Omega_{Z/T}^\bullet \rightarrow 0.$$

When $T = \text{Spec}(K)$ we drop it from the notation. It follows that $\Omega_{X/S,Z}^i = \Omega_{X/S}^i$ for $i > \dim_S Z$. Note that $\Omega_{X,Z}^0 = \Omega_{X/S,Z}^0$ is the sheaf of ideals of Z on X . We define $H_{DR}^i(X/S, Z)$ to be the i -th hypercohomology group of the complex $\Omega_{X/S,Z}^\bullet$. We set $H_{DR}^i(X/S) = H_{DR}^i(X/S, \emptyset)$. If X is affine, then $H_{DR}^i(X/S, Z)$ is the i -th cohomology group of the complex of $K[S]$ modules $\Gamma(\Omega_{X/S,Z}^\bullet)$. If X is affine, K has characteristic zero and U is a dense open subscheme of X then the natural map from $H_{DR}^i(X/S, Z)$ to $H_{DR}^i(U/S, U \cap Z)$ is an injection.

From the last short exact sequence with $T = S$, we obtain a long exact sequence

$$(2.1) \quad \dots \rightarrow H_{DR}^{i-1}(Z/S) \rightarrow H_{DR}^i(X/S, Z) \rightarrow H_{DR}^i(X/S) \rightarrow \dots$$

The Gauss-Manin connection $\nabla: H_{DR}^i(X/S, Z) \rightarrow \Omega_S^1 \otimes H_{DR}^i(X/S, Z)$ is the boundary map in the long exact sequence obtained by taking hypercohomology of the short exact sequence of complexes:

$$(2.2) \quad 0 \rightarrow f^* \Omega_S^1 \otimes \Omega_{X/S,Z}^\bullet(-1) \rightarrow \Omega_{X/S,Z}^\bullet / f^* \Omega_S^2 \otimes \Omega_X^\bullet(-2) \rightarrow \Omega_{X/S,Z}^\bullet \rightarrow 0$$

(which is exact because X and Z are smooth over S). It is an integrable connection. If K has characteristic zero and f is surjective and has geometrically connected fibers, then $H_{DR}^0(X/S) = K[S]$ and the Gauss-Manin

connection is the trivial connection on this module. Moreover, it is easy to show that the sequence (2.1) is horizontal with respect to the respective Gauss-Manin connections.

Suppose now that S is an affine curve over K and $Z = \emptyset$. Then the short exact sequence (2.2) becomes

$$0 \rightarrow f^* \Omega_S^1 \otimes \Omega_{X/S}^1(-1) \rightarrow \Omega_{X/S}^1 \rightarrow \Omega_{X/S}^1 \rightarrow 0 .$$

Taking cohomology of this sequence yields the Leray long exact sequence

$$(2.3) \quad \dots \rightarrow H_{DR}^i(X/S) \xrightarrow{\nabla} \Omega_S^1 \otimes H_{DR}^i(X/S) \rightarrow H_{DR}^{i+1}(X) \rightarrow H_{DR}^{i+1}(X/S) \xrightarrow{\nabla} \dots$$

3. SECTIONS OF A FAMILY AND EXTENSIONS OF CONNECTIONS

Suppose now S is a smooth connected affine curve over a field K of characteristic zero and $f: X \rightarrow S$ is a smooth proper morphism of schemes over K , with geometrically connected fibers. These assumptions will be in force throughout the remainder of this paper. Suppose Z is a closed subscheme of X finite over S . Suppose the normalization $n: \tilde{Z} \rightarrow Z$ of Z is smooth over S . After repeated blowing ups at closed points we find a scheme $m: \tilde{X}' \rightarrow X$, which contains \tilde{Z} and is such that the restriction of m to \tilde{Z} is n . Let \tilde{X} equal the complement in \tilde{X}' of the singular locus of \tilde{X}'/S . This locus is a closed subscheme of \tilde{X}' disjoint from \tilde{Z} . The long exact sequence 2.1 becomes

$$(3.1) \quad 0 \rightarrow K[S] \rightarrow K[\tilde{Z}] \rightarrow H_{DR}^1(\tilde{Z}/S, \tilde{Z}) \rightarrow H_{DR}^1(\tilde{X}/S) \rightarrow 0$$

Let H denote the pullback of $H_{DR}^1(\tilde{X}/S, \tilde{Z})$ by means of the horizontal monomorphism from $H_{DR}^1(X/S)$ into $H_{DR}^1(\tilde{X}/S)$. We claim that H is independent of the choice of \tilde{X} . Indeed, there exists a non-empty affine open subscheme S' of S such that the map from $\tilde{X} \times_S S'$ to $X' = : X \times_S S'$ is an isomorphism. If $Z' = Z \times_S S'$, then Z' is smooth over S' and it is easy to see that $H \otimes K[S'] \cong H_{DR}^1(X'/S', Z')$. Hence H is an extension of the connection $H_{DR}^1(X'/S', Z')$ on S' to a connection on S . Since such an extension is unique if it exists, it follows that H is independent of the choice of \tilde{X} and so we set $H_{DR}^1(X/S, Z) = H$. We obtain from the previous exact sequence, a natural exact sequence

$$0 \rightarrow K[S] \rightarrow K[\tilde{Z}] \rightarrow H_{DR}^1(X/S, Z) \rightarrow H_{DR}^1(X/S) \rightarrow 0 .$$

For a section s of X/S , we will also use s to denote the induced reduced closed subscheme $s(S)$ of X when convenient. Now suppose s and t are two distinct sections of X/S . Let $Z = s \cup t$. Then \tilde{Z} , the normalization of Z , is just two disjoint copies of S and so is étale over S . (The sections s and t