

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 36 (1990)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: MANIN'S PROOF OF THE MORDELL CONJECTURE OVER
FUNCTION FIELDS
Autor: Coleman, Robert F.
Kapitel: 2. PICARD-FUCHS COMPUTATIONS
DOI: <https://doi.org/10.5169/seals-57915>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 06.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

2. PICARD-FUCHS COMPUTATIONS

We will need an explicit formula for $\mu(s, t)$ in some cases. Suppose that X/S has relative dimension one. Suppose $z \in K[S]$ such that $\Omega_S^1(S) = K[S]dz$ and suppose U is an affine open of $X, s \in U(S)$ and $v \in \mathcal{O}_X(U)$, such that $s^*v = 0$ and $\Omega_{X/S}^1(U) = \mathcal{O}_X(U)d_{X/S}v$. For $u \in \mathcal{O}_X(U)$ we define $\partial_z u$ and $\partial_v u$ by

$$du = \partial_z u dz + \partial_v u dv$$

Clearly ∂_z is a lifting of $\partial = : \partial / \partial z$ to a derivation of $\mathcal{O}_X(U)$. For $\omega = ud_{X/S}v \in \Omega_{X/S}^1(U)$ we set $\partial_z \omega = \partial_z u d_{X/S}v$ (the image of the Lie derivative of udv with respect to ∂_z in $\Omega_{X/S}^1(U)$). Since ∂ generates \mathcal{D} over $K[S]$ we can and will also make \mathcal{D} act on $\Omega_{X/S}^1(U)$ using ∂_z .

LEMMA 2.2.1. *Suppose $\omega = ud_{X/S}v \in \Omega_{X/S}^1(U)$ is of the second kind and $[\omega]$ is its class in $H_{DR}^1(X/S)$. Then*

$$\partial[\omega] = [\partial_z \omega] .$$

Proof. The element udv is a lifting of $ud_{X/S}v$ to $\Omega_X^1(U)$, and $d(udv) = du \wedge dv = \partial_z u dz \wedge dv$. Since this is the image of $dz \otimes \partial_z \omega$ in Ω_X^2 the lemma follows. \square

COROLLARY 2.2.2. *Suppose $\sum D_i \otimes \omega_i \in PF$. Then*

$$\sum D_i \omega_i = d_{X/S} w$$

for some $w \in \mathcal{O}_X(U)$.

Suppose $t \neq s$ is an element of $U(S)$ and $Z = s \cup t$. Let l denote the map from $K[S]$ into $H_{DR}^1(U/S, Z)$ associated to the pair (s, t) . For $\omega \in \Omega_{X/S}^1(U)$ let $[\omega]_Z$ denote the class of ω in $H_{DR}^1(U/S, Z)$.

LEMMA 2.2.3. *Suppose U, s and v are as above, $t \in U(S)$ and $t^*v \neq 0$. Suppose $\omega = ud_{X/S}v \in \Omega_{X/S}^1(U)$. Then $\partial^k[\omega]_Z$ equals*

$$[\partial_z^k \omega]_Z + l(\sum \partial^{i-1}(t^*(\partial_z^{k-i} u) \partial t^*v))$$

where i runs from 1 to k .

Proof. By shrinking S we may suppose that t^*v is invertible. We want to compute $\nabla[\omega]_Z$. First we must lift $ud_{X/S}v$ to section of $\Omega_{X,Z}^1(U)$. Let $y = f^*(t^*v)$. Then $\eta = u y dy^{-1} v$ is such a lifting and it equals $udv - u y y^{-1} \partial_z y dz$. Then $\nabla[\omega]_Z$ is the class of

$$d\eta = \partial_z u dz \wedge dv - d(uvy^{-1}) \wedge dy = dz \wedge \partial_z u dv + dz \wedge d(uvy^{-1} \partial_z y) .$$

which is the image of

$$dz \otimes (\partial_z \omega + d_{X/S}(uvy^{-1} \partial_z y)) \in \Omega_S^1 \otimes \Omega_{X/S}^1(U) .$$

Hence $\partial[\omega]$ is the class of $\partial_z \omega + d_{X/S}(uvy^{-1} \partial_z y)$ in $H_{DR}^1(U/S, Z)$. Since $(t^* - s^*)(uvy^{-1} \partial_z y) = t^* u \partial(t^* v)$ the lemma follows in the case $k = 1$. Since $\partial \circ l = l \circ \partial$ the lemma follows in general by induction. \square

COROLLARY 2.2.4. *Suppose U, s, z and v are as above, $t \in X(S)$ which meets U and $t^* v \neq 0$. Suppose ω, ω' and ω'' are elements $\omega_{X/S}$. Let $\omega = u d_{X/S} v$ and $\omega' = u' d_{X/S} v$ on U . Then we have:*

(i) *Suppose $\mu = \partial \otimes \omega - 1 \otimes \omega' \in PF$, $\omega = u d_{X/S} v$ and $\partial_z \omega - \omega' = d_{X/S} w$, with $w \in \mathcal{O}_X(U)$. Then*

$$\mu(s, t) = t^* w - s^* w + (t^* u) \partial t^* v .$$

(ii) *Suppose $\mu = \partial^2 \otimes \omega + \partial \otimes \omega' + 1 \otimes \omega'' \in PF$ and $\partial^2 \omega + \partial \omega' + \omega'' = d_{X/S} w$ with $w \in \mathcal{O}_X(U)$. Then*

$$\mu(s, t) = t^* ((w - s^* w, (u' + 2\partial_z u), \partial_v u, u) \cdot (1, x_t, x_t^2, \partial x_t))$$

and where $x_t = \partial t^* v$.

Proof. First shrink S so that s and t satisfy the hypotheses of the lemma and then apply it and the definition of $\mu(s, t)$. \square

Suppose $g: X \rightarrow A$ is a morphism over S from a curve to an Abelian scheme. Suppose $\kappa_{A/S}$ is an isomorphism. If $\eta = g^* \omega$ where $\omega \in \omega_{A/S}$ we will set $\mu_\eta = g^* \mu_\omega$. This is independent of the choice of ω . As an immediate consequence of the previous corollary we obtain:

COROLLARY 2.2.5. *Let U, z, s and v be as above. Set $X(S)' = \{t \in X(S) : t \text{ meets } U \text{ and } t^* v \neq 0\}$. Then there exist maps*

$$V = : V_{z, v} : T_{U, v} \rightarrow K(S)^4$$

and

$$L = : L_{z, v, s} : \omega_{X/S} \rightarrow K(X)^4$$

such that L is K -linear and for $t \in X(S)'$ and $\omega \in g^* \omega_{A/S}$,

$$\mu_\omega(s, t) = t^*(L(\omega) \cdot V(t)) .$$